

## **SUMMARY OF PRODUCT CHARACTERISTICS**

### **1 NAME OF THE MEDICINAL PRODUCT**

Clonidine Hydrochloride 25 mcg Tablets

### **2 QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each tablet contains Clonidine Hydrochloride 25 mcg.

Excipients with known effect:

Each tablet contains 78.275 mg of lactose monohydrate

For the full list of excipients, see section 6.1.

### **3 PHARMACEUTICAL FORM**

White to off white, round uncoated biconvex tablet, both side plain with approximately 6.0 mm diameter

### **4 CLINICAL PARTICULARS**

#### **4.1 Therapeutic indications**

- a) The prophylactic management of migraine or recurrent vascular headache.
- b) The management of vasomotor conditions commonly associated with the menopause and characterised by flushing.

#### **4.2 Posology and method of administration**

Posology

*Adults*

Initially 2 tablets twice daily. If after two weeks there has been no remission, increase to 3 tablets twice daily. The duration of treatment depends upon the severity of the condition.

If symptoms continue to occur the patient should be informed that it may take 2 - 4 weeks until Clonidine Hydrochloride 25 mcg Tablets are fully effective.

#### Paediatric population

There is insufficient information for the application of Clonidine in children and adolescents younger than 18 years. Therefore, the use of Clonidine is not recommended in paediatric subjects under 18 years.

#### Elderly

No specific information on the use of this product in the elderly is available. Clinical trials have included patients over 65 years and no adverse reactions specific to this age group have been reported.

#### Patient with Renal impairment

Clonidine Hydrochloride should be used with caution in patients with renal insufficiency. Careful monitoring of blood pressure is required.

#### Method of administration

For oral administration.

### **4.3 Contraindications**

Clonidine Hydrochloride 25 mcg Tablets should not be used in patients with severe bradyarrhythmia resulting from either sick-sinus syndrome or AV block of 2nd or 3rd degree, or in patients with known hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

In case of rare hereditary conditions that may be incompatible with an excipient of the product (please refer to section 4.4 Special warnings and precautions for use) the use of this product is contraindicated.

### **4.4 Special warnings and precautions for use**

Clonidine Hydrochloride 25 mcg Tablets should be used with caution in patients with cerebrovascular disease, coronary insufficiency, heart failure, occlusive peripheral vascular disorders, such as Raynaud's disease, polyneuropathy, constipation or those with a history of depression.

At doses higher than those recommended above, Clonidine is an effective antihypertensive agent. Caution should therefore be observed where antihypertensive agents are being used, as potentiation of the hypotensive effect may occur. Provided the recommended clonidine dosage regimen is followed, no difficulty with hypotension should arise during the routine management of patients with either migraine or menopausal flushing.

Depending on the dose given, Clonidine Hydrochloride can cause bradycardia. In patients with pre-existing cardiac conduction abnormalities, arrhythmias have been observed after high doses of clonidine hydrochloride.

Patients with renal failure require extreme care. (See Section 4.2).

Patients should be instructed not to discontinue therapy without consulting their physician. Following sudden discontinuation of Clonidine after prolonged treatment with high doses, agitation, restlessness, palpitations, rapid rise in blood pressure, nervousness, tremor, headache or nausea have been reported. When discontinuing therapy with Clonidine, the physician should reduce the dose gradually over 2-4 days.

Patients who wear contact lenses should be warned that treatment with clonidine may cause decreased lacrimation.

The use and the safety of clonidine in children and adolescents under 18 years have insufficient evidence in randomized controlled trials and therefore cannot be recommended for use in this population.

Serious adverse events, including sudden death, have been reported in concomitant use with methylphenidate. The safety of using methylphenidate in combination with clonidine has not been systematically evaluated.

### **Excipients**

This medicine contains lactose. Patients with rare hereditary problems of galactose intolerance, total lactase deficiency or glucose-galactose malabsorption should not take this medicine.

## **4.5 Interaction with other medicinal products and other forms of interaction**

Concurrent administration of antihypertensive agents, vasodilators or diuretics may lead to an increased hypotensive effect.

Substances with alpha2-receptor blocking properties, such as mirtazapine, may abolish the alpha2-receptor mediated effects of clonidine in a dose-dependent manner.

Concomitant use of beta-blockers and/or cardiac glycosides can cause bradycardia or dysrhythmia (AV-block) in isolated cases.

It cannot be ruled out that concomitant administration of a beta-receptor blocker will cause or potentiate peripheral vascular disorders.

If during combined treatment with a beta-blocker there is a need to interrupt or discontinue antihypertensive therapy, the beta-blocker must always be discontinued slowly first (reducing the dose gradually to avoid sympathetic hyperactivity), and then the Clonidine hydrochloride 25 mcg Tablets, which should also be reduced gradually over several days if previously given in high doses.

Orthostatic hypotension may be provoked or aggravated by concomitant administration of tricyclic antidepressants or neuroleptics with alpha-receptor blocking properties.

As the effects of Clonidine can be antagonised by tricyclic anti-depressants, it may be necessary to adjust the dosage of Clonidine Hydrochloride 25 mcg Tablets, if these agents are administered concurrently.

Although there is no experience from clinical trials, the effect of tranquillisers, hypnotics or alcohol could theoretically be potentiated by Clonidine Hydrochloride 25 mcg Tablets.

## **4.6 Fertility, pregnancy and lactation**

### **Pregnancy**

There is limited amount of data from the use of Clonidine in pregnant women. As with all medicines, Clonidine should not be used in pregnancy, especially the first trimester, unless the expected benefit is thought to outweigh any possible risk to the foetus.

In animal studies involving doses higher than the equivalent maximum therapeutic dose in man, effects on foetal development were only seen in one species. Foetal malformations did not occur.

Careful monitoring of mother and child is recommended.

Clonidine passes the placental barrier and may lower the heart rate of the foetus. Post partum a transient rise in blood pressure in the newborn cannot be excluded.

There is no adequate experience regarding the long-term effects of prenatal exposure.

### **Breast-feeding**

Clonidine is excreted in human milk.

However, there is insufficient information on the effect on newborns. The use of Clonidine is therefore not recommended during breast feeding.

### **Fertility**

No clinical studies on the effect on human fertility have been conducted with Clonidine. Non-clinical studies with clonidine indicate no direct or indirect harmful effects with respect to the fertility index.

#### **4.7 Effects on ability to drive and use machines**

No studies on the effects on the ability to drive and use machines have been performed. However, patients should be advised that they may experience undesirable effects such as dizziness, sedation and accommodation disorder during treatment with clonidine. If patients experience the above mentioned side effects they should avoid potentially hazardous tasks such as driving or operating machinery.

#### **4.8 Undesirable effects**

Most adverse effects are mild and tend to diminish with continued therapy.

Adverse events have been ranked under headings of frequency using the following convention: very common ( $\geq 1/10$ ), common ( $\geq 1/100, < 1/10$ ), uncommon ( $\geq 1/1,000, < 1/100$ ), rare ( $\geq 1/10,000, < 1/1,000$ ), very rare ( $< 1/10,000$ ) and unknown (cannot be calculated from the available data).

##### ***Cardiac disorders:***

Bradyarrhythmia: not known

Sinus bradycardia: uncommon

AV-block: rare

##### ***Nervous system disorders:***

Dizziness, Sedation: very common

Headache: common

Paraesthesia: uncommon

##### ***Eye disorders:***

Lacrimation decreased: rare.

Accommodation disorder: not known

##### ***Respiratory, thoracic and mediastinal disorders:***

Nasal dryness: rare

##### ***Gastrointestinal disorders:***

Dry mouth: very common

Constipation, Nausea, Salivary gland pain, Vomiting: common

Colonic pseudo-obstruction: rare

***Skin and subcutaneous tissue disorders:***

Pruritus, Rash, Urticaria: uncommon

Alopecia: rare

***Vascular disorders:***

Orthostatic hypotension: very common

Raynaud's phenomenon: uncommon

General disorders and administration site conditions:

Fatigue common Malaise: uncommon

***Reproductive system and breast disorders:***

Erectile dysfunction: common.

***Endocrine disorders:***

Gynaecomastia: rare

***Psychiatric disorders:***

Sleep disorder, depression: common

Hallucination, delusional perception, nightmare: uncommon

Confusional state, libido decreased: not known

***Investigations:***

Blood glucose increased: rare

**Reporting of suspected adverse reactions:**

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via Yellow Card Scheme; website: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store.

## **4.9 Overdose**

**Symptoms**

Manifestations of intoxication are due to a generalised sympathetic depression and include pupillary constriction, somnolence including coma, hypotension, orthostatic hypotension, bradycardia, hypothermia, respiratory depression including apnoea, occasionally vomiting, very occasionally hypertension, dryness of the mouth.

**Management**

There is no specific antidote for clonidine overdose. Administration of activated charcoal should be performed where appropriate.

Supportive care may include atropine sulfate for symptomatic bradycardia, and intravenous fluids and/or inotropic sympathomimetic agents for hypotension. Severe persistent hypertension may require correction with alpha- adrenoceptor blocking drugs.

Naloxone may be a useful adjunct for the management of clonidine-induced respiratory depression.

## **5 PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Antimigraine preparations ATC code: N02C X02

#### **Mechanism of action**

Clonidine is an antihypertensive agent which acts centrally by stimulating alpha<sub>2</sub>-adrenergic receptors and producing a reduction in sympathetic tone, resulting in a fall in diastolic and systolic blood pressure and a reduction in heart rate.

Treatment with Clonidine Hydrochloride 25 mcg Tablets diminishes the responsiveness of peripheral vessels to constrictor and dilator stimuli, thereby preventing the vascular changes associated with migraine. The same direct action on peripheral vessels moderates the vascular changes associated with menopausal flushing.

#### **Paediatric population**

The efficacy of Clonidine in the treatment of hypertension has been investigated in five clinical studies in paediatric patients. The efficacy data confirms the properties of Clonidine in reduction of systolic and diastolic blood pressure. However, due to limited data and methodological insufficiencies, no definitive conclusion can be drawn on the use of Clonidine for hypertensive children.

The efficacy of Clonidine has also been investigated in a few clinical studies with paediatric patients with ADHD, Tourette syndrome and stuttering. The efficacy of Clonidine in these conditions has not been demonstrated.

There were also two small paediatric studies in migraine, neither of which demonstrated efficacy.

In the paediatric studies the most frequent adverse events were drowsiness, dry mouth, headache, dizziness and insomnia. These adverse events might have serious impact on daily functioning in paediatric patients.

Overall, the safety and efficacy of Clonidine in children and adolescents have not been established. (see section 4.2).

## **5.2 Pharmacokinetic properties**

### **Absorption and distribution**

The pharmacokinetics of clonidine is dose-proportional in the range of 75-300 micrograms; over this range, dose linearity has not been fully demonstrated. Clonidine, the active ingredient of Clonidine Tablets, is highly absorbed and undergoes a minor first pass effect.

Peak plasma concentrations are reached within 1-3 h after oral administration. The plasma protein binding is 30-40 %. Clonidine is rapidly and extensively distributed into tissues and crosses the blood-brain barrier, as well as the placental barrier. Clonidine is excreted in human milk. However, there is insufficient information on the effect on newborns.

### **Biotransformation and elimination**

The terminal elimination half-life of clonidine has been found to range from 5 to 25.5 hours. It can be prolonged in patients with severely impaired renal function up to 41 hours.

About 70 % of the dose administered is excreted with the urine mainly in form of the unchanged parent drug (40-60 % of the dose). The main metabolite p-hydroxy-clonidine is pharmacologically inactive. Approximately 20% of the total amount is excreted with the faeces. There is no definitive data about food or race effects on the pharmacokinetics of clonidine.

The antihypertensive effect is reached at plasma concentrations between about 0.2 and 2.0 ng/ml in patients with normal renal function. The hypotensive effect is attenuated or decreases with plasma concentrations above 2.0 ng/ml.

## **5.3 Preclinical safety data**

There are no preclinical data of relevance to the prescriber which are additional to that already included in other sections of the SmPC.

# **6 PHARMACEUTICAL PARTICULARS**

## **6.1 List of excipients**

Lactose Monohydrate, Povidone, Maize Starch, Talc, Magnesium stearate.

## **6.2 Incompatibilities**

Not applicable

## **6.3 Shelf life**

36 Months

## **6.4 Special precautions for storage**

This medicinal product does not require any special storage conditions.

## **6.5 Nature and contents of container**

PVC/PVdC/Aluminium foil blister packs, with blister packs contained in a carton together with the patient information leaflet.

Clonidine Hydrochloride 25 mcg Tablets are available in pack of 30,90 and 112 Tablets.

Not all pack sizes may be marketed.

## **6.6 Special precautions for disposal**

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

## **7 MARKETING AUTHORISATION HOLDER**

Orbit Pharma Limited  
128 City Road, London,  
United Kingdom, EC1V 2NX

## **8 MARKETING AUTHORISATION NUMBER(S)**

PL 56854/0016

**9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

01/10/2024

**10 DATE OF REVISION OF THE TEXT**

20/05/2025