

**FONTS USED**

HelveticaNeueLTStd-Bd  
HelveticaNeueLTStd-Bk  
HelveticaNeueLTStd-It  
HelveticaNeueLTStd-Roman  
HelveticaWorld-Regular  
Wingdings-Regular

**SMALLEST TEXT SIZE**

6.00 pt

**AVE. TEXT SIZE**

8 pt

**LARGEST TEXT SIZE**

14.00 pt

# Imodium®

## Original <sup>2mg Capsules</sup> Loperamide hydrochloride

- This medicine is used for two different types of diarrhoea. They have **different age limits**. See [Section 1](#) ▶
- **Do not take this medicine:**
  - There are some **people who should not use** this medicine. *To find out if you are one of them, see [Section 2](#)* ▶
  - If you have ever had a **bad reaction** to any of the ingredients. *For the list of ingredients, see [Section 6](#)* ▶
- **Speak to your doctor:**
  - If you suffer from any of the conditions mentioned in [Section 2](#) ▶
  - If you are taking any **other medicines**. See [Section 2](#) ▶
- **If you have Irritable Bowel Syndrome (IBS)** see also [Section 2](#) Extra warnings for IBS patients ▶
- **Follow the dosage instructions carefully.** See [Section 3](#) ▶  
Always take this medicine exactly as described in this leaflet or as your doctor or pharmacist has told you.

Now read this whole leaflet carefully before you use this medicine. Keep the leaflet: you might need it again.

## 1 What the medicine is for

Imodium® Original 2mg Capsules is used to treat **two types of diarrhoea**. The two types have different age limits.

### **I** Short-term diarrhoea

- For adults and children aged **12 and over**.
- To treat attacks that last up to **48 hours**.
- If your attack lasts longer than 48 hours, **talk to your doctor**.

### **I** IBS diarrhoea

- For adults and young people aged **18 and over** who have been diagnosed with IBS (Irritable Bowel Syndrome).
- To treat attacks that last up to **48 hours**.
- You can use this medicine **for up to 2 weeks** for repeated attacks, but if any one attack lasts continuously for **longer than 48 hours, talk to your doctor**.

The capsules contain *loperamide hydrochloride*, a substance that helps reduce diarrhoea by slowing down an overactive bowel. This allows water and salts that are usually lost in diarrhoea to be absorbed by the body.

## 2 Before taking this medicine

### **I** Warnings for everyone

This medicine is suitable for most people, but a few people should not use it:

### **X** Do not take this medicine...

- If you have ever had a **bad reaction** to any of the ingredients.
  - If it is for a child aged **under 12** (or under 18 for an IBS patient).
  - If you have **severe diarrhoea** after taking **antibiotics**.
  - If you are having a flare-up of an **inflammatory bowel condition** like **ulcerative colitis**.
  - If you are **constipated**, or your **stomach appears swollen** (especially in children with severe dehydration).
  - If you have **acute dysentery**, the symptoms of which may include **blood in your stools** and a **high temperature**.
- If any of these applies to you, **talk to a doctor or pharmacist and do not take IMODIUM®**.

*This section is continued over the page.*

Turn over ▶

## **i** Warnings for everyone (continued)

### **A** Talk to your doctor first...

- If you have **AIDS** and your **stomach becomes swollen**, stop taking the capsules immediately and contact your doctor.
- If you suffer from **liver disease**.
- If you have diarrhoea that lasts for **more than 48 hours**.
- If you have been told by your doctor that you have an **intolerance to some sugars**.
- If you have **severe diarrhoea** as your body loses more fluid, sugars and salts than normal.
- If you are taking any **other medicines**, including:
  - *ritonavir* (used to treat **HIV**).
  - *quinidine* (used to treat **abnormal heart rhythms** or **malaria**).
  - *oral desmopressin* (used to treat **excessive urination**).
  - *itraconazole* or *ketoconazole* (used to treat **fungal infections**).
  - *gemfibrozil* (used to treat **high cholesterol**).

If you are unsure about any of the medicines you are taking, show the bottle or pack to your pharmacist.

If any of these applies to you (now or in the past), **talk to a doctor or pharmacist**.

### **Pregnancy or breast-feeding**

- **If you are pregnant**, think you are pregnant or planning a pregnancy: ask your doctor or pharmacist for advice before taking this medicine.
- **If you are breast-feeding** do not take this medicine.

Small amounts may get into your milk. Talk to your doctor about a suitable treatment.

### **Driving and using machines**

- This medicine may make you feel dizzy, tired or sleepy. You may feel less alert, feel faint or pass out. If you're affected do not drive, cycle or use machines.

### **Special warnings about this medicine**

- Your body can lose large amounts of fluids and salts when you have diarrhoea. You need to replace the fluid by drinking more liquid than usual. Ask your pharmacist about **rehydration therapy** to replace lost salts. **This is especially important for children, and frail or older people.**

- Consult with a doctor if you have a history of drug abuse.

- Loperamide is a type of opioid that acts directly on the gut when used as recommended but can cause other opioid effects when misused at high doses. Addiction is observed with opioids as a class.

- Abuse and misuse of loperamide, the active ingredient in IMODIUM® Original 2mg Capsules, have been reported. Do not take this product for anything other than its intended use (see section 1) and never take more than the recommended amount (see section 3).

- Serious heart problems (symptoms of which include fast or irregular heartbeat) have been reported in patients who have taken too much loperamide, the active ingredient in IMODIUM® Original 2mg Capsules.

### **Some of the ingredients can cause problems**

- This medicine contains lactose. If you have been told by your doctor that you have an intolerance to some sugars contact your doctor before taking this medicine.
- This medicine contains less than 1 mmol sodium (23 mg) per capsule, that is to say essentially 'sodium-free'.

### **i** Extra warnings for IBS patients

Use only if your doctor has previously diagnosed IBS. Check the following:

### **X** Do not take this medicine...

- If you are aged **under 18**.

### **A** Talk to your doctor first...

- If you are aged **40 or over** and it is **some time since your last IBS attack**.
- If you are aged **40 or over** and your **IBS symptoms are different** this time.
- If you have recently **passed blood from the bowel**.
- If you suffer from **severe constipation**.
- If you are feeling **sick or vomiting**.
- If you have **lost your appetite or lost weight**.
- If you have **difficulty or pain passing urine**.
- If you have a **fever**.
- If you have **recently travelled abroad**.

If any of these applies to you, **talk to your doctor before taking IMODIUM®**.

### 3 How to take this medicine

Always take this medicine exactly as described in this leaflet or as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

**Check the tables below to see how much medicine to take.**

- Swallow the correct number of capsules whole with a drink of water. For oral use only.
- Do not use more than the dose shown in the tables.
- The capsules are **not for long-term treatment**.

#### Short-term diarrhoea

Age	Dose
<b>Adults and children aged 12 and over</b>	Take <b>two capsules</b> to start treatment. Take <b>one capsule</b> after each loose bowel movement.
■ Do not take for attacks lasting <b>longer than 48 hours</b> .	■ Do not take more than <b>six capsules</b> in a 24- hour period.
■ <i>Replace lost fluid by drinking more liquid than usual.</i>	■ Not for children aged under 12.

### i How long to take IMODIUM® for short-term diarrhoea

**You can use this medicine for up to 48 hours.**

If your attack lasts longer than 48 hours, stop taking IMODIUM® and **talk to your doctor**.

#### IBS diarrhoea

Age	Dose
<b>Adults aged 18 and over</b>	Take <b>two capsules</b> to start treatment. Take <b>one capsule</b> after each loose bowel movement (or as advised by your doctor).
■ You can use this medicine <b>for up to 2 weeks</b> for repeated attacks, but do not take for any one attack lasting <b>longer than 48 hours</b> .	■ Do not take more than <b>six capsules</b> in a 24-hour period.
■ <i>Replace lost fluid by drinking more liquid than usual.</i>	■ Not for children and young people aged under 18.

**Talk to your doctor and stop taking this medicine:**

- if you have been using this medicine **continuously for 48 hours**.
- if you develop new IBS symptoms.
- if your IBS symptoms get worse.
- if your IBS symptoms have not improved **after 2 weeks**.

### i How long to take IMODIUM® for IBS diarrhoea

**You can use this medicine for up to 2 weeks** for repeated attacks of IBS diarrhoea. But if any one attack lasts for **longer than 48 hours**, stop taking IMODIUM® and **talk to your doctor**.

**If anyone takes too much of this medicine**

If you have taken too many Imodium® Original 2mg Capsules, immediately contact a doctor or hospital for advice. Symptoms may include: increased heart rate, irregular heartbeat, changes to your heartbeat (these symptoms can have potentially serious, life-threatening consequences), muscle stiffness, uncoordinated movements, drowsiness, difficulty urinating, or weak breathing.

Children react more strongly to large amounts of Imodium® Original 2mg Capsules than adults. If a child takes too much or shows any of the above symptoms, call a doctor immediately.

**If you forget to take the medicine**

You should only take this medicine as you need it, following the dosage instructions above carefully.

If you forget to take a dose, take a dose after the next loose stool (bowel movement). **Do not** take a double dose.

Turn over ►



## 4 Possible side effects

IMODIUM® can have side effects, like all medicines, although these don't affect everyone and most are usually mild.

### Get medical help at once

- Rare:** (may affect up to 1 in 1,000 people)
- Allergic reactions including unexplained wheezing, shortness of breath, passing out or swelling of face and throat.
  - Skin rashes, which may be severe and include blistering or peeling skin.
  - Loss of consciousness or reduced level of consciousness (passing out, feeling faint or less alert), muscle stiffness and uncoordinated movements.
  - Not known (frequency cannot be estimated from the available data): Upper abdominal pain, abdominal pain that radiates to back, tenderness when touching the abdomen, fever, rapid pulse, nausea, vomiting, which may be symptoms of inflammation of the pancreas (acute pancreatitis).

If you get any of these, **stop using the medicine and get medical help at once**.

### Talk to a doctor as soon as possible

- Uncommon:** (may affect up to 1 in 100 people)
- Itchiness or hives.
  - Stomach pain or swollen stomach.

**Rare:** (may affect up to 1 in 1,000 people)

- Difficulties passing water.
- Severe constipation.
- Miosis (narrowing of the pupils of the eye).

- Hypertonia (muscle tension).
- Coordination abnormality (uncoordinated movements).

If you notice any of the above, stop using the medicine and **talk to a doctor**.

### Other effects that may occur

**Common:** (may affect up to 1 in 10 people)

- Feeling sick, constipation or wind.
- Headache.

**Uncommon:** (may affect up to 1 in 100 people)

- Dizziness or drowsiness.
- Vomiting, indigestion.
- Dry mouth.

**Rare:** (may affect up to 1 in 1,000 people)

- Tiredness.

### Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

## 5 Storing this medicine

Keep this medicine out of the sight and reach of children.

Store in the original package.

Do not use your medicine after the date shown as the expiry date on the packaging.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

## 6 Further information

### What's in this medicine

**The active ingredient in Imodium® Original 2mg**

**Capsules is:** Loperamide hydrochloride 2 mg per capsule.

**Other ingredients are:** Lactose, maize starch, talc and magnesium stearate (E572). The capsule itself is made up of titanium dioxide (E171), black ferrous oxide (E172), yellow ferric oxide (E172), indigo carmine (E132), erythrosine (E127) and gelatin.

### What the medicine looks like

Imodium® Original 2mg Capsules are green / grey capsules marked 'Imodium' and 'JANSSEN'. They are available in packs of 2, 4 or 6 capsules. Not all pack sizes may be marketed.

**Product Licence holder:** McNeil Products Ltd, 50 - 100 Holmers Farm Way, High Wycombe, Buckinghamshire, HP12 4EG, UK.

**Manufacturer:** JNTL Consumer Health (France) SAS, Domaine de Maigremont, 27100 Val de Reuil, France.

**This leaflet was revised September 2024.**

IMODIUM® is a registered trade mark.

© **McNeil Products Limited 2024**