

Pentasa® 1g Suppositories

(mesalazine)

1950
26.09.25[12]

PATIENT INFORMATION LEAFLET

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

Your medicine is available using the above name but will be referred to as Pentasa Suppository throughout this leaflet.

What is in this leaflet:

1. What Pentasa Suppository is and what it is used for
2. What you need to know before you use Pentasa Suppository
3. How to use Pentasa Suppository
4. Possible side effects
5. How to store Pentasa Suppository
6. Contents of the pack and other information

1. WHAT PENTASA SUPPOSITORY IS AND WHAT IT IS USED FOR

Each Pentasa Suppository contains mesalazine 1g as the active ingredient. Mesalazine belongs to a group of medicines called salicylates.

Pentasa Suppository is used to treat inflammation in the gut caused by conditions such as ulcerative colitis. Sometimes ulcerative colitis only affects the back passage (rectum), this is known as ulcerative proctitis. Pentasa Suppositories are used to treat ulcerative proctitis and are intended for rectal use only.

The suppositories release the active ingredient slowly which then acts locally to reduce the inflammation and help relieve or stop the pain.

2. What you need to know before you use PENTASA Suppository

Do Not use Pentasa Suppository if you:

- are **allergic (hypersensitive) to mesalazine** or any of the other ingredients of this medicine (see section 6)
- are **allergic to other salicylates e.g. acetylsalicylic acid**
- have **severe liver and/or kidney** problems

Warnings and precautions

Talk to your doctor or pharmacist before using these suppositories if you:

- are allergic to sulphasalazine (risk of allergy to salicylates)
- currently have or have previously had liver or kidney disease
- have a medical condition that can make you prone to bleeding
- are on medication that may affect kidney function e.g. non-steroidal anti-inflammatory drugs (NSAIDs) such as aspirin
- have lung problems, in particular asthma
- suddenly develop abdominal cramps, abdominal pain, fever, severe headache and rash. In such circumstances you should stop taking Pentasa Suppository immediately.
- Kidney stones may develop with use of mesalazine. Symptoms may include pain in sides of abdomen and blood in urine. Take care to drink sufficient amount of liquid during treatment with mesalazine
- have ever developed a severe skin rash or skin peeling, blistering and/or mouth sores after using mesalazine.

Mesalazine may produce red-brown urine discoloration after contact with sodium hypochlorite bleach in the toilet water. It concerns a chemical reaction between mesalazine and bleach and is harmless.

Take special care with mesalazine:

Serious skin reactions including Drug reaction with eosinophilia and systemic symptoms (DRESS), Stevens-Johnson syndrome (SJS) and toxic epidermal necrolysis (TEN) have been reported in association with mesalazine treatment. Stop using Pentasa Suppository and seek medical attention immediately if you notice any of the symptoms related to these serious skin reactions described in section 4.

If you experience strong or recurrent headache, disturbed vision, or ringing or buzzing in the ears contact your doctor immediately.

While you are on treatment with this medicine, your doctor will normally arrange blood and urine tests to check your kidney function especially at the beginning of treatment.

You should make sure that you don't become dehydrated, while you are using this medicine. This can occur after severe or prolonged attacks of vomiting and/or diarrhoea, high fever or heavy sweating. If this occurs, you should talk to your doctor or a pharmacist for advice as soon as possible.

Other medicines and Pentasa Suppository

Tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription.

This is especially important if you are taking any of the following:

- **azathioprine** (used after transplantations or to treat autoimmune diseases)
- **6-mercaptopurine or thioguanine** (chemotherapy, used to treat leukaemia)
- certain agents that inhibit blood clotting (medicines for thrombosis or to thin your blood, e.g. warfarin)

Pregnancy, breast-feeding and fertility

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before using this medicine.

There is limited experience with the use of mesalazine during pregnancy and breast-feeding. Blood disorders have been reported in newborns of mothers being treated with this medicine. The newborn may develop allergic reactions after breast-feeding, e.g. diarrhoea. If the newborn develops diarrhoea, breast-feeding should be discontinued.

Driving and using machines

This medicine is not known to affect the ability to drive and/or use machines.

3. HOW TO USE PENTASA SUPPOSITORY

For Adult use only, not recommended in children.

Always use Pentasa Suppository exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure. The suppository should be inserted into the back passage (rectum) only. It is intended to be left in place for as long as possible and is therefore best inserted before going to sleep.

To treat an attack of proctitis your doctor will usually prescribe one suppository to be used daily for two to four weeks. To help prevent further attacks it may be used for longer.

How to insert the suppository

You should make sure that you empty your bowels before inserting the suppository.

- a. Tear along the perforation to remove one suppository blister from a blister strip
- b. Put a protector on the finger with which you will insert the suppository
- c. Remove the suppository from the blister before insertion. This is done by pushing the suppository through the printed foil
- d. Insertion may be made easier if you moisten the suppository with water first
- e. Insert the suppository gently and fully into the back passage. It may help to lie down on one side and raise the other knee.
- f. Remain still for a few minutes after insertion to help keep the suppository in place. If it comes out within 10 minutes, a new one should be inserted
- g. Dispose of the empty blister and used finger protector safely and wash your hands.

If you use more Pentasa Suppository than you should

If you accidentally use too many suppositories, you should go to your nearest emergency department or contact your doctor immediately. Take the pack and any remaining suppositories with you.

If you forget to use Pentasa Suppository

If you forget to use a dose, use the next dose as soon as you remember, unless it is less than 6 hours until your next dose. Do not use a double dose to make up for the forgotten one.

If you have any further questions on the use of this product, ask your doctor or pharmacist.

4. POSSIBLE SIDE EFFECTS

Like all medicines, Pentasa Suppository can cause side effects, although not everyone gets them. Following rectal administration local reactions such as itching, rectal discomfort and urge may occur.

STOP using Pentasa Suppository and seek medical attention immediately if you notice any of the following symptoms:

- reddish non-elevated, target-like or circular patches on the trunk, often with central blisters, skin peeling, ulcers of mouth, throat, nose, genitals and eyes, widespread rash, fever and enlarged lymph nodes. These serious skin rashes can be preceded by fever and flu-like symptoms.
- itching, skin rashes, swelling of the face, lips or throat, difficulty in breathing or wheeziness (signs of an allergic reaction).
- unexplained bleeding, bruising, skin rashes, fever or sore throat (signs of blood disorder).
- a change in the colour or amount of urine produced (signs of kidney problems).

If you experience any of the above side effects, you should contact your doctor or go to the nearest hospital emergency department immediately.

The following common side effects may affect up to 1 in 10 people:

- diarrhoea
- abdominal pain
- nausea
- vomiting
- headache
- rash
- flatulence (passing wind)
- rectal discomfort,
- irritation at the site of application

The following rare side effects may affect up to 1 in 1,000 people:

- inflammation of some areas of the heart (myocarditis and pericarditis) which can cause shortness of breath and chest pain or palpitations (rapid or irregular heart beats).
- inflammation of the pancreas (symptoms include back and/or stomach pain).
- increased amylase (enzyme that helps digest carbohydrates).
- dizziness.
- increased sensitivity of your skin to sun and ultraviolet light (photosensitivity).

The following very rare side effects may affect up to 1 in 10,000 people:

- anaemia and other blood disorders (decrease in the numbers of certain blood cells, which can cause unexplained bleeding, bruising, fever or sore throat)
- liver disorders (symptoms include jaundice (yellowing of the skin and/or eyes) and/or pale bowel motions)
- kidney disorders (symptoms include blood in the urine, and/or oedema (swelling due to build-up of fluid))
- peripheral neuropathy (a condition affecting the nerves of the hands and feet symptoms include tingling and numbness)
- allergic and fibrotic lung reactions, inflammation of the lining of the lungs or lung scarring (symptoms include coughing, bronchospasm, chest discomfort or pain on breathing, breathing difficulties, bloody and/or excessive phlegm)
- pancolitis (a kind of inflammatory bowel disorder (IBD) that affects the entire internal lining of the large bowel)
- hair loss (this is reversible)
- muscle or joint pain
- inflammation which can affect different parts of the body such as joints, skin, kidneys, heart etc. (symptoms include painful joints, fatigue, fever, abnormal or unexplained bleeding (e.g. nose bleeds), bruising, purple discoloration of the skin (including severe skin erosions and severe blistering that may affect the skin as the protective barrier of the body))
- semen with a low concentration of sperm (oligospermia) (this is reversible)
- severe diarrhoea and abdominal pain because of an allergic reaction to this medicine within the bowel
- allergic reactions and fever may occasionally occur.

Not known (frequency cannot be estimated from the available data)

- kidney stones and associated kidney pain (see also section 2)
- change in urine colour.
- if you experience strong or recurrent headache, disturbed vision, or ringing or buzzing in the ears. These could be symptoms of increased pressure within your skull (idiopathic intracranial hypertension).

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme Website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects, you can help provide more information on the safety of this medicine.

5. HOW TO STORE PENTASA SUPPOSITORY

Keep out of the sight and reach of children.

Do not store above 25°C.

Store this medicine in the original package.

Do not use after the expiry date marked on the packaging. The expiry date refers to the last day of that month.

If your suppositories become discoloured or show any other signs of deterioration, consult your doctor or pharmacist who will tell you what to do.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines that are no longer required. These measures will help to protect the environment.

6. CONTENTS OF THE PACK AND OTHER INFORMATION

What Pentasa Suppository contains

Each suppository contains 1g of the active ingredient, mesalazine (also known as 5-aminosalicylic acid or 5-ASA).

Other ingredients in this medicine are povidone, magnesium stearate, macrogol 6000 and talc.

What Pentasa Suppository looks like and the contents of the pack

Pentasa Suppositories are oval, compressed, white to pale brown, speckled suppositories.

They are presented in blister strips of 7 suppositories per strip.

Each carton contains 28 suppositories together with 28 finger protectors for use when inserting the suppository.

Manufacturer and product licence holder

Manufactured by Ferring GmbH, Wittland 11, D-24109 Kiel, Germany.

Procured from within the EU by product licence holder Tenolol Ltd, 5 Sandridge Close, Harrow, Middlesex, HA1 1XD. Repackaged by Servipharm Ltd.

POM

PL 30900/1950

Leaflet revision and issue (Ref) 26.09.25[12]

Pentasa is a trademark of Ferring B.V.

**Blind or partially sighted?
Is this leaflet hard to see or read?
Call 020 8423 2111 to obtain the
leaflet in a format suitable for you.**