
 <span style="float: right; font-size: 24px; font-weight: bold;">Research</span>	
<b>Technical information</b>	
Ref. Doc Source Bepharbel:	FP00000240_20251_NO_V0_CommentsYOU sent by Youness Ougnoun on 2025-06-26
Weight Paper:	40 g/m <sup>2</sup>
Artwork's Type:	Leaflet (PIL)
Product Name:	Amversio 1g
Customer :	SERB
Market:	GB
Supplier	UNIT
Opened Size:	150 x 280 mm
Folded Size:	53 x 150 mm
Artwork Reference Cutomer	FP00000240_20251_NO
Artwork Code Bepharbel:	010NO01
Technical Colors:	Cut = Crop marks Guide
Printing Colors:	Black
Laetus Code Bepharbel:	840
Spot Marks :	50 mm
Minimum Font Size:	9 Pt. Helvetica Neue TT Regular - Equivalent to Times New Roman 9 pt
<b>Revision History:</b>	
Date:	History:
22-JUL-2025	Creation
01-SEP-2025	Correction
10-SEP-2025	Finalization
06-OCT-2025	Correction
30-OCT-2025	Update Artwork code
<b>Artwork Creation:</b> Atwork Customer Approval	
Title: Artwork Product Designer First Name: Gilles Name: Renneson	Title: First Name: Name:
Date & Signature (or Visa)  30-OCT-2025  	Date & Signature (or Visa)

**APPROVED**

By BINDA Cindy at 5:12 pm, Oct 31, 2025

**APPROVED**

By Marion Gardin at 3:33 pm, Nov 11, 2025

**Package leaflet: Information for the patient**  
**Amversio 1 g oral powder**  
betaine anhydrous

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

**What is in this leaflet**

1. What Amversio is and what it is used for
2. What you need to know before you take Amversio
3. How to take Amversio
4. Possible side effects
5. How to store Amversio
6. Contents of the pack and other information

**1. What Amversio is and what it is used for**

Amversio contains betaine anhydrous which is intended to be an adjunctive treatment of homocystinuria, an inherited (genetic) disease where the amino acid methionine cannot be broken down completely by the body.

Methionine is present in regular food protein (such as meat, fish, milk, cheese, eggs). It is converted into homocysteine which is then normally converted into cysteine during digestion. Homocystinuria is a disease caused by the accumulation of homocysteine which is not converted to cysteine and is characterised by formation of clots in the veins, bone weakness, and skeletal and crystalline lens abnormalities. The use of Amversio together with other treatments such as vitamin B6, vitamin B12, folate and a specific diet aims to reduce the elevated homocysteine levels in your body.

**2. What you need to know before you take Amversio**

**Do not take Amversio**

- if you are allergic to betaine anhydrous.

**Warnings and precautions**

Talk to your doctor or pharmacist before taking Amversio.

- If you notice side effects like headaches, vomiting or a change in your vision and you are of the homocystinuria subtype called CBS (cystathionine beta-synthase deficiency), please contact your doctor immediately, they could be signs of a swelling in the brain (cerebral oedema). In that case your doctor will monitor your methionine level in your body and may review your diet. Your treatment with Amversio may need to be interrupted.
- If you are treated with Amversio and with an amino-acid mixture and if you need to take other medicines at the same time, leave 30 minutes between the intake (see "section "Other medicines and Amversio).

**Other medicines and Amversio**

Tell your doctor if you are taking, have recently taken or might take any other medicines.

If you are taking amino-acid mixture or medicines such as vigabatrin or Gaba analogues (medicine used to treat epilepsy), please tell your doctor as they might interact with your treatment with Amversio.

**Pregnancy and breast-feeding**

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will decide if you can use this medicine during pregnancy and breast-feeding.

**Driving and using machines**

Amversio has no or negligible influence on the ability to drive and use machines.

**3. How to take Amversio**

The use of this medicine will be supervised by a doctor experienced in the treatment of patients with homocystinuria.

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

The recommended dose in children and adults is 100 mg/kg/day divided in 2 doses per day. In some patients doses above 200 mg/kg/day were needed to reach therapeutic goals. Your doctor may adapt the dose depending on your laboratory values.

You will therefore need regular blood tests to determine the correct daily dose.

## Instructions for use

You should take Amversio orally (by mouth).

To measure the dose:

- shake the bottle lightly before opening
- take the correct measuring spoon:
  - the green measuring spoon dispenses 100 mg;
  - the blue measuring spoon dispenses 150 mg;
  - the purple measuring spoon dispenses 1 g.
- take a heaped spoonful of powder out of the bottle
- pass the flat back of a knife over the top of the spoon
- the powder left in the spoon is one spoonful
- take the correct number of spoonfuls of powder from the bottle

Mix the measured dose of powder with water, juice, milk, formula or food until completely dissolved and ingest immediately after mixing.

### If you take more Amversio than you should

If you accidentally take too much Amversio, talk to a doctor or pharmacist immediately.

### If you forget to take Amversio

Do not take a double dose to make up for forgotten doses. If you forget to take a dose take it as soon as you remember it and continue the next dose as planned.

### If you stop taking Amversio

Do not stop the treatment without consulting your doctor. Contact your doctor or pharmacist before stopping.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

## 4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Methionine level can be related to swelling in the brain (cerebral swelling), which may affect up to 1 in 100 people (frequency uncommon). If you experience morning headaches with vomiting and/or visual changes, **contact immediately your doctor (they could be signs of a swelling in the brain).**

Other side effects include:

Very common (may affect more than 1 in 10 people)

- high levels of methionine in the blood

Uncommon (may affect up to 1 in 100 people)

- gastrointestinal disorders like diarrhoea, nausea, vomiting, stomach discomfort and inflammation of the tongue,
- decreased appetite (anorexia),
- agitation,
- irritability,
- hair loss,
- hives,
- abnormal skin odour,
- lack of control over passing urine (urinary incontinence).

### Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

## 5. How to store Amversio

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the bottle label and the carton after EXP. The expiry date refers to the last day of that month.

Store in the original package.

Keep the bottle tightly closed in order to protect from moisture.

This medicine does not require any special temperature storage conditions.

After the first opening of the bottle, the medicine should be used within 3 months.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## 6. Contents of the pack and other information

### What Amversio contains

- The active substance is betaine anhydrous. 1 g of oral powder contains 1 g of betaine anhydrous.
- There is no other ingredient.

### What Amversio looks like and contents of the pack

Amversio is a white crystalline free flowing powder. It is presented in a bottle with a child resistant closure. Each bottle contains 180 g of powder. Each carton contains one bottle and three measuring spoons.

The green measuring spoon dispenses 100 mg.

The blue measuring spoon dispenses 150 mg.

The purple measuring spoon dispenses 1 g.

### Marketing Authorisation Holder

SERB S.A., Avenue Louise 480, 1050 Brussels, Belgium

This leaflet was last revised in November 2025.

### Manufacturer

BePharBel Manufacturing S.A.,  
rue du Luxembourg 13,  
6180 Courcelles, Belgium