

## **SUMMARY OF PRODUCT CHARACTERISTICS**

### **1 NAME OF THE MEDICINAL PRODUCT**

Lutonaze 137 micrograms /50 micrograms per actuation, Nasal Spray, Suspension

Azelastine hydrochloride/Fluticasone propionate Zentiva 137 micrograms /50 micrograms per actuation, Nasal Spray, Suspension

### **2 QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each gram of suspension contains 1000 micrograms azelastine hydrochloride and 365 micrograms fluticasone propionate.

One actuation (0.14 g) delivers 137 micrograms azelastine hydrochloride (equivalent to 125 micrograms azelastine) and 50 micrograms fluticasone propionate.

Excipient with known effect:

One actuation (0.14 g) delivers 0.014 mg benzalkonium chloride.

For the full list of excipients, see section 6.1.

### **3 PHARMACEUTICAL FORM**

Nasal spray, suspension. (Nasal spray)

White, homogeneous suspension.

### **4 CLINICAL PARTICULARS**

#### **4.1 Therapeutic indications**

Relief of symptoms of moderate to severe seasonal and perennial allergic rhinitis if monotherapy with either intranasal antihistamine or glucocorticoid is not considered sufficient.

## **4.2 Posology and method of administration**

### Posology

For full therapeutic benefit regular usage is essential.

Contact with the eyes should be avoided.

### *Adults and adolescents (12 years and older)*

One actuation in each nostril twice daily (morning and evening).

### *Children below 12 years*

Azelastrine hydrochloride/Fluticasone propionate Nasal Spray is not recommended for use in children below 12 years of age as safety and efficacy has not been established in this age group.

### *Elderly*

No dose adjustment is required in this population.

### *Renal and hepatic impairment*

There are no data in patients with renal and hepatic impairment.

### Duration of treatment

Azelastrine hydrochloride/Fluticasone propionate Nasal Spray is suitable for long-term use.

The duration of treatment should correspond to the period of allergenic exposure.

### Method of administration

Azelastrine hydrochloride/Fluticasone propionate Nasal Spray is for nasal use only.

### Instructions for use

#### *Preparing the spray:*

The bottle should be shaken gently before use for about 5 seconds by tilting it upwards and downwards and the protective cap be removed afterwards. Prior to first

use Azelastine hydrochloride /Fluticasone propionate Nasal Spray must be primed by pressing down and releasing the pump 6 times. If Azelastine hydrochloride/Fluticasone propionate Nasal Spray has not been used for more than 7 days it must be reprimed once by pressing down and releasing the pump.

*Using the spray:*

After blowing the nose the suspension is to be sprayed once into each nostril keeping the head tilted downward (see figure). After use the spray tip is to be wiped and the protective cap to be replaced.

### **4.3 Contraindications**

Hypersensitivity to the active substances or to any of the excipients listed in section 6.1.

### **4.4 Special warnings and precautions for use**

During post-marketing use, there have been reports of clinically significant drug interactions in patients receiving fluticasone propionate and ritonavir, resulting in systemic corticosteroid effects including Cushing's syndrome and adrenal suppression. Therefore, concomitant use of fluticasone propionate and ritonavir should be avoided, unless the potential benefit to the patient outweighs the risk of systemic corticosteroid side-effects (see section 4.5).

Systemic effects of nasal corticosteroids may occur, particularly when prescribed at high doses for prolonged periods. These effects are much less likely to occur than with oral corticosteroids and may vary in individual patients and between different corticosteroid preparations. Potential systemic effects may include Cushing's syndrome, Cushingoid features, adrenal suppression, growth retardation in children and adolescents, cataract, glaucoma and more rarely, a range of psychological or behavioural effects including psychomotor hyperactivity, sleep disorders, anxiety, depression or aggression (particularly in children).

Azelastine hydrochloride/Fluticasone propionate Nasal Spray undergoes extensive first-pass metabolism, therefore the systemic exposure of intranasal fluticasone propionate in patients with severe liver disease is likely to be increased. This may result in a higher frequency of systemic adverse events. Caution is advised when treating these patients.

Treatment with higher than recommended doses of nasal corticosteroids may result in clinically significant adrenal suppression. If there is evidence for higher than recommended doses being used, then additional systemic corticosteroid cover should be considered during periods of stress or elective surgery.

In general, the dose of intranasal fluticasone formulations should be reduced to the lowest dose at which effective control of the symptoms of rhinitis is maintained. Higher doses than the recommended one (see section 4.2) have not been tested for Azelastine/Fluticasone. As with all intranasal corticosteroids, the total systemic burden of corticosteroids should be considered whenever other forms of corticosteroid treatment are prescribed concurrently.

Growth retardation has been reported in children receiving nasal corticosteroids at licensed doses. Regular monitoring of height growth in children and adolescents, who are on long-term treatment with nasal corticosteroids are recommended. If growth is slowed, treatment should be reviewed aiming at reducing the dose of nasal corticosteroid, if possible, to the lowest dose at which effective control of symptoms is maintained.

Visual disturbance may be reported with systemic and topical corticosteroid use. If a patient presents with symptoms such as blurred vision or other visual disturbances, the patient should be considered for referral to an ophthalmologist for evaluation of possible causes which may include cataract, glaucoma or rare diseases such as central serous chorioretinopathy (CSCR) which have been reported after use of systemic and topical corticosteroids.

Close monitoring is required in patients with a change in vision or with a history of increased ocular pressure, glaucoma and/or cataracts.

If there is any reason to believe that adrenal function is impaired, care must be taken when transferring patients from systemic steroid treatment to Azelastine/Fluticasone Nasal Spray.

In patients who have tuberculosis, any type of untreated infection, or have had a recent surgical operation or injury to the nose or mouth, the possible benefits of the treatment with Azelastine/Fluticasone Nasal Spray should be weighed against possible risk.

Infections of the nasal airways should be treated with antibacterial or antimycotical therapy, but do not constitute a specific contraindication to treatment with Azelastine hydrochloride/Fluticasone propionate Nasal Spray.

Excipient(s) with known effect:

Azelastine hydrochloride/Fluticasone propionate Nasal Spray contains benzalkonium chloride. Long term use may cause oedema of the nasal mucosa. Each actuation (0.14 g) contains less than 1 mmol (23 mg) sodium, that is to say essentially “sodium-free”.

## 4.5 Interaction with other medicinal products and other forms of interaction

### *Fluticasone propionate*

Under normal circumstances, low plasma concentrations of fluticasone propionate are achieved after intranasal dosing, due to extensive first pass metabolism and high systemic clearance mediated by cytochrome P450 3A4 in the gut and liver. Hence, clinically significant drug interactions mediated by fluticasone propionate are unlikely.

A drug interaction study in healthy subjects has shown that ritonavir (a highly potent cytochrome P450 3A4 inhibitor) can greatly increase fluticasone propionate plasma concentrations, resulting in markedly reduced serum cortisol concentrations. During post-marketing use, there have been reports of clinically significant drug interactions in patients receiving intranasal or inhaled fluticasone propionate and ritonavir, resulting in systemic corticosteroid effects. Co-treatment with other CYP 3A4 inhibitors, including cobicistat-containing products is also expected to increase the risk of systemic side-effects. The combination should be avoided unless the benefit outweighs the increased risk of systemic corticosteroid side-effects, in which case patients should be monitored for systemic corticosteroid side-effects.

Studies have shown that other inhibitors of cytochrome P450 3A4 produce negligible (erythromycin) and minor (ketoconazole) increases in systemic exposure to fluticasone propionate without notable reductions in serum cortisol concentrations. Nevertheless, care is advised when co-administering potent cytochrome P450 3A4 inhibitors (e.g. ketoconazole), as there is potential for increased systemic exposure to fluticasone propionate.

### *Azelastine hydrochloride*

No specific interaction studies with azelastine hydrochloride nasal spray have been performed. Interaction studies at high oral doses have been performed. However, they bear no relevance to azelastine nasal spray as given recommended nasal doses result in much lower systemic exposure. Nevertheless, care should be taken when administering azelastine hydrochloride in patients taking concurrent sedative or central nervous medications because sedative effect may be enhanced. Alcohol may also enhance this effect (see section 4.7).

## 4.6 Fertility, pregnancy and lactation

### Fertility

There are only limited data with regard to fertility (see section 5.3).

### Pregnancy

There are no or limited amount of data from the use of azelastine hydrochloride and fluticasone propionate in pregnant women. Therefore, Azelastine/Fluticasone Nasal Spray should be used during pregnancy only if the potential benefit justifies the potential risk to the foetus (see section 5.3)

### Lactation

It is unknown whether nasally administered azelastine hydrochloride/metabolites or fluticasone propionate/metabolites are excreted in human breast milk.

Azelastine/Fluticasone Nasal Spray should be used during lactation only if the potential benefit justifies the potential risk to the newborns/infant (see section 5.3).

## **4.7 Effects on ability to drive and use machines**

Azelastine hydrochloride/Fluticasone propionate Nasal Spray has minor influence on the ability to drive and use machines. In isolated cases fatigue, weariness, exhaustion, dizziness or weakness that may also be caused by the disease itself, may occur when using Azelastine hydrochloride /Fluticasone propionate Nasal Spray. In these cases, the ability to drive and use machines may be impaired. Alcohol may enhance this effect.

## **4.8 Undesirable effects**

Commonly, dysgeusia, a substance-specific unpleasant taste, may be experienced after administration (often due to incorrect method of application, namely tilting the head too far backwards during administration).

Adverse reactions are listed below by system organ class and frequency. Frequencies are defined as:

Very common ( $\geq 1/10$ )

Common ( $\geq 1/100$  to  $< 1/10$ )

Uncommon ( $\geq 1/1,000$  to  $< 1/100$ )

Rare ( $\geq 1/10,000$  to  $< 1/1,000$ )

Very rare ( $< 1/10,000$ )

Not known (cannot be estimated from the available data)

<b>Frequency</b>	<b>Very common</b>	<b>Common</b>	<b>Uncommon</b>	<b>Rare</b>	<b>Very rare</b>	<b>Not known</b>
<b>System Organ Class</b>						

<b>Immune system disorders</b>					Hypersensitivity including anaphylactic reactions, angioedema (oedema of the face or tongue and skin rash), bronchospasm	
<b>Nervous system disorder</b>		Headache, Dysgeusia (unpleasant taste), unpleasant smell			Dizziness, somnolence (drowsiness, sleepiness)	
<b>Eye disorders*</b>					Glaucoma, increased intraocular pressure, cataract	Vision, blurred (see also section 4.4)
<b>Respiratory, thoracic and mediastinal disorders</b>	Epistaxis		Nasal discomfort (including nasal irritation, stinging, itching), sneezing, nasal dryness, cough, dry throat, throat irritation		Nasal septal perforation**, mucosal erosion	Nasal ulcers
<b>Gastrointestinal disorders</b>				Dry mouth	Nausea	
<b>Skin and subcutaneous tissue disorders</b>					Rash, Pruritus, Urticaria	
<b>General disorders and administration site conditions</b>					Fatigue (weariness, exhaustion), weakness (see section 4.7)	

\* A very small number of spontaneous reports have been identified following prolonged treatment with intranasal fluticasone propionate.

\*\* Nasal septal perforation has been reported following the use of intranasal corticosteroids.

Systemic effects of some nasal corticosteroids may occur, particularly when administered at high doses for prolonged periods (see section 4.4).

Growth retardation has been reported in children receiving nasal corticosteroids. Growth retardation may be possible in adolescents, too (see section 4.4).

In rare cases osteoporosis was observed if nasal glucocorticoids were administered long-term.

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store.

## **4.9 Overdose**

With the nasal route of administration overdose reactions are not anticipated.

There are no data from patients available on the effects of acute or chronic overdosage with intranasal fluticasone propionate.

Intranasal administration of 2 milligrams fluticasone propionate (10 times the recommended daily dose) twice daily for seven days to healthy human volunteers has no effect on hypothalamo-pituitary-adrenal (HPA) axis function.

Administration of doses higher than those recommended over a long period of time may lead to temporary suppression of adrenal function.

In these patients, treatment with Azelastine hydrochloride/Fluticasone propionate Nasal Spray should be continued at a dose sufficient to control symptoms; the adrenal function will recover in a few days and can be verified by measuring plasma cortisol.

In the event of overdose after incidental oral uptake, disturbances of the central nervous system (including drowsiness, confusion, coma, tachycardia and hypotension) caused by azelastine hydrochloride are to be expected based on the results of animal experiments.

Treatment of these disorders must be symptomatic. Depending on the amount swallowed, gastric lavage is recommended. There is no known antidote.

## 5 PHARMACOLOGICAL PROPERTIES

### 5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Decongestants and other nasal preparations for topical use, corticosteroids/ fluticasone, combinations, ATC code: R01AD58.

#### Mechanism of action and pharmacodynamic effects

Azelastine hydrochloride/Fluticasone propionate Nasal Spray contains azelastine hydrochloride and fluticasone propionate, which have different modes of action and show synergistic effects in terms of improvement of allergic rhinitis and rhinoconjunctivitis symptoms.

#### *Fluticasone propionate*

Fluticasone propionate is a synthetic trifluorinated corticosteroid that possesses a very high affinity for the glucocorticoid receptor and has a potent anti-inflammatory action, e.g. 3-5 fold more potent than dexamethasone in cloned human glucocorticoid receptor binding and gene expression assays.

#### *Azelastine hydrochloride*

Azelastine, a phthalazinone derivative is classified as a potent long-acting anti-allergic compound with selective H<sub>1</sub>-antagonist, mast cell stabilizing and anti-inflammatory properties. Data from *in vivo* (preclinical) and *in vitro* studies show that azelastine inhibits the synthesis or release of the chemical mediators known to be involved in early and late-stage allergic reactions, e.g. leukotrienes, histamine, platelet-activating factor (PAF) and serotonin.

Relief of nasal allergic symptoms is observed within 15 minutes after administration.

#### Azelastine hydrochloride/Fluticasone propionate nasal spray

In 4 clinical studies in adults and adolescents with allergic rhinitis azelastine/fluticasone nasal spray one spray in each nostril twice daily significantly improved nasal symptoms (comprising rhinorrhoea, nasal congestion, sneezing and nasal itching) compared with placebo, azelastine hydrochloride alone and fluticasone propionate alone. It significantly improved ocular symptoms (comprising itching, tearing/watering and redness of the eyes) and the patients' disease-related quality of life (Rhinoconjunctivitis Quality of Life Questionnaire – RQLQ) in all 4 studies.

In comparison to a marketed fluticasone propionate nasal spray, substantial symptom improvement (50% reduction in nasal symptoms severity) was achieved significantly earlier (3 days and more) with azelastine /fluticasone nasal spray. The superior effect of azelastine/fluticasone nasal spray to fluticasone propionate nasal spray was maintained throughout a one-year study in patients with chronic persistent allergic rhinitis and nonallergic/vasomotor rhinitis.

In a ragweed pollen allergen exposure chamber study, first statistically significant relief of nasal symptoms was observed at 5 minutes after administration of azelastine/fluticasone nasal spray (compared to placebo). At 15 minutes after administration of azelastine /fluticasone nasal spray , 60% of patients reported a clinically relevant reduction in symptom scores of at least 30%.

## 5.2 Pharmacokinetic properties

### Absorption

After intranasal administration of two sprays per nostril (548 mcg of azelastine hydrochloride and 200 mcg of fluticasone) of azelastine/fluticasone nasal spray, the mean ( $\pm$  standard deviation) peak plasma exposure ( $C_{max}$ ) was  $194.5 \pm 74.4$  pg/ml for azelastine and  $10.3 \pm 3.9$  pg/ml for fluticasone propionate and the mean total exposure (AUC) was  $4217 \pm 2618$  pg/ml\*hr for azelastine and  $97.7 \pm 43.1$  pg/ml\*hr for fluticasone. The median time to peak exposure ( $t_{max}$ ) from a single dose was 0.5 hours for azelastine and 1.0 hours for fluticasone.

Fluticasone systemic exposure was ~50% increased comparing azelastine /fluticasone nasal spray with a marketed fluticasone nasal spray. Azelastine/fluticasone nasal spray was equivalent to a marketed azelastine nasal spray with respect to azelastine systemic exposure. There was no evidence of pharmacokinetic interactions between azelastine hydrochloride and fluticasone propionate.

### Distribution

Fluticasone propionate has a large volume of distribution at steady-state (approximately 318 liters). Plasma protein binding is 91%.

The volume of distribution of azelastine is high indicating distribution predominantly into the peripheral tissue. The level of protein binding is 80-90%. Additionally, both drugs have broad therapeutic windows. Therefore, drug displacement reactions are unlikely.

### Biotransformation

Fluticasone propionate is cleared rapidly from the systemic circulation, principally by hepatic metabolism to an inactive carboxylic acid metabolite, by the cytochrome P450 enzyme CYP3A4. Swallowed fluticasone propionate is also subject to extensive first pass metabolism. Azelastine is metabolized to N-desmethylazelastine via various CYP isoenzymes, mainly CYP3A4, CYP2D6 and CYP2C19.

### Elimination

The elimination rate of intravenous administered fluticasone propionate is linear over the 250-1000 microgram dose range and are characterized by a high plasma clearance (CL=1.1 l/min). Peak plasma concentrations are reduced by approximately 98%

within 3 - 4 hours and only low plasma concentrations were associated with the 7.8 h terminal half-life. The renal clearance of fluticasone propionate is negligible (<0.2%) and less than 5% as the carboxylic acid metabolite. The major route of elimination is the excretion of fluticasone propionate and its metabolites in the bile.

Plasma elimination half-lives after a single dose of azelastine are approximately 20 - 25 hours for azelastine and about 45 hours for the therapeutically active metabolite N-desmethy laz elastine. Excretion occurs mainly via the faeces. The sustained excretion of small amounts of the dose in the faeces suggests that some enterohepatic circulation may take place.

### **5.3 Preclinical safety data**

#### *Fluticasone propionate*

Findings in general toxicology studies were similar to those observed with other glucocorticoids and are associated with exaggerated pharmacological activity. These findings are not likely to be relevant for humans given recommended nasal doses which results in minimal systemic exposure. No genotoxic effects of fluticasone propionate have been observed in conventional genotoxicity tests. Further, there were no treatment-related increases in the incidence of tumours in two-year inhalation studies in rats and mice.

In animal studies glucocorticoids have been shown to induce malformations including cleft palate and intra-uterine growth retardation. Again, this is not likely to be relevant for humans given recommended nasal doses which results in minimal systemic exposure (see section 5.2).

#### *Azelastine hydrochloride*

Azelastine hydrochloride displayed no sensitizing potential in the guinea pig. Azelastine demonstrated no genotoxic potential in a battery of in vitro and in vivo tests, nor any carcinogenic potential in rats or mice. In male and female rats, azelastine at oral doses greater than 3 mg/kg/day caused a dose-related decrease in the fertility index.

No substance-related alterations were found in the reproductive organs of males or females during chronic toxicity studies, however, embryotoxic and teratogenic effects in rats, mice and rabbits occurred only at maternal toxic doses (for example, skeletal malformations were observed in rats and mice at doses of 68.6 mg/kg/day).

#### *Azelastine/fluticasone nasal spray*

Repeated-dose intranasal toxicity studies in rats for a period up to 90 days and in dogs for 14 days with azelastine/fluticasone nasal spray revealed no new adverse effects in comparison to the individual components.

## **6 PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Disodium edetate  
Glycerol  
Microcrystalline cellulose  
Carmellose sodium  
Polysorbate 80  
Benzalkonium chloride  
Phenylethyl alcohol  
Purified water

### **6.2 Incompatibilities**

Not applicable

### **6.3 Shelf life**

36 months  
In-use shelf-life (after first use): 6 months.

### **6.4 Special precautions for storage**

Do not refrigerate or freeze

### **6.5 Nature and contents of container**

Type I amber glass bottle fitted with a spray pump, a nasal polypropylene applicator (actuator) and a dust cap, containing 23 g (at least 120 actuations) suspension.

Pack sizes:

1 bottle with 23 g suspension in 25 mL bottles (at least 120 actuations)

## **6.6 Special precautions for disposal**

Not applicable

## **7 MARKETING AUTHORISATION HOLDER**

Zentiva Pharma UK Limited,  
12 New Fetter Lane,  
London, EC4A 1JP,  
United Kingdom

## **8 MARKETING AUTHORISATION NUMBER(S)**

PL 17780/1213

## **9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

07/11/2024

## **10 DATE OF REVISION OF THE TEXT**

07/11/2024