

# Arimidex® 1 mg film-coated tablets

(anastrozole)

The medicine is available as the above name but will be referred to as Arimidex throughout this leaflet.

## Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, or pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, or pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

## What is in this leaflet

1. What Arimidex is and what it is used for
2. What you need to know before you take Arimidex
3. How to take Arimidex
4. Possible side effects
5. How to store Arimidex
6. Contents of the pack and other information

### 1. What Arimidex is and what it is used for

Arimidex contains a substance called anastrozole. This belongs to a group of medicines called 'aromatase inhibitors'. Arimidex is used to treat breast cancer in women who have gone through the menopause and as a preventative treatment in postmenopausal women at moderate or high risk of breast cancer.

Arimidex works by cutting down the amount of the hormone called oestrogen that your body makes. It does this by blocking a natural substance (an enzyme) in your body called 'aromatase'.

### 2. What you need to know before you take Arimidex

#### Do not take Arimidex if you:

- are allergic to anastrozole or any of the other ingredients of this medicine (listed in section 6).
- are pregnant or breast-feeding (see the section called 'Pregnancy and breast-feeding').

Do not take Arimidex if any of the above apply to you. If you are not sure, talk to your doctor or pharmacist before taking Arimidex.

#### Warnings and precautions

Talk to your doctor, or pharmacist or nurse before taking Arimidex

- if you still have menstrual periods and have not yet gone through the menopause.
- if you are taking a medicine that contains tamoxifen or medicines that contain oestrogen (see the section called 'Other medicines and Arimidex').
- if you have or have ever had a condition that affects the strength of your bones (osteoporosis or osteopenia). Arimidex lowers the levels of female hormones and this may lead to a loss of the mineral content of bones, which might decrease their strength. You may have to have bone density tests during treatment. Your doctor can give you medicine to prevent or treat the bone loss. Women with severe osteoporosis are not suitable for anastrozole treatment.
- if you have problems with your liver or kidneys.
- if you have heart problems or have had a stroke.

If you are not sure if any of the above applies to you, talk to your doctor or pharmacist before taking Arimidex.

If you go into the hospital, let the medical staff know you are taking Arimidex.

## Other medicines and Arimidex

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. This includes medicines that you buy without a prescription and herbal medicines. This is because Arimidex can affect the way some medicines work and some medicines can have an effect on Arimidex.

Do not take Arimidex if you are already taking any of the following medicines:

- Certain medicines used to treat breast cancer (selective oestrogen receptor modulators), e.g. medicines that contain tamoxifen. This is because these medicines may stop Arimidex from working properly.
- Medicines that contain oestrogen, such as hormone replacement therapy (HRT).

If this applies to you, ask your doctor or pharmacist for advice.

Tell your doctor or pharmacist if you are taking the following:

- A medicine known as an 'LHRH analogue', this includes gonadorelin, buserelin, goserelin, leuprorelin and triptorelin. These medicines are used to treat breast cancer, certain female health (gynaecological) conditions, and infertility.

## Pregnancy and breast-feeding

Do not take Arimidex if you are pregnant or breast-feeding. Stop Arimidex if you become pregnant and talk to your doctor.

If you are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

## Driving and using machines

Arimidex is not likely to affect your ability to drive or use any tools or machines. However, some people may occasionally feel weak or sleepy while taking Arimidex. If this happens to you, ask your doctor or pharmacist for advice.

## Arimidex contains lactose

Arimidex contains lactose which is a type of sugar. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

## Information on sodium content

This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium free'.

### 3. How to take Arimidex

Always take this medicine exactly as your doctor or pharmacist has told you.

Check with your doctor or pharmacist if you are not sure.

- The recommended dose is one tablet once a day.
- Try to take your tablet at the same time each day.
- Swallow the tablet whole with a drink of water.
- It does not matter if you take Arimidex before, with or after food.

Keep taking Arimidex for as long as your doctor or pharmacist tells you to. It is a long-term treatment and you may need to take it for several years.

Check with your doctor or pharmacist if you are not sure.

## Use in children and adolescents

Arimidex should not be given to children and adolescents.

## If you take more Arimidex than you should

If you take more Arimidex than you should, talk to a doctor straight away.

## If you forget to take Arimidex

If you forget to take a dose, just take your next dose as normal. Do not take a double dose (two doses at the same time) to make up for a forgotten dose.

## If you stop taking Arimidex

Do not stop taking your tablets unless your doctor tells you to.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

## 4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

If any of the side effects get worse, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

**Stop taking Arimidex and seek urgent medical treatment, if you experience any of the following serious side effects:**

**Common** (may affect up to 1 in 10 people):

- Allergic (hypersensitivity) reactions including face, lips, or tongue.

**Rare** (may affect up to 1 in 1,000 people):

- Rare inflammation of your skin that may include red patches or blisters (erythema multiforme).

**Very rare** (may affect up to 1 in 10,000 people):

- An extremely severe skin reaction with ulcers or blisters on the skin. This is known as 'Stevens-Johnson syndrome'
- Swelling of the throat that may cause difficulty in swallowing or breathing. This is known as 'angioedema'.

### Other side effects

**Very common side effects (affect more than 1 in 10 people)**

- Headache
- Hot flushes
- Feeling sick (nausea)
- Skin rash
- Pain or stiffness in your joints
- Inflammation of the joints (arthritis)
- Feeling weak
- Bone loss (osteoporosis)
- Depression.

**Common side effects (affect 1 to 10 people in 100)**

- Loss of appetite
- Raised or high levels of fatty substance known as cholesterol in your blood. This would be seen in a blood test
- Feeling sleepy
- Carpal tunnel syndrome (tingling, pain, coldness, weakness in parts of the hand)
- Tickling, tingling or numbness of skin, loss/lack of taste
- Diarrhoea
- Being sick (vomiting)
- Changes in blood tests that show how well your liver is working
- Thinning of your hair (hair loss)
- Bone pain
- Vaginal dryness
- Bleeding from the vagina (usually in the first few weeks of treatment – if the bleeding continues, talk to your doctor)
- Muscle pain.

**Uncommon side effects (affect 1 to 10 people in 1,000)**

- Changes in special blood tests that show how your liver is working (gamma-GT and bilirubin)
- Inflammation of the liver (hepatitis)
- Hives or nettle rash
- Trigger finger (a condition in which your finger or thumb catches in a bent position)
- Increased amounts of calcium in your blood. If you experience nausea, vomiting and thirst, you should tell your doctor, or pharmacist or nurse as you may need to have blood tests.

**Rare side effects (affect 1 to 10 people in 10,000)**

- Inflammation of the small blood vessels causing red or purple colouring of the skin. Very rarely symptoms of joint, stomach, and kidney pain may occur; this is known as 'Henoch-Schönlein purpura'.

**Side effects with frequency not known (frequency cannot be estimated from the available data)**

- Dry eye
- Lichenoid eruption (small, red or purple itchy bumps on the skin)
- Inflammation of a tendon or tendonitis (connective tissues that connect muscles to bones)
- Rupture of a tendon (connective tissues that connect muscles to bones)
- Memory impairment

## Effects on your bones

Arimidex lowers the amount of the hormone called oestrogen that is in your body. This may lower the mineral content of your bones. Your bones may be less strong and may be more likely to fracture. Your doctor will manage these risks according to treatment guidelines for managing bone health in women who have gone through the menopause. You should talk to your doctor about the risks and treatment options.

## Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects via the Yellow Card Scheme Website: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects, you can help provide more information on the safety of this medicine.

## 5. How to store Arimidex

**Keep this medicine out of the sight and reach of children.** Keep your tablets in a safe place where children cannot see or reach them. Your tablets could harm them.

Do not use this medicine after the expiry date which is stated on the carton and blister strip after 'EXP'. The expiry date refers to the last day of that month.

Do not store above 30°C.

Keep your tablets in the container they came in.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

If your tablets become discoloured or show signs of any deterioration, consult your doctor or pharmacist who will tell you what to do.

## 6. Contents of the pack and other information

### What Arimidex contains

- The active substance is anastrozole. Each film-coated tablet contains 1 mg anastrozole.
- The other ingredients are: lactose monohydrate, povidone, sodium starch glycollate, magnesium stearate, hypromellose, macrogol 300, titanium dioxide.

### What Arimidex looks like and contents of the pack

White, round, biconvex film-coated tablets of about 6.1 mm marked 'A' on one side and 'Adx1' on the other side. Arimidex comes in blister packs of 28 tablets.

PL: 15814/1973

POM

Manufactured by Haupt Pharma Muenster GmbH, Schleebrueggenkamp 15, Muenster Nordrhein-Westfalen, 48 159, Germany.

Procured from within the EU and repackaged by the Product Licence holder: O.P.D. Laboratories Ltd., Unit 6 Colonial Way, Watford, Herts WD24 4PR.

Leaflet revision and issue date (Ref.): 15.07.2025.

Arimidex is a registered trademark of Laboratoires Juvise Pharmaceuticals.

**To request a copy of this leaflet in Braille, large print or audio please call 01923 332 796.**

## Anastrozole 1 mg film-coated tablets

The medicine is available as the above name but will be referred to as Anastrozole tablets throughout this leaflet.

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, or pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, or pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

### What is in this leaflet

1. What Anastrozole tablets are and what they are used for
2. What you need to know before you take Anastrozole tablets
3. How to take Anastrozole tablets
4. Possible side effects
5. How to store Anastrozole tablets
6. Contents of the pack and other information

### 1. What Anastrozole tablets are and what they are used for

Anastrozole tablets contain a substance called anastrozole. This belongs to a group of medicines called 'aromatase inhibitors'. Anastrozole tablets are used to treat breast cancer in women who have gone through the menopause and as a preventative treatment in postmenopausal women at moderate or high risk of breast cancer.

Anastrozole tablets work by cutting down the amount of the hormone called oestrogen that your body makes. It does this by blocking a natural substance (an enzyme) in your body called 'aromatase'.

### 2. What you need to know before you take Anastrozole tablets

#### Do not take Anastrozole tablets if you:

- are allergic to anastrozole or any of the other ingredients of this medicine (listed in section 6).
- are pregnant or breast-feeding (see the section called 'Pregnancy and breast-feeding').

Do not take Anastrozole tablets if any of the above apply to you. If you are not sure, talk to your doctor or pharmacist before taking Anastrozole tablets.

#### Warnings and precautions

Talk to your doctor, or pharmacist or nurse before taking Anastrozole tablets

- if you still have menstrual periods and have not yet gone through the menopause.
- if you are taking a medicine that contains tamoxifen or medicines that contain oestrogen (see the section called 'Other medicines and Anastrozole tablets').
- if you have or have ever had a condition that affects the strength of your bones (osteoporosis or osteopenia).  
Anastrozole tablets lowers the levels of female hormones and this may lead to a loss of the mineral content of bones, which might decrease their strength. You may have to have bone density tests during treatment. Your doctor can give you medicine to prevent or treat the bone loss. Women with severe osteoporosis are not suitable for anastrozole treatment.
- if you have problems with your liver or kidneys.
- if you have heart problems or have had a stroke.

If you are not sure if any of the above applies to you, talk to your doctor or pharmacist before taking Anastrozole tablets.

If you go into the hospital, let the medical staff know you are taking Anastrozole tablets.

#### Other medicines and Anastrozole tablets

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. This includes medicines that you buy without a prescription and herbal medicines. This is because Anastrozole tablets can affect the way some medicines work and some medicines can have an effect on Anastrozole tablets.

Do not take Anastrozole tablets if you are already taking any of the following medicines:

- Certain medicines used to treat breast cancer (selective oestrogen receptor modulators), e.g. medicines that contain tamoxifen. This is because these medicines may stop Anastrozole tablets from working properly.
- Medicines that contain oestrogen, such as hormone replacement therapy (HRT).

If this applies to you, ask your doctor or pharmacist for advice.

Tell your doctor or pharmacist if you are taking the following:

- A medicine known as an 'LHRH analogue', this includes gonadorelin, buserelin, goserelin, leuprorelin and triptorelin. These medicines are used to treat breast cancer, certain female health (gynaecological) conditions, and infertility.

#### Pregnancy and breast-feeding

Do not take Anastrozole tablets if you are pregnant or breast-feeding. Stop Anastrozole tablets if you become pregnant and talk to your doctor.

If you are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

#### Driving and using machines

Anastrozole tablets are not likely to affect your ability to drive or use any tools or machines. However, some people may occasionally feel weak or sleepy while taking Anastrozole tablets. If this happens to you, ask your doctor or pharmacist for advice.

#### Anastrozole tablets contains lactose

Anastrozole tablets contains lactose which is a type of sugar. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

#### Information on sodium content

This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium free'.

### 3. How to take Anastrozole tablets

Always take this medicine exactly as your doctor or pharmacist has told you.

Check with your doctor or pharmacist if you are not sure.

- The recommended dose is one tablet once a day.
- Try to take your tablet at the same time each day.
- Swallow the tablet whole with a drink of water.
- It does not matter if you take Anastrozole tablets before, with or after food.

Keep taking Anastrozole tablets for as long as your doctor or pharmacist tells you to. It is a long-term treatment and you may need to take it for several years.

Check with your doctor or pharmacist if you are not sure.

#### Use in children and adolescents

Anastrozole tablets should not be given to children and adolescents.

#### If you take more Anastrozole tablets than you should

If you take more Anastrozole tablets than you should, talk to a doctor straight away.

#### If you forget to take Anastrozole tablets

If you forget to take a dose, just take your next dose as normal. Do not take a double dose (two doses at the same time) to make up for a forgotten dose.

#### If you stop taking Anastrozole tablets

Do not stop taking your tablets unless your doctor tells you to. If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

## 4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

If any of the side effects get worse, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

**Stop taking Anastrozole tablets and seek urgent medical treatment, if you experience any of the following serious side effects:**

**Common** (may affect up to 1 in 10 people):

- Allergic (hypersensitivity) reactions including face, lips, or tongue.

**Rare** (may affect up to 1 in 1,000 people):

- Rare inflammation of your skin that may include red patches or blisters (erythema multiforme).

**Very rare** (may affect up to 1 in 10,000 people):

- An extremely severe skin reaction with ulcers or blisters on the skin. This is known as 'Stevens-Johnson syndrome'
- Swelling of the throat that may cause difficulty in swallowing or breathing. This is known as 'angioedema'.

### Other side effects

**Very common side effects (affect more than 1 in 10 people)**

- Headache
- Hot flushes
- Feeling sick (nausea)
- Skin rash
- Pain or stiffness in your joints
- Inflammation of the joints (arthritis)
- Feeling weak
- Bone loss (osteoporosis)
- Depression.

**Common side effects (affect 1 to 10 people in 100)**

- Loss of appetite
- Raised or high levels of fatty substance known as cholesterol in your blood. This would be seen in a blood test
- Feeling sleepy
- Carpal tunnel syndrome (tingling, pain, coldness, weakness in parts of the hand)
- Tickling or numbness of skin, loss/lack of taste
- Diarrhoea
- Being sick (vomiting)
- Changes in blood tests that show how well your liver is working
- Thinning of your hair (hair loss)
- Bone pain
- Vaginal dryness
- Bleeding from the vagina (usually in the first few weeks of treatment – if the bleeding continues, talk to your doctor)
- Muscle pain.

**Uncommon side effects (affect 1 to 10 people in 1,000)**

- Changes in special blood tests that show how your liver is working (gamma-GT and bilirubin)
- Inflammation of the liver (hepatitis)
- Hives or nettle rash
- Trigger finger (a condition in which your finger or thumb catches in a bent position)
- Increased amounts of calcium in your blood. If you experience nausea, vomiting and thirst, you should tell your doctor, or pharmacist or nurse as you may need to have blood tests.

**Rare side effects (affect 1 to 10 people in 10,000)**

- Inflammation of the small blood vessels causing red or purple colouring of the skin. Very rarely symptoms of joint, stomach, and kidney pain may occur; this is known as 'Henoch-Schönlein purpura'.

**Side effects with frequency not known (frequency cannot be estimated from the available data)**

- Dry eye
- Lichenoid eruption (small, red or purple itchy bumps on the skin)
- Inflammation of a tendon or tendonitis (connective tissues that connect muscles to bones)
- Rupture of a tendon (connective tissues that connect muscles to bones)
- Memory impairment

## Effects on your bones

Anastrozole tablets lower the amount of the hormone called oestrogen that is in your body. This may lower the mineral content of your bones. Your bones may be less strong and may be more likely to fracture. Your doctor will manage these risks according to treatment guidelines for managing bone health in women who have gone through the menopause. You should talk to your doctor about the risks and treatment options.

## Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects via the Yellow Card Scheme Website: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects, you can help provide more information on the safety of this medicine.

## 5. How to store Anastrozole tablets

**Keep this medicine out of the sight and reach of children.** Keep your tablets in a safe place where children cannot see or reach them. Your tablets could harm them.

Do not use this medicine after the expiry date which is stated on the carton and blister strip after 'EXP'. The expiry date refers to the last day of that month.

Do not store above 30°C.

Keep your tablets in the container they came in.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

If your tablets become discoloured or show signs of any deterioration, consult your doctor or pharmacist who will tell you what to do.

## 6. Contents of the pack and other information

### What Anastrozole tablets contain

- The active substance is anastrozole. Each film-coated tablet contains 1 mg anastrozole.
- The other ingredients are: lactose monohydrate, povidone, sodium starch glycollate, magnesium stearate, hypromellose, macrogol 300, titanium dioxide.

### What Anastrozole tablets look like and contents of the pack

White, round, biconvex film-coated tablets of about 6.1 mm marked 'A' on one side and 'Adx1' on the other side.

Anastrozole tablets comes in blister packs of 28 tablets.

PL: 15814/1973

POM

Manufactured by Haupt Pharma Muenster GmbH, Schleebrueggkamp 15, Muenster Nordrhein-Westfalen, 48 159, Germany.

Procured from within the EU and repackaged by the Product Licence holder: O.P.D. Laboratories Ltd., Unit 6 Colonial Way, Watford, Herts WD24 4PR.

Leaflet revision and issue date (Ref.): 15.07.2025

**To request a copy of this leaflet in Braille, large print or audio please call 01923 332 796.**