

PACKAGE LEAFLET: INFORMATION FOR THE USER
CYPROHEPTADINE HYDROCHLORIDE 4MG TABLETS
(Cyproheptadine Hydrochloride)

Read all of this leaflet carefully because it contains important information for you.

This medicine is available without prescription. However, you still need to take Cyproheptadine Hydrochloride tablets carefully to get the best results from it.

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- You must contact a doctor if your symptoms worsen or do not improve.
- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

In this leaflet:

1. What Cyproheptadine Hydrochloride tablets is and what it is used for
2. Before you take Cyproheptadine Hydrochloride tablets
3. How to take Cyproheptadine Hydrochloride tablets
4. Possible side effects
5. How to store Cyproheptadine Hydrochloride tablets
6. Further Information

1. WHAT CYPROHEPTADINE HYDROCHLORIDE TABLETS IS AND WHAT IT IS USED FOR

Cyproheptadine Hydrochloride tablets contains a medicine called cyproheptadine. This belongs to a group of medicines called ‘antihistamines’. In some illnesses too much ‘histamine’ and ‘serotonin’ are produced in the body.

These occur naturally in your body. Cyproheptadine Hydrochloride tablets helps in these illnesses by stopping the unwanted effects of histamine and serotonin.

Cyproheptadine Hydrochloride tablets is used for:

- skin allergies with itchiness, swelling, or red skin, eczema or nettle rash
- itchiness of chicken pox and reactions to insect bites and stings
- sneezing, watery eyes and blocked or runny nose caused by an allergy (allergic rhinitis)
- conjunctivitis caused by an allergy- with red, watery, inflamed eyes
- migraine and other similar headaches.

2. BEFORE YOU TAKE CYPROHEPTADINE HYDROCHLORIDE TABLETS

Do not take Cyproheptadine Hydrochloride tablets if:

- you are allergic (hypersensitive) to cyproheptadine or any of the other ingredients of Cyproheptadine Hydrochloride tablets (listed in section 6 below)
- you are taking a type of medicine called a monoamine oxidase inhibitor (MAOI) for depression, such as meclobemide, phenelzine, tranylcypamine, linezolid or pargyline.
- you have a narrowing of the opening from the stomach (‘stricture’)
- you have an eye problem called ‘glaucoma’

- you are being treated for an asthmatic attack
- you are elderly and not very mobile
- you have problems passing urine
- you have a stomach ulcer.

Do not take Cyproheptadine Hydrochloride tablets if any of the above apply to you. If you are not sure talk to your doctor or pharmacist before taking Cyproheptadine Hydrochloride tablets.

Take special care with Cyproheptadine Hydrochloride tablets

Check with your doctor before taking your medicine if:

- you have ever been told that you have high pressure inside your eye (high intra-ocular pressure)
- you have an over-active thyroid gland
- you have high blood pressure
- you have heart problems
- you have asthma.

If you are not sure if any of the above apply to you, talk to your doctor or pharmacist before taking Cyproheptadine Hydrochloride tablets.

Taking other medicines

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription. This includes herbal medicines. This is because Cyproheptadine Hydrochloride tablets can affect the way some other medicines work. Also some medicines can affect the way Cyproheptadine Hydrochloride tablets work.

Do not take Cyproheptadine Hydrochloride tablets if you are taking a type of medicine called a 'monoamine oxidase inhibitor' (MAO inhibitor) such as meclobemide, phenelzine, tranylcypamine, linezolid or pargyline.

Also, it is important to tell your doctor or pharmacist if you take:

- sleeping pills - such as temazepam, nitrazepam, zopiclone, zaleplon or zolpidem
- medicines for worry or anxiety - such as diazepam, alprazolam, chlordiazepoxide, clorazepate
- medicines for depression- such as fluoxetine, paroxetine, amitriptyline or imipramine.
- any medicines that make you sleepy - such as sedatives or tranquillisers.

If you are not sure if any of the above apply to you, talk to your doctor or pharmacist before taking Cyproheptadine Hydrochloride tablets.

Taking Cyproheptadine Hydrochloride tablets with food and drink

Do not drink alcohol while you are taking Cyproheptadine Hydrochloride tablets. This is because Cyproheptadine Hydrochloride tablets can make you sleepy and drinking alcohol at the same time can make you even more sleepy.

Pregnancy and breast-feeding

Do not take Cyproheptadine Hydrochloride tablets if you are or think you may be pregnant. Ask your doctor or pharmacist for advice before taking any medicine.

Driving and using machines

Do not drive or use any tools or machines while taking this medicine. This is because Cyproheptadine Hydrochloride tablets can make you sleepy.

Important information about some of the ingredients of Cyproheptadine Hydrochloride tablets

Cyproheptadine Hydrochloride tablets contains lactose, which is a type of sugar. If you have been told by your doctor that you can not tolerate or digest some sugars (have an intolerance to some sugars), talk to your doctor before taking this medicine.

3. HOW TO TAKE CYPROHEPTADINE HYDROCHLORIDE TABLETS

If this medicine is from your doctor or pharmacist, do exactly as they have told you. Otherwise follow the instructions below.

Taking this medicine

- Take this medicine by mouth.
- If you do not understand the instructions, or you are not sure, ask your doctor or pharmacist.

The usual doses of Cyproheptadine Hydrochloride tablets are

For allergies

Adults and children over 14 years

- Start with 1 tablet three times a day.
- Most patients need 3-4 tablets a day.
- Do not take more than 8 tablets a day.

Children aged 7-14 years

- The usual dose is 1 tablet two or three times a day as needed.
- If you need to give an extra dose, give it at bedtime.
- Do not give more than 4 tablets a day.

Children aged 2-6 years

- The usual dose is half a tablet two or three times a day.
- If you need to give an extra dose, give it at bedtime.
- Do not give more than 3 tablets a day.

For migraine

- Take 1 tablet when you are getting the migraine
- Take another one, if needed, after 30 minutes.
- Then take 1 tablet every 4-6 hours to maintain relief.
- Do not take more than 2 tablets in any 4-6 hour period.

Children under 2 years

- Do not give Cyproheptadine Hydrochloride tablets to children under 2 years of age.

If you take more Cyproheptadine Hydrochloride tablets than you should

If you take too much Cyproheptadine Hydrochloride tablets by mistake, contact your doctor immediately or seek medical help as soon as possible.

If you forget to take Cyproheptadine Hydrochloride tablets

- If you forget to take a tablet, skip the missed dose.
- Take the next dose as usual.
- Do not take a double dose to make up for a forgotten dose.

If you have any further questions on the use of this product, ask your doctor or pharmacist.

4. POSSIBLE SIDE EFFECTS

Like all medicines, Cyproheptadine Hydrochloride tablets can cause side effects, although not everybody gets them.

Stop taking Cyproheptadine Hydrochloride tablets and tell your doctor straight away, if you notice any of the following serious side effects- you may need urgent medical treatment:

- allergic skin rash and hives
- severe allergic reaction which causes difficulty in breathing or dizziness

Other side effects include:

Heart and circulation

- increased heart rate
- low blood pressure
- palpitations
- feeling like you have a missed heart beat

Blood

- nosebleeds

Nervous system

- feeling sleepy, dizzy, problems with co-ordination, or feeling tired
- feeling confused, restless, over excited, nervous, irritable or trembling
- being unable to sleep, pins and needles or feeling numb
- inflammation of your nerves-signs include being more sensitive to pain and feeling that you can not move (paralysis), fits (convulsions)
- feeling aggressive
- seeing, hearing, or feeling something that is not there (hallucinations)
- feeling faint
- headache
- an exaggerated feeling of well being (euphoria)
- being very upset (hysteria)
- anti-histamine medicines such as this medicine can sometimes make young children over excited

Eyes or ears

- Inflammation inside the ear, ringing in the ears
- blurred or double vision
- vertigo

Chest

- dry throat and nose
- presence of phlegm
- blocked nose
- feeling wheezy
- tight chest

Stomach and gut

- dry mouth
- stomach ache, loss of appetite
- feeling sick (nausea), being sick (vomiting)
- increase in your appetite and weight gain
- diarrhoea or constipation

Urinary

- passing urine more often
- difficulty passing urine
- being unable to empty your bladder of urine

Skin and hair

- your skin being sensitive to sunlight
- you may sweat more than usual
- severe shivering
- swelling of your skin

Liver

- liver problems, including jaundice, hepatitis and hepatic failure - signs include yellowing of your skin and eyes, sometimes with a fever, pale stools and dark urine

Sexual

- menstrual problems

If any of the side effects get serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

5. HOW TO STORE CYPROHEPTADINE HYDROCHLORIDE TABLETS

Keep out of the sight and reach of children.

Do not store above 25°C. Store in the original package in order to protect from light.

Do not transfer your medicines into another box or container - they might get mixed up.

Do not use Cyproheptadine Hydrochloride tablets after the expiry date which is stated on the carton after EXP. The expiry date refers to the last day of that month.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist to dispose of medicines no longer required. These measures will help to protect the environment.

6. FURTHER INFORMATION

What Cyproheptadine Hydrochloride tablets contains

Active ingredient:

The active substance is cyproheptadine hydrochloride. Each Cyproheptadine Hydrochloride tablets contains 4 mg anhydrous cyproheptadine Hydrochloride.

Other ingredients:

The other ingredients are calcium hydrogen phosphate E341, lactose, magnesium stearate E572, potato starch, pregelatinised maize starch.

What Cyproheptadine Hydrochloride tablets looks like and contents of the pack

Cyproheptadine Hydrochloride tablets are white round bevel-edged tablets marked 'MSD 62' on one side with a single scoreline on the other.

Cyproheptadine Hydrochloride tablets are supplied in blister packs of 30 tablets.

Marketing Authorisation Holder and Manufacturer

The Marketing Authorisation is held by Teva UK Limited, Whistler Drive, Castleford, WF10 5HX, United Kingdom.

Cyproheptadine Hydrochloride tablets are made by Tiofarma B.V., Hermanus Boerhaavestraat 1, 3261 ME Oud-Beijerland, The Netherlands.

This leaflet was last revised in July 2025.

PL 00289/2271