

**Package leaflet: Information for the user**  
**Zibor 25,000 IU anti-Xa/ml solution for injection in pre-filled syringes**  
Bemiparin sodium

**Read all of this leaflet carefully before you start using this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

**What is in this leaflet**

1. What Zibor is and what it is used for
2. What you need to know before you use Zibor
3. How to use Zibor
4. Possible side effects
5. How to store Zibor
6. Contents of the pack and other information

**1. What Zibor is and what it is used for**

The active ingredient in Zibor is bemiparin sodium, which belongs to a group of medicines called anticoagulants. These help to stop blood from clotting in the blood vessels. Zibor is used to treat dangerous blood clots, which have formed in, for example, the veins of the legs and/or the lungs (deep vein thrombosis and/or pulmonary embolism).

**2. What you need to know before you use Zibor**

**Do not use Zibor if you:**

- are allergic to bemiparin sodium, heparin or a similar product (such as enoxaparin, dalteparin, nadroparin) or any other ingredients of this medicine (listed in section 6).
- have had an allergic reaction after being given any medicine containing heparin.
- are allergic to any substance derived from pigs.
- suffer from Heparin Induced Thrombocytopenia (HIT), a condition that produces a severe decrease in your number of blood-clotting cells (platelets), or, as a result of HIT, you suffer from another condition called, Disseminated Intravascular Coagulation (DIC), where your blood-clotting cells would clump together if Zibor is used.
- suffer from a condition known as endocarditis (inflammation of the lining of the heart and heart valves).
- suffer from any condition which results in a tendency to bleed excessively.
- suffer from a serious liver and/or pancreas disease.
- have any damage to your internal organs which may lead to a high risk of internal bleeding (for example, an active stomach ulcer, cerebral aneurisms [swelling in the artery walls in the brain], or brain tumours).
- have suffered from a brain haemorrhage.

- have an injury to, or are going to have an operation on, your brain, spine, eyes and/or ears.
- are being treated with Zibor, you must not have epidural or spinal anaesthesia (an anaesthetic injected into your spine) because it could be dangerous. Therefore, make sure your doctor knows that you are being treated with Zibor before any surgery.

### **Warnings and precautions**

Talk to your doctor before using Zibor if you:

- suffer from liver disease.
- suffer from kidney disease. Your doctor may consider conducting special monitoring. In case your disease is severe, your doctor may consider adjusting your dose.
- have high and/or difficult to control blood pressure. have ever had a stomach ulcer which is no longer active.
- suffer from thrombocytopenia, a condition where there are fewer than normal blood-clotting cells (platelets) in your blood, making you bruise and bleed easily.
- have kidney stones and/or bladder stones.
- suffer from any condition which may cause you to bleed more easily.
- suffer from eye problems, due to problems in your blood vessels.
- suffer from diabetes.
- have blood test results showing that you have high levels of potassium in your blood.
- are going to have a lumbar puncture (a puncture in the lower part of the spine for laboratory tests) make doubly sure your doctor knows you are being treated with Zibor.

### **Other medicines and Zibor**

Check with your doctor if you think you may already be taking:

- any medicine, which is injected into a muscle, because such injections must be avoided during treatment with Zibor.
- other anticoagulants such as warfarin and/or acenocoumarol (Vitamin K antagonists), to treat and/or prevent blood clots.
- non-steroidal anti-inflammatory drugs, such as ibuprofen, for example, for arthritis.
- steroids, such as prednisolone, to treat inflammatory diseases, such as arthritis.
- platelet inhibitors, such as aspirin, ticlopidine or clopidogrel, to prevent blood clots.
- medicines which can increase levels of potassium in your blood, such as some diuretics (water pills) and anti-hypertensives (used to reduce blood pressure).
- medicines to increase your blood volume, such as dextran.
- an injected drug used to treat heart problems, called glyceryl nitrate.

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines, including medicines obtained without a prescription.

### **Special tests you may need**

- Some patients may need to have the level of blood clotting cells in their blood checked. Your doctor will decide whether this is necessary and when (e.g. before treatment, on the first day of treatment, then every 3-4 days and at the end of treatment).
- If you suffer from certain conditions (diabetes, kidney disease) or if you are taking medicines to prevent the loss of potassium, your doctor may check the potassium level in your blood.

### **Pregnancy and breast-feeding**

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

### **Driving and using machines**

Zibor has no effect on the ability to drive or operate machinery.

## **3. How to use Zibor**

Always use this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

The recommended dose is:

### **Adults (aged 18 to 64)**

Your daily injection dose will depend upon your body weight. If you weigh:

- less than 50 Kilos, the dose will be 0.2 millilitres (= 5,000 IU).
- between 50 and 70 kilos, the dose will be 0.3 millilitres (=7,500 IU).
- between 71 and 100 kilos, the dose will be 0.4 millilitres (=10,000 IU).
- more than 100 kilos, the dose will be adjusted, depending on your exact weight, to the equivalent of 115 IU a day for each kilogram you weigh.

*IU: The potency of this medicine is described in International anti-Xa activity units.*

Zibor is usually injected under your skin, usually into a skin fold at the side of your waist (abdomen) or in the upper part of the thigh. Your doctor or nurse will usually administer the injection in hospital. You may need to continue to receive Zibor when you return home.

- This medicine must never be injected into a muscle or mixed with any other injection.
- It is usually given once a day.
- Your doctor will tell you for how long you should be given this medicine (usually for approximately 5 – 9 days).
- If your doctor has told you that you can inject this medicine yourself, please follow your doctor's instructions extremely carefully. (See below "How do I inject Zibor?").

**Elderly patients (65 years old and above)** are normally given the same dosages as other adult patients. If you have liver or kidney problems, please tell your doctor who may wish to keep a close eye on you. If you have severe kidney problems, your doctor may give you a specific dose.

### **Use in children (under 18 years of age)**

Zibor is not recommended for children.


### **How do I inject Zibor?**

Zibor should never be injected into muscle because that could cause bleeding into the muscle.


You should receive instructions on the correct way to use this medicine and the correct technique for self-injection before you give yourself an injection for the first time. These instructions should be given by a doctor or other qualified healthcare professional.

You should follow these steps:

- Wash your hands well and sit or lie in a comfortable position.
- Choose an area of the waist, at least 5 centimetres away from your belly button and from existing scars or bruises and clean the skin carefully.
- Use different places for the injection on different days, for example, first on the left hand side, next time on the right.
- Pull the needle cap off the Zibor syringe.
- To keep the needle sterile, make sure it doesn't touch anything.
- This pre-filled syringe is now ready for use.
- Before injecting, do not push the plunger to get rid of any air bubbles, because you might lose the medicine.



- Hold the syringe in one hand and with your other hand, using your forefinger and thumb, gently pinch the area of skin which you've cleaned and make a skin fold.
- Insert the full length of the needle into the folded skin, straight in at a 90° angle.
- Press down on the plunger, making sure you hold the skin fold in position throughout the injection.
- 



- Remove the needle by pulling it straight out and let go of the skin fold.
- Don't rub the skin where you put the needle in. This will help to avoid bruises.
- Don't try to put the needle cap back on the syringe. Just drop it (needle first) into a sharps bin, close the container lid tightly and place it out of reach of children.
- If you get the impression that the dose is either too strong (for example, you are experiencing unexpected bleeding) or too weak (for example, the dose doesn't seem to be working), talk to your doctor or pharmacist.

### **If you use more Zibor than you should**

This may result in bleeding. If this happens, tell your doctor immediately or go immediately to the casualty department at your nearest hospital with this leaflet.

### **If you forget to use Zibor**

Do not take a double dose to make up for a forgotten dose. You should consult your doctor as soon as possible so that he may tell you what to do.

### **If you stop using Zibor**

Always check with your doctor before you stop using this medicine.

If you have any further questions on the use of this medicine, ask your doctor.

## **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

**Stop using Zibor and tell the doctor or nurse immediately (or go immediately to your nearest Casualty or Accident and Emergency Department), if you notice any of the following side effects:**

#### **Common (these may affect up to 1 in 10 patients):**

- Unusual or unexpected bleeding, for example blood in your urine and/or stools which may cause haemorrhagic anaemia.

#### **Rare (these may affect up to 1 in 1,000 patients):**

- Severe decrease in your number of blood-clotting cells (thrombocytopenia type II) which can lead to bruising, bleeding in the mouth, gums and nose, rash.
- Dark painful skin reactions at the injection site (cutaneous necrosis).
- Intra-spinal haematomas following spinal or lumbar anaesthesia (back pain, numbness and weakness in the lower limbs, bowel or bladder dysfunction). These haematomas could cause various degrees of neurological impairment, including prolonged or permanent paralysis.
- Serious allergic reactions (raised temperature, shivering, breathlessness, swelling in your vocal cords, light-headedness, sweating, nettle rash/hives, itchy skin, low blood pressure, hot flushes, flushing, black out, contraction of the bronchial tube, swelling of the larynx).

#### **Other side effects:**

#### **Very Common (affect more than 1 in 10 patients):**

- Bruising, blotchy skin, itching and some pain at the places where the medicine was injected.

#### **Common (these may affect up to 1 in 10 patients):**

- A slight and temporary increase in certain enzymes (transaminases), which would show up in blood tests.

#### **Uncommon (these may affect up to 1 in 100 patients):**

- Mild and transient temporary decrease in your number of blood-clotting cells (thrombocytopenia type I), which would show up in blood tests.
- Mild allergic skin reactions: skin rash, nettle rash/hives, weals.

#### **Not known (cannot be estimated from the available data):**

- Increase in potassium levels, which could show up in blood tests.

Brittle bones (osteoporosis) may develop through use of this or similar medicines over a long period of time. The frequency is unknown.

### **Reporting of side effects**

If you get any side effects, talk to your doctor. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for 'MHRA Yellow Card' in the Google Play or Apple App Store.

By reporting side effects you can help provide more information on the safety of this medicine.

## **5. How to store Zibor**

Keep this medicine out of the sight and reach of children.  
Do not store above 25 °C. Do not freeze.

Do not use this medicine if you notice:

- the protective package has already been opened.
- the protective package is damaged.
- the medicine in the syringe appears cloudy.
- it contains small particles.

After the blister containing the syringe has been opened, the medicine should be used immediately.

### **Expiry date**

Do not use this medicine after the expiry date, which is stated on the carton.  
The expiry date refers to the last day of that month.

### **Disposal**

This medicine comes in single-dose syringes.

Drop used syringes into a sharps bin.

Do not keep them after use.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## **6. Contents of the pack and other information**

### **What Zibor contains**

- The active substance is bemiparin sodium.
- The other ingredient is water for injections.

### **What Zibor looks like and contents of the pack**

The medicine contained in the syringes is a clear, colourless or slightly yellowish solution, free of particles.

Zibor 25,000 IU is available in packs of 2, 10, 30 and 100 pre-filled syringes containing 0.2 ml, 0.3 or 0.4 ml of solution for injection.

Not all pack sizes may be marketed.

Each 0.2 ml syringe provides a dose of bemiparin sodium of 5,000 IU.  
Each 0.3 ml syringe provides a dose of bemiparin sodium of 7,500 IU.  
Each 0.4 ml syringe provides a dose of bemiparin sodium of 10,000 IU.

### **Marketing Authorisation Holder and Manufacturer**

#### **Marketing Authorisation Holder**

GINELADIUS, S.L.  
Rufino González 50, 28037 Madrid-Spain

#### **Manufacturer**

ROVI Pharma Industrial Services, S.A.  
Julián Camarillo 35, 28037  
Madrid – Spain

Laboratorios Farmacéuticos ROVI, S.A.  
Julián Camarillo 35,  
28037 Madrid – Spain

Local representative in UK:  
ROVI Biotech Limited  
One Elmfield Park, Office 2.06, Bromley, BR1 1LU,  
United Kingdom

#### **This medicinal product is authorised in the Member States of the EEA under the following names:**

Ivor: Italy, Portugal

Zibor: Czech Republic, Estonia, Ireland, Latvia, Lithuania, Poland, Romania, Slovakia,  
Slovenia, United Kingdom

Phivor: Spain

**This leaflet was last revised in November 2025.**