

Package leaflet: Information for the user

Trazodone Hydrochloride 100 mg/5 ml Oral Solution

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Trazodone Oral Solution is and what it is used for
2. What you need to know before you take Trazodone Oral Solution
3. How to take Trazodone Oral Solution
4. Possible side effects
5. How to store Trazodone Oral Solution
6. Contents of the pack and other information

1. What Trazodone Oral Solution is and what it is used for

The name of your medicine is Trazodone 100 mg/5 ml Oral Solution (called Trazodone Oral Solution throughout this leaflet). It contains a medicine called trazodone hydrochloride. This belongs to a group of medicines called antidepressants.

Trazodone Oral Solution can be used to treat anxiety and depression.

2. What you need to know before you take Trazodone Oral Solution

Do not take Trazodone Oral Solution if you:

- are allergic to trazodone hydrochloride or any of the other ingredients of this medicine (listed in section 6).
Signs of an allergic reaction can include: red and lumpy skin rash, swallowing or breathing problems, swelling of your eyelids, lips, face, throat or tongue, unexplained high temperature (fever) and feeling faint
- have recently had a heart attack
- are a heavy drinker or are taking sleeping tablets
- are under 18 years of age.

Do not take this medicine if any of the above applies to you. If you are not sure, talk to your doctor or pharmacist before taking Trazodone Oral Solution.

Warnings and precautions

Take special care with Trazodone Oral Solution

Thoughts of suicide and worsening of your depression or anxiety disorder.

If you are depressed and/or have anxiety disorders you can sometimes have thoughts of harming or killing yourself. These may be increased when first starting antidepressants, since these medicines all take time to work, usually about two weeks but sometimes longer.

You may be more likely to think like this if you:

- have previously had thoughts about killing or harming yourself.
- are a young adult. Information from clinical trials has shown an increased risk of suicidal behaviour in adults aged less than 25 years with psychiatric conditions who were treated with an antidepressant.

If you have thoughts of harming or killing yourself at any time, **contact your doctor or go to a hospital straight away.**

You may find it helpful to tell a relative or close friend that you are depressed or have an anxiety disorder, and ask them to read this leaflet. You might ask them to tell you if they think your depression or anxiety is getting worse, or if they are worried about changes in your behaviour.

If you are elderly, you may be more prone to side effects, increased caution is necessary especially when taking other medicines at the same time as Trazodone Oral Solution or if you have some other diseases.

Talk to your doctor or pharmacist before taking Trazodone Oral Solution if you:

- have or have ever had fits or seizures
- have severe liver, kidney or heart problems
- are pregnant, trying to become pregnant or are breast-feeding
- have an overactive thyroid gland (hyperthyroidism)
- have problems passing water or need to pass water (urine) frequently
- have narrow angle glaucoma (an eye disorder)
- have schizophrenia or other type of mental disorder
- are elderly, as you may be more prone to side effects
- are taking buprenorphine (a strong pain killer). The use of buprenorphine together with Trazodone Oral Solution can lead to serotonin syndrome, a potentially life-threatening condition (see "Other medicines and Trazodone Oral Solution").

If you are not sure if any of the above applies to you, talk to your doctor or pharmacist before taking Trazodone Oral Solution.

Children and adolescents

Trazodone Oral Solution should not be used in children and adolescents under 18 years of age.

Other medicines and Trazodone Oral Solution

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. This includes medicines you buy without prescription,

including herbal medicines. This is because trazodone can affect the way some other medicines work. Also some medicines can affect the way Trazodone Oral Solution works.

Tell your doctor if you are taking any of the following medicines:

- MAOI (Monoamine Oxidase Inhibitors) medicines such as tranylcypromine, phenelzine and isocarboxazid (for depression) or selegiline (for Parkinson's disease), or have taken them in the last 2 weeks
- other antidepressants (such as amitriptyline or fluoxetine)
- **buprenorphine/opioids.** These medicines may interact with Trazodone Oral Solution and you may experience symptoms such as involuntary, rhythmic contractions of muscles, including the muscles that control movement of the eye, agitation, hallucinations, coma, excessive sweating, tremor, exaggeration of reflexes, increased muscle tension, body temperature above 38°C. Contact your doctor when experiencing such symptoms
- sedatives (such as tranquilizers or sleeping pills)
- medicines used to treat epilepsy such as carbamazepine and phenytoin
- medicines used to treat high blood pressure, for example, clonidine
- digoxin (used to treat heart problems)
- medicines used to treat fungal infections such as ketoconazole and itraconazole
- some medicines used to treat HIV such as ritonavir and indinavir
- erythromycin, an antibiotic used to treat infections
- levodopa (used to treat Parkinson's disease)
- St. John's Wort (a herbal remedy)
- warfarin (used to stop your blood from clotting).

Anaesthetics If you are going to have an anaesthetic (for an operation), tell your doctor or dentist that you are taking Trazodone Oral Solution.

Trazodone Oral Solution with food, drink and alcohol

You should avoid drinking alcohol while taking Trazodone Oral Solution.

Pregnancy, breast-feeding and fertility

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

Taking Trazodone Oral Solution in the late stages of pregnancy may lead to your baby experiencing withdrawal symptoms when they are born.

Driving and using machines

Trazodone Oral Solution may make you feel sleepy or dizzy. If this happens do not drive or use any tools or machines.

Important information about some of the ingredients of Trazodone Oral Solution

- **Sorbitol** If your doctor has told you that you have an intolerance to some sugars or if you have been diagnosed with hereditary fructose intolerance (HFI), a rare genetic disorder in which a person cannot break down fructose, talk to your doctor before you take or receive this medicine. If you are given more than 25 ml of this medicine daily, the sorbitol in this medicine may cause gastrointestinal discomfort and have a mild laxative effect.
- **Ethanol** This medicinal product contains small amounts of ethanol (alcohol), less than 100 mg per 30 ml.
- This medicine contains less than 1 mmol sodium (23 mg) per 5 ml, that is to say essentially 'sodium-free'.

3. How to take Trazodone Oral Solution

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

Taking this medicine

- take this medicine by mouth.
- take with or after food. This can help lower the chances of side effects.
- if you have been told to take Trazodone Oral Solution only once each day then you should take it before going to bed.
- if you feel the effect of your medicine is too weak or strong, do not change the dose yourself, but ask your doctor.

How much to take

Adults:

Depression

- Adults usually start by taking 150 mg (7.5 ml) each day.
- For adults in hospital the dose may be as high as 600 mg (30 ml) each day.

Anxiety

- Adults usually start by taking 75 mg (3.75 ml) each day.

Your doctor may increase the dose to 300 mg (15 ml) day depending on your condition.

Elderly:

- Older people or those who are frail will usually be given a starting dose of 100 mg (5 ml) each day.

Doses of more than 300 mg (15 ml) will not normally be given.

Note:

If necessary, Trazodone Oral Solution can be administered via a nasogastric, nasoduodenal or nasojejunal feeding tube that should be rinsed twice with 10 ml of water immediately after administration.

If you take more Trazodone Oral Solution than you should

If you take more Trazodone Oral Solution than you should, tell a doctor or go to a hospital casualty department straight away. Take the medicine pack with you. This is so the doctor knows what you have taken.

The following effects may happen: feeling sick or being sick, feeling sleepy, dizzy or faint, fits (seizures), confusion,

breathing or heart problems.

If you forget to take Trazodone Oral Solution

If you forget to take a dose, take it as soon as you remember it. However, if it is nearly time for the next dose, skip the missed dose.

Do not take a double dose to make up for a forgotten dose.

If you stop taking Trazodone Oral Solution

Keep taking Trazodone Oral Solution until your doctor tells you to stop. Do not stop taking Trazodone Oral Solution just because you feel better.

When your doctor tells you to stop taking this medicine he/she will help you to stop taking it gradually.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Stop taking Trazodone Oral Solution and see a doctor or go to a hospital straight away if you:

- get swelling of the hands, feet, ankles, face, lips or throat which may cause difficulty swallowing or breathing, itching of the skin and nettle rash. This may mean you are having an allergic reaction to Trazodone Oral Solution.
- have painful erection of the penis, unrelated to sexual activity, that will not go away (priapism).
- have yellowing of the eyes or skin. This could be a liver problem (such as jaundice).
- are getting infections more easily than usual. This could be because of a blood disorder (agranulocytosis).
- are bruising more easily than usual. This could be because of a blood disorder (thrombocytopenia).
- have severe abdominal pain and bloating, are being sick (vomiting) and have constipation. These may be signs that your intestine is not working properly (paralytic ileus).

Talk to your doctor straight away if you notice the following side effects:

- you have thoughts of harming or killing yourself.
- feeling tired, faint, dizzy, having pale skin. These could be signs of anaemia.
- convulsions/fits.
- unusual skin sensations such as numbness, tingling, pricking, burning or creeping on the skin (paraesthesia).
- feeling confused, restless, sweating, shaking, shivering, hallucinations (strange visions or sounds), sudden jerks of the muscles or a fast heartbeat, you may have something called Serotonin syndrome.
- feeling very unwell, possibly with shortness of breath (dyspnoea), difficulty in walking or walking with a shuffling gait, shaking, uncontrolled muscle twitching and a high temperature (above 38°C). This could be a rare condition known as Neuroleptic Malignant Syndrome.
- rapid, slow or irregular heartbeat

Below is a list of other side effects that have been reported:

- feeling drowsy or sleepy, tiredness
- feeling less alert than usual
- feeling sick (nausea) or being sick (vomiting), indigestion
- constipation, diarrhoea
- dry mouth, altered taste, increased amounts of saliva, blocked nose
- sweating more than usual
- dizziness, headache, confusion, weakness, tremor (shaking)
- blurred vision
- loss of appetite and weight loss
- feeling dizzy or light-headed on standing or sitting up quickly (postural hypotension), fainting (syncope)
- feeling restless and having difficulty sleeping
- water retention which may cause swollen arms or legs
- skin rash, itching
- chest pain
- pain in limbs, back pain, pain in your muscles, pain in your joints
- jerking movements that you cannot control, mainly in the arms and legs, uncontrolled muscle movements or twitches
- frequent infections with high temperature, severe chills, sore throat or mouth ulcers. These could be signs of a blood problem called leucopenia.
- feeling anxious or more nervous than usual, feeling agitated
- overactive behaviour or thoughts (mania), believing things that are not true (delusions), memory disturbance
- nightmares
- decreased sex drive
- feeling dizzy, possibly with a “spinning” feeling (vertigo)
- high blood pressure
- high temperature
- flu type symptoms
- difficulty with speaking
- higher than normal number of white blood cells (seen by a blood test)
- high levels of liver enzymes in your blood (shown by a blood test)
- severe liver disorders such as hepatitis
- liver failure with potential fatal outcome
- feeling tired, weak and confused, having muscles that ache, are stiff or do not work well. There may also be headache, loss of appetite, nausea or vomiting, convulsion. This may be due to low sodium levels in your blood.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme, website www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Trazodone Oral Solution

Keep this medicine out of the sight and reach of children.

Do not use Trazodone Oral Solution after the expiry date which is stated on the label and carton after EXP. The expiry date refers to the last day of that month.

Keep the bottle in the outer carton in order to protect from light.

After first opening do not store above 25°C and use within 1 month.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

6. Contents of the pack and other information

What Trazadone Oral Solution contains

- Each 5 ml of Trazodone Oral Solution contains 100 mg of the active substance, trazodone hydrochloride.
- Other ingredients include sorbitol solution 70% (E420), glycerol (E422), sodium benzoate (E211), saccharin sodium (E954), hydrochloric acid solution 1.0N, purified water, nitrogen and orange flavour 14104/04 (contains ethanol).

What Trazadone Oral Solution looks like and contents of the pack

Trazodone Oral Solution is a colourless to yellowish clear solution with an orange odour. It is available in an amber glass bottle containing 120 ml of medicine.

The pack also contains a 15 ml plastic measuring cup with intermediate graduations.

Marketing Authorisation Holder

Zentiva Pharma UK Limited
12 New Fetter Lane
London
EC4A 1JP
United Kingdom

Manufacturer

Rafarm SA,
Thesi Pousi-Xatzis,
Agiou Louka,
Paiania Attiki,
19902,
PO Box 37,
Greece

This leaflet was last revised in November 2021.

ZENTIVA
1065028918

ZENTIVA

GENERAL INFO:

PM CODE:	1065028918
PRODUCT NAME:	PIL TRAZODONE HCL 2.4G/120ML BT1 GB
SAP ID / GMID:	921820
AW VERSION:	V6
CREATION DATE:	26.11.2021
AW BY:	TV
SUPPLIER:	RAFARM S.A. (FDP)

REASON OF REVISION:

Change of design from the Creo Style to the Zentiva style.;Zentiva Carton and label colours;Base colour Purple 265;50 mg/5 ml (GMID 920359) Dark Red 201; 100 mg/5 ml (GMID 921820) Dark Green 3308; Leaflet style to be updated including update to wording in line with QRD.;ZN4373 - 74, TRW; Artwork required for submission by the end of the month.

TECHNICAL INFO:

FORMAT (size):	135 x 550 mm
LAETUS (pharma code):	N/A
FONT + MIN. SIZE:	Helvetica Neue LT W1G 9 pt
MATERIAL TYPE (TS):	See VistaLink

COLOURS: [1]

■ Black

TECH. COLOURS: [1]

■ DieCut

ZENTIVA

1065028918