

# **SUMMARY OF PRODUCT CHARACTERISTICS**

## **1 NAME OF THE MEDICINAL PRODUCT**

Sereflo Ciphaler 50 microgram / 250 microgram per dose inhalation powder, predispensed

## **2 QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each single inhalation provides a delivered dose (the dose leaving the mouthpiece) of 47 micrograms of salmeterol (as salmeterol xinafoate) and 231 micrograms of fluticasone propionate. This corresponds to a pre-dispensed dose of 50 micrograms of salmeterol (as salmeterol xinafoate) and 250 micrograms fluticasone propionate.

### Excipients with known effect

Each delivered dose contains up to 11.6 mg of lactose.

For the full list of excipients, see section 6.1.

## **3 PHARMACEUTICAL FORM**

Inhalation powder, pre-dispensed.

Disposable rubine red-white plastic inhaler containing a blister strip with 60 regularly placed blisters. The blister contains a white to off-white powder.

## **4 CLINICAL PARTICULARS**

### **4.1 Therapeutic indications**

Asthma

Sereflo Ciphaler is indicated in the regular treatment of asthma where use of a combination product (long- acting  $\beta_2$  agonist and inhaled corticosteroid) is appropriate:

- patients not adequately controlled with inhaled corticosteroids and 'as needed' inhaled short-acting  $\beta_2$  agonist
- or
- patients already adequately controlled on both inhaled corticosteroid and long-acting  $\beta_2$  agonist

## 4.2 Posology and method of administration

### Posology

Patients should be made aware that Sereflo Ciphaler must be used daily for optimum benefit, even when asymptomatic.

Patients should be regularly reassessed by a doctor, so that the strength of Sereflo Ciphaler they are receiving remains optimal and is only changed on medical advice. **The dose should be titrated to the lowest dose at which effective control of symptoms is maintained. Where the control of symptoms is maintained with the lowest strength of the combination given twice daily then the next step could include a test of inhaled corticosteroid alone.** As an alternative, patients requiring a long- acting  $\beta_2$  agonist could be titrated to Sereflo Ciphaler given once daily if, in the opinion of the prescriber, it would be adequate to maintain disease control. In the event of once daily dosing when the patient has a history of nocturnal symptoms the dose should be given at night and when the patient has a history of mainly daytime symptoms the dose should be given in the morning.

Patients should be given the strength of Sereflo Ciphaler containing the appropriate fluticasone propionate dosage for the severity of their disease. If an individual patient should require dosages outside the recommended regimen, appropriate doses of  $\beta_2$  agonist and/or corticosteroid should be prescribed.

When it is appropriate to prescribe a different strength than is available for Sereflo Ciphaler, a change to an alternative fixed-dose combination of salmeterol and fluticasone propionate containing a different dose of the inhaled corticosteroid, respectively, is required.

### Recommended doses

#### Asthma

*Adults and adolescents 12 years and older*

- One inhalation of 50 micrograms salmeterol and 250 micrograms fluticasone propionate twice daily.

A short-term trial of salmeterol/fluticasone propionate may be considered as initial maintenance therapy in adults or adolescents with moderate persistent asthma (defined as patients with daily symptoms, daily rescue use and moderate to severe airflow limitation) for whom rapid control of asthma is essential. In these cases, the recommended initial dose is one inhalation of 50 micrograms salmeterol and 100 micrograms fluticasone propionate twice daily, a strength which is available for other similar fixed-dose combination products containing these two active ingredients. Once control of asthma is attained treatment should be reviewed and consideration given as to whether patients should be stepped down to an inhaled corticosteroid alone. Regular review of patients as treatment is stepped down is important.

A clear benefit has not been shown as compared to inhaled fluticasone propionate alone used as initial maintenance therapy when one or two of the criteria of severity are missing. In general inhaled corticosteroids remain the first line treatment for most patients. Sereflo Ciphaler is not intended for the initial management of mild asthma. Salmeterol/fluticasone propionate 50 microgram/100 micrograms strength is not appropriate in adults and adolescents with severe asthma; it is recommended to establish the appropriate dosage of inhaled corticosteroid before any fixed-combination can be used in patients with severe asthma.

#### *Paediatric population*

Sereflo Ciphaler is not recommended in children below 12 years of age due to lack of data on safety and efficacy.

#### *Special patient groups*

There is no need to adjust the dose in elderly patients or in those with renal impairment. There are no data available for use of Sereflo Ciphaler in patients with hepatic impairment.

#### Method of administration

For inhalation use.

#### Using the inhaler

1. Patient should hold the inhaler in one hand and place the thumb of other hand in the thumb grip. The patient should then push the thumb grip away as far as it will go until the patient hear a click. This will open a small hole in the mouthpiece.
2. Patient should hold the inhaler with the mouthpiece towards themselves. Patient should slide the lever away from the mouthpiece as far as it will go until it clicks. This places a dose of medicine in the mouthpiece.

3. Every time the lever is pulled back a blister is opened inside and the powder made ready for patient to inhale. Patient should not play with the lever as this opens the blisters and wastes medicine.
4. Before the patient breathes in the dose from the inhaler, the patient should hold the inhaler away from their mouth and breathe out as far as is comfortable. The patient should not breathe into the mouthpiece.
5. The patient should then put the mouthpiece to the lips. The patient should steadily and deeply breathe through the inhaler. The patient should not breathe in through the nose.
6. Patient should remove the inhaler from the mouth and hold the breath for about 10 seconds, or for as long as it is comfortable.
7. Patient should breathe out slowly.
8. Patient should rinse the mouth with water after breathing in the medicine and spit out the water. This may help the patients from getting thrush and becoming hoarse.
9. To close the inhaler, patient should slide the thumbgrip back towards him/her as far as it will go. He/she should make sure that the inhaler clicks. The lever will return to its original position and will reset.
10. The inhaler is now ready for use again.

The counter on top of the inhaler shows how many doses are left. It counts down to 0. The numbers 5 to 0 will appear in red to warn the patient that there will be only few doses left. Once the counter shows 0, the inhaler will be empty.

#### Cleaning your inhaler

The mouthpiece of the Sereflo Ciphaler should be wiped with a dry tissue to clean it.

### **4.3 Contraindications**

Hypersensitivity to the active substances or to any of the excipients listed in section 6.1.

### **4.4 Special warnings and precautions for use**

#### Deterioration of disease

Sereflo Ciphaler should not be used to treat acute asthma symptoms for which a fast- and short- acting bronchodilator is required. Patients should be advised to have their inhaler to be used for relief in an acute asthma attack available at all times.

Patients should not be initiated on Sereflo Ciphaler during an exacerbation, or if they have significantly worsening or acutely deteriorating asthma.

Serious asthma-related adverse events and exacerbations may occur during treatment with Sereflo Ciphaler. Patients should be asked to continue treatment but to seek medical advice if asthma symptoms remain uncontrolled or worsen after initiation on Sereflo Ciphaler.

Increased requirements for use of reliever medication (short-acting bronchodilators), or decreased response to reliever medication indicate deterioration of control and patients should be reviewed by a physician.

Sudden and progressive deterioration in control of asthma is potentially life-threatening and the patient should undergo urgent medical assessment. Consideration should be given to increasing corticosteroid therapy.

Once asthma symptoms are controlled, consideration may be given to gradually reducing the dose of Sereflo Ciphaler. Regular review of patients as treatment is stepped down is important. The lowest effective dose of Sereflo Ciphaler should be used (see section 4.2).

Treatment with Sereflo Ciphaler should not be stopped abruptly in patients with asthma due to risk of exacerbation. Therapy should be down-titrated under physician supervision.

As with all inhaled medication containing corticosteroids, Sereflo Ciphaler should be administered with caution in patients with active or quiescent pulmonary tuberculosis and fungal, viral or other infections of the airway. Appropriate treatment should be promptly instituted, if indicated.

#### Cardiovascular effects

Rarely, Sereflo Ciphaler may cause cardiac arrhythmias e.g. supraventricular tachycardia, extrasystoles and atrial fibrillation, and a mild transient reduction in serum potassium at high therapeutic doses. Sereflo Ciphaler should be used with caution in patients with severe cardiovascular disorders or heart rhythm abnormalities and in patients with diabetes mellitus, thyrotoxicosis, uncorrected hypokalaemia or patients predisposed to low levels of serum potassium.

#### Hyperglycaemia

There have been very rare reports of increases in blood glucose levels (see section 4.8) and this should be considered when prescribing to patients with a history of diabetes mellitus.

#### Paradoxical bronchospasm

As with other inhalation therapy paradoxical bronchospasm may occur with an immediate increase in wheezing and shortness of breath after dosing. Paradoxical bronchospasm responds to a rapid-acting bronchodilator and should be treated straightaway. Sereflo Ciphaler should be discontinued immediately, the patient assessed and alternative therapy instituted if necessary.

The pharmacological side effects of  $\beta_2$  agonist treatment, such as tremor, palpitations and headache, have been reported, but tend to be transient and reduce with regular therapy.

### Systemic corticosteroid effects

Systemic effects may occur with any inhaled corticosteroid, particularly at high doses prescribed for long periods. These effects are much less likely to occur than with oral corticosteroids. Possible systemic effects include Cushing's syndrome, Cushingoid features, adrenal suppression, decrease in bone mineral density, cataract and glaucoma and more rarely, a range of psychological or behavioural effects including psychomotor hyperactivity, sleep disorders, anxiety, depression or aggression (particularly in children) (see *Paediatric population* sub-heading below for information on the systemic effects of inhaled corticosteroids in children and adolescents). **It is important, therefore, that the patient is reviewed regularly and the dose of inhaled corticosteroid is reduced to the lowest dose at which effective control of asthma is maintained.**

Prolonged treatment of patients with high doses of inhaled corticosteroids may result in adrenal suppression and acute adrenal crisis. Very rare cases of adrenal suppression and acute adrenal crisis have also been described with doses of fluticasone propionate between 500 and less than 1000 micrograms. Situations, which could potentially trigger acute adrenal crisis include trauma, surgery, infection or any rapid reduction in dosage. Presenting symptoms are typically vague and may include anorexia, abdominal pain, weight loss, tiredness, headache, nausea, vomiting, hypotension, decreased level of consciousness, hypoglycaemia, and seizures. Additional systemic corticosteroid cover should be considered during periods of stress or elective surgery.

The benefits of inhaled fluticasone propionate therapy should minimise the need for oral steroids, but patients transferring from oral steroids may remain at risk of impaired adrenal reserve for a considerable time. Therefore these patients should be treated with special care and adrenocortical function regularly monitored. Patients who have required high dose emergency corticosteroid therapy in the past may also be at risk. This possibility of residual impairment should always be borne in mind in emergency and elective situations likely to produce stress, and appropriate corticosteroid treatment must be considered. The extent of the adrenal impairment may require specialist advice before elective procedures.

Ritonavir can greatly increase the concentration of fluticasone propionate in plasma. Therefore, concomitant use should be avoided, unless the potential benefit to the patient outweighs the risk of systemic corticosteroid side effects. There is also an

increased risk of systemic side effects when combining fluticasone propionate with other potent CYP3A inhibitors (see section 4.5).

#### Interactions with potent CYP3A4 inhibitors

Concomitant use of systemic ketoconazole significantly increases systemic exposure to salmeterol. This may lead to an increase in the incidence of systemic effects (e.g. prolongation in the QTc interval and palpitations). Concomitant treatment with ketoconazole or other potent CYP3A4 inhibitors should therefore be avoided unless the benefits outweigh the potentially increased risk of systemic side effects of salmeterol treatment (see section 4.5).

#### Visual disturbance

Visual disturbance may be reported with systemic and topical corticosteroid use. If a patient presents with symptoms such as blurred vision or other visual disturbances, the patient should be considered for referral to an ophthalmologist for evaluation of possible causes, which may include cataract, glaucoma or rare diseases such as central serous chorioretinopathy (CSCR) which have been reported after use of systemic and topical corticosteroids.

#### Paediatric population

Children and adolescents <16 years taking high doses of fluticasone propionate (typically  $\geq 1000$  micrograms/day) may be at particular risk. Systemic effects may occur, particularly at high doses prescribed for long periods. Possible systemic effects include Cushing's syndrome, Cushingoid features, adrenal suppression, acute adrenal crisis and growth retardation in children and adolescents and more rarely, a range of psychological or behavioural effects including psychomotor hyperactivity, sleep disorders, anxiety, depression or aggression. Consideration should be given to referring the child or adolescent to a paediatric respiratory specialist.

It is recommended that the height of children receiving prolonged treatment with inhaled corticosteroid is regularly monitored. **The dose of inhaled corticosteroid should be reduced to the lowest dose at which effective control of asthma is maintained.**

#### Lactose intolerance

This medicinal product contains lactose. Patients with rare hereditary problems of galactose intolerance, total lactase deficiency or glucose-galactose malabsorption should not take this medicine. The excipient lactose contains small amounts of milk proteins, which may cause allergic reactions.

#### 4.5 Interaction with other medicinal products and other forms of interaction

$\beta$  adrenergic blockers may weaken or antagonise the effect of salmeterol. Both non-selective and selective  $\beta$  blockers should be avoided unless there are compelling reasons for their use. Potentially serious hypokalaemia may result from  $\beta_2$  agonist therapy. Particular caution is advised in acute severe asthma as this effect may be potentiated by concomitant treatment with xanthine derivatives, steroids and diuretics.

Concomitant use of other  $\beta$  adrenergic containing drugs can have a potentially additive effect.

##### Fluticasone propionate

Under normal circumstances, low plasma concentrations of fluticasone propionate are achieved after inhaled dosing, due to extensive first pass metabolism and high systemic clearance mediated by cytochrome CYP3A4 in the gut and liver. Hence, clinically significant drug interactions mediated by fluticasone propionate are unlikely.

In an interaction study in healthy subjects with intranasal fluticasone propionate, ritonavir (a highly potent cytochrome CYP3A4 inhibitor) 100 mg b.i.d. increased the fluticasone propionate plasma concentrations several hundred fold, resulting in markedly reduced serum cortisol concentrations. Information about this interaction is lacking for inhaled fluticasone propionate, but a marked increase in fluticasone propionate plasma levels is expected. Cases of Cushing's syndrome and adrenal suppression have been reported. The combination should be avoided unless the benefit outweighs the increased risk of systemic glucocorticoid side effects.

In a small study in healthy volunteers, the slightly less potent CYP3A inhibitor ketoconazole increased the exposure of fluticasone propionate after a single inhalation by 150%. This resulted in a greater reduction of plasma cortisol as compared with fluticasone propionate alone. Co-treatment with other potent CYP3A inhibitors, such as itraconazole and cobicistat-containing products, and moderate CYP3A inhibitors, such as erythromycin, is also expected to increase the systemic fluticasone propionate exposure and the risk of systemic side effects. Combinations should be avoided unless the benefit outweighs the potential increased risk of systemic corticosteroid side-effects, in which case patients should be monitored for systemic corticosteroid side-effects.

##### Salmeterol

###### Potent CYP3A4 inhibitors

Co-administration of ketoconazole (400 mg orally once daily) and salmeterol (50 micrograms inhaled twice daily) in 15 healthy subjects for 7 days resulted in a significant increase in plasma salmeterol exposure (1.4-fold  $C_{max}$  and 15-fold AUC). This may lead to an increase in the incidence of other systemic

effects of salmeterol treatment (e.g. prolongation of QTc interval and palpitations) compared with salmeterol or ketoconazole treatment alone (see section 4.4).

Clinically significant effects were not seen on blood pressure, heart rate, blood glucose and blood potassium levels. Co-administration with ketoconazole did not increase the elimination half-life of salmeterol or increase salmeterol accumulation with repeat dosing.

The concomitant administration of ketoconazole should be avoided, unless the benefits outweigh the potentially increased risk of systemic side effects of salmeterol treatment. There is likely to be a similar risk of interaction with other potent CYP3A4 inhibitors (e.g. itraconazole, telithromycin, ritonavir).

#### Moderate CYP 3A4 inhibitors

Co-administration of erythromycin (500 mg orally three times a day) and salmeterol (50 micrograms inhaled twice daily) in 15 healthy subjects for 6 days resulted in a small but non-statistically significant increase in salmeterol exposure (1.4-fold  $C_{max}$  and 1.2-fold AUC). Co-administration with erythromycin was not associated with any serious adverse effects.

## **4.6 Fertility, Pregnancy and lactation**

### Fertility

There are no data in humans. However, animal studies showed no effects of salmeterol or fluticasone propionate on fertility.

### Pregnancy

A large amount of data on pregnant women (more than 1000 pregnancy outcomes) indicates no malformative or feto/neonatal toxicity related to Sereflo Ciphaler. Animal studies have shown reproductive toxicity after administration of  $\beta_2$  adrenoreceptor agonists and glucocorticosteroids (see section 5.3).

Administration of Sereflo Ciphaler to pregnant women should only be considered if the expected benefit to the mother is greater than any possible risk to the fetus.

The lowest effective dose of fluticasone propionate needed to maintain adequate asthma control should be used in the treatment of pregnant women.

### Breastfeeding

It is unknown whether salmeterol and fluticasone propionate/metabolites are excreted in human milk.

Studies have shown that salmeterol and fluticasone propionate, and their metabolites, are excreted into the milk of lactating rats.

A risk to breastfed newborns/infants cannot be excluded. A decision must be made whether to discontinue breastfeeding or to discontinue Sereflo Ciphaler therapy taking into account the benefit of breastfeeding for the child and the benefit of therapy for the woman.

#### 4.7 Effects on ability to drive and use machines

Sereflo Ciphaler has no or negligible influence on the ability to drive and use machines.

#### 4.8 Undesirable effects

As Sereflo Ciphaler contains salmeterol and fluticasone propionate, the type and severity of adverse reactions associated with each of the compounds may be expected. There is no incidence of additional adverse events following concurrent administration of the two compounds.

Adverse events which have been associated with salmeterol/fluticasone propionate are given below, listed by system organ class and frequency. Frequencies are defined as: very common ( $\geq 1/10$ ), common ( $\geq 1/100$  to  $< 1/10$ ), uncommon ( $\geq 1/1000$  to  $< 1/100$ ), rare ( $\geq 1/10,000$  to  $< 1/1000$ ) and not known (cannot be estimated from the available data). Frequencies were derived from clinical trial data. The incidence in placebo was not taken into account.

System Organ Class	Adverse Event	Frequency
Infections and Infestations	Candidiasis of the mouth and throat	Common
	Bronchitis	Common <sup>1, 3</sup>
	Oesophageal candidiasis	Rare
Immune System Disorders	Hypersensitivity reactions with the following manifestations:	
	Cutaneous hypersensitivity reactions	Uncommon
	Angioedema (mainly facial and oropharyngeal oedema)	Rare
	Respiratory symptoms (dyspnoea)	Uncommon
	Respiratory symptoms (bronchospasm)	Rare

	Anaphylactic reactions including anaphylactic shock	Rare
Endocrine Disorders	Cushing's syndrome, Cushingoid features, Adrenal suppression, Growth retardation in children and adolescents, Decreased bone mineral density	Rare <sup>4</sup>
Metabolism and Nutrition Disorders	Hypokalaemia Hyperglycaemia	Common <sup>3</sup> Uncommon <sup>4</sup>
Psychiatric Disorders	Anxiety Sleep disorders Behavioural changes, including psychomotor hyperactivity and irritability (predominantly in children) Depression, aggression (predominantly in children)	Uncommon Uncommon Rare Not Known
Nervous System Disorders	Headache Tremor	Very Common <sup>1</sup> Uncommon
Eye Disorders	Cataract Glaucoma Vision, blurred	Uncommon Rare <sup>4</sup> Not Known <sup>4</sup>
Cardiac Disorders	Palpitations Tachycardia Cardiac arrhythmias (including supraventricular tachycardia and extrasystoles). Atrial fibrillation Angina pectoris	Uncommon Uncommon Rare Uncommon Uncommon
Respiratory, Thoracic and Mediastinal Disorders	Nasopharyngitis Throat irritation Hoarseness/dysphonia Sinusitis Paradoxical bronchospasm	Very Common <sup>2,3</sup> Common Common Common <sup>1,3</sup> Rare <sup>4</sup>
Skin and Subcutaneous Tissue Disorders	Contusions	Common <sup>1,3</sup>
Musculoskeletal & Connective Tissue Disorders	Muscle cramps Traumatic fractures Arthralgia Myalgia	Common Common <sup>1,3</sup> Common Common

1. Reported commonly in placebo
2. Reported very commonly in placebo
3. Reported over 3 years in a COPD study
4. See section 4.4

#### Description of selected adverse reactions

The pharmacological side effects of  $\beta_2$  agonist treatment, such as tremor, palpitations and headache, have been reported, but tend to be transient and reduce with regular therapy.

As with other inhalation therapy paradoxical bronchospasm may occur with an immediate increase in wheezing and shortness of breath after dosing. Paradoxical bronchospasm responds to a rapid-acting bronchodilator and should be treated straightaway. Sereflo Ciphaler should be discontinued immediately, the patient assessed and alternative therapy instituted if necessary.

Due to the fluticasone propionate component, hoarseness and candidiasis (thrush) of the mouth and throat and, rarely, of the oesophagus can occur in some patients. Both hoarseness and incidence of mouth and throat candidiasis may be relieved by rinsing the mouth with water and/or brushing the teeth after using the product. Symptomatic mouth and throat candidiasis can be treated with topical anti-fungal therapy whilst still continuing with the Sereflo Ciphaler.

#### Paediatric population

Possible systemic effects include Cushing's syndrome, Cushingoid features, adrenal suppression and growth retardation in children and adolescents (see section 4.4). Children may also experience anxiety, sleep disorders and behavioural changes, including hyperactivity and irritability.

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the Yellow Card Scheme at: <https://yellowcard.mhra.gov.uk/> or search for MHRA Yellow Card in the Google Play or Apple App Store.

## **4.9 Overdose**

### Symptoms

There are no data available from clinical trials on overdose with Sereflo Ciphaler, however data on overdose with both drugs are given below:

The signs and symptoms of salmeterol overdose are dizziness, increases in systolic blood pressure, tremor, headache and tachycardia. If Sereflo Ciphaler therapy has to be withdrawn due to overdose of the  $\beta$  agonist component of the drug, provision of appropriate replacement steroid therapy should be considered. Additionally, hypokalaemia can occur and therefore serum potassium levels should be monitored. Potassium replacement should be considered.

### Acute

Acute inhalation of fluticasone propionate doses in excess of those recommended may lead to temporary suppression of adrenal function. This does not need emergency action as adrenal function is recovered in a few days, as verified by plasma cortisol measurements.

### Chronic overdose of inhaled fluticasone propionate

Adrenal reserve should be monitored and treatment with a systemic corticosteroid may be necessary. When stabilised, treatment should be continued with an inhaled corticosteroid at the recommended dose. Refer to section 4.4: risk of adrenal suppression.

### Management

In cases of both acute and chronic fluticasone propionate overdose, Sereflo Ciphaler therapy should be continued at a suitable dosage for symptom control.

## **5 PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Adrenergics in combination with corticosteroids or other drugs, excl. Anticholinergics.

ATC Code: R03AK06

#### Mechanism of action and pharmacodynamic effects

Sereflo Ciphaler contains salmeterol and fluticasone propionate which have differing modes of action. The respective mechanisms of action of both drugs are discussed below.

#### *Salmeterol*

Salmeterol is a selective long-acting (12 hour)  $\beta_2$  adrenoceptor agonist with a long side chain which binds to the exo-site of the receptor.

Salmeterol produces a longer duration of bronchodilation, lasting for at least 12 hours, than recommended doses of conventional short-acting  $\beta_2$  agonists.

#### *Fluticasone propionate*

Fluticasone propionate given by inhalation at recommended doses has a glucocorticoid anti-inflammatory action within the lungs, resulting in reduced

symptoms and exacerbations of asthma, with less adverse effects than when corticosteroids are administered systemically.

Clinical efficacy and safety

Salmeterol and fluticasone propionate inhalation powder asthma clinical trials

A twelve month study (Gaining Optimal Asthma Control, GOAL), in 3416 adult and adolescent patients with persistent asthma, compared the safety and efficacy of salmeterol and fluticasone propionate inhalation powder versus inhaled corticosteroid (Fluticasone Propionate) alone to determine whether the goals of asthma management were achievable. Treatment was stepped up every 12 weeks until **\*\*total control** was achieved or the highest dose of study drug was reached. GOAL showed more patients treated with salmeterol and fluticasone propionate inhalation powder achieved asthma control than patients treated with ICS alone and this control was attained at a lower corticosteroid dose.

*\*Well controlled* asthma was achieved more rapidly with salmeterol and fluticasone propionate inhalation powder than with ICS alone. The time on treatment for 50% of subjects to achieve a first individual *well controlled* week was 16 days for salmeterol and fluticasone propionate inhalation powder compared to 37 days for the ICS group. In the subset of steroid naive asthmatics the time to an individual *well controlled* week was 16 days in the salmeterol and fluticasone propionate inhalation powder treatment compared to 23 days following treatment with ICS.

The overall study results showed:

<b>Percentage of Patients Attaining *Well Controlled (WC) and **Totally Controlled (TC) Asthma over 12 months</b>				
<b>Pre-Study Treatment</b>	<b>Salmeterol/FP</b>		<b>FP</b>	
	<b>WC</b>	<b>TC</b>	<b>WC</b>	<b>TC</b>
<b>No ICS (SABA alone)</b>	78%	50%	70%	40%
<b>Low dose ICS (<math>\leq</math>500 micrograms BDP or equivalent/day)</b>	75%	44%	60%	28%
<b>Medium dose ICS (<math>&gt;</math>500 to 1000 micrograms BDP or equivalent/day)</b>	62%	29%	47%	16%
<b>Pooled results across the 3 treatment levels</b>	71%	41%	59%	28%

\*Well controlled asthma; less than or equal to 2 days with symptom score greater than 1 (symptom score 1 defined as 'symptoms for one short period during the day'), SABA use on less than or equal to 2 days and less than or equal to 4 occasions/week, greater than or equal to 80% predicted morning peak expiratory flow, no night-time awakenings, no exacerbations and no side effects enforcing a change in therapy

\*\*Total control of asthma; no symptoms, no SABA use, greater than or equal to 80% predicted morning peak expiratory flow, no night-time awakenings, no exacerbations and no side effects enforcing a change in therapy

The results of this study suggest that salmeterol and fluticasone propionate inhalation powder 50/100 micrograms bd may be considered as initial maintenance therapy in patients with moderate persistent asthma for whom rapid control of asthma is deemed essential (see section 4.2).

A double blind, randomised, parallel group study in 318 patients with persistent asthma aged  $\geq 18$  years evaluated the safety and tolerability of administering two inhalations twice daily (double dose) of salmeterol and fluticasone propionate inhalation powder for two weeks. The study showed that doubling the inhalations of each strength of salmeterol and fluticasone propionate inhalation powder for up to 14 days resulted in a small increase in  $\beta$  agonist-related adverse events (tremor; 1 patient [1%] vs 0, palpitations; 6 [3%] vs 1 [ $<1\%$ ], muscle cramps; 6[3%] vs 1 [ $<1\%$ ]) and a similar incidence of inhaled corticosteroid-related adverse events (e.g. oral candidiasis; 6 [6%] vs 16 [8%], hoarseness; 2 [2%] vs 4 [2%]) compared to one inhalation twice daily. The small increase in  $\beta$  agonist-related adverse events should be taken into account if doubling the dose of salmeterol and fluticasone propionate inhalation powder is considered by the physician in adult patients requiring additional short-term (up to 14 days) inhaled corticosteroid therapy.

### Asthma

#### *The Salmeterol Multi-center Asthma Research Trial (SMART)*

The Salmeterol Multi-center Asthma Research Trial (SMART) was a 28-week US study that evaluated the safety of salmeterol compared to placebo added to usual therapy in adult and adolescent subjects. Although there were no significant differences in the primary endpoint of the combined number of respiratory-related deaths and respiratory-related life-threatening experiences, the study showed a significant increase in asthma-related deaths in patients receiving salmeterol (13 deaths out of 13,176 patients treated with salmeterol versus 3 deaths out of 13,179 patients on placebo). The study was not designed to assess the impact of concurrent inhaled corticosteroid use, and only 47% of subjects reported ICS use at baseline.

#### Safety and efficacy of salmeterol-FP versus FP alone in asthma

Two multi-centre 26-week studies were conducted to compare the safety and efficacy of salmeterol-FP versus FP alone, one in adult and adolescent subjects (AUSTRI trial), and the other in paediatric subjects 4-11 years of age (VESTRI trial). For both studies, enrolled subjects had moderate to severe persistent asthma with history of asthma-related hospitalisation or asthma exacerbation in the previous year. The primary objective of each study was to determine whether the addition of LABA to ICS therapy (salmeterol-FP) was non-inferior to ICS (FP) alone in terms of the risk of serious asthma related events (asthma-related hospitalisation, endotracheal intubation, and death). A secondary efficacy objective of these studies was to evaluate whether

ICS/LABA (salmeterol-FP) was superior to ICS therapy alone (FP) in terms of severe asthma exacerbation (defined as deterioration of asthma requiring the use of systemic corticosteroids for at least 3 days or an in-patient hospitalisation or emergency department visit due to asthma that required systemic corticosteroids).

A total of 11,679 and 6,208 subjects were randomized and received treatment in the AUSTRI and VESTRI trials, respectively. For the primary safety endpoint, non-inferiority was achieved for both trials (see Table below).

***Serious Asthma-Related Events in the 26-Week AUSTRI and VESTRI Trials***

	AUSTRI		VESTRI	
	Salmeterol-FP (n = 5,834)	FP Alone (n = 5,845)	Salmeterol-FP (n = 3,107)	FP Alone (n = 3,101)
Composite endpoint (Asthma-related hospitalisation, endotracheal intubation, or death)	34 (0.6%)	33 (0.6%)	27 (0.9%)	21 (0.7%)
Salmeterol-FP/FP Hazard ratio (95% CI)	1.029 (0.638-1.662) <sup>a</sup>		1.285 (0.726-2.272) <sup>b</sup>	
Death	0	0	0	0
Asthma-related hospitalisation	34	33	27	21
Endotracheal intubation	0	2	0	0

<sup>a</sup>. If the resulting upper 95% CI estimate for the relative risk was less than 2.0, then non-inferiority was concluded.

<sup>b</sup>. If the resulting upper 95% CI estimate for the relative risk was less than 2.675, then non-inferiority was concluded.

For the secondary efficacy endpoint, reduction in time to first asthma exacerbation for salmeterol-FP relative to FP was seen in both studies, however only AUSTRI met statistical significance:

	AUSTRI		VESTRI	
	Salmeterol-FP (n = 5,834)	FP Alone (n = 5,845)	Salmeterol-FP (n = 3,107)	FP Alone (n = 3,101)
Number of subjects with an asthma exacerbation	480 (8%)	597 (10%)	265 (9%)	309 (10%)
Salmeterol-FP/FP Hazard ratio (95% CI)	0.787 (0.698, 0.888)		0.859 (0.729, 1.012)	

#### Paediatric population

In trial SAM101667, in 158 children aged 6 to 16 years with symptomatic asthma, the combination of salmeterol/fluticasone propionate is equally efficacious to doubling the dose of fluticasone propionate regarding symptom control and lung function. This study was not designed to investigate the effect on exacerbations.

In a 12 week trial of children aged 4 to 11 years [n=257] treated with either salmeterol/fluticasone propionate 50/100 or salmeterol 50 micrograms + fluticasone propionate 100 micrograms both twice daily, both treatment arms experienced a 14% increase in peak expiratory flow rate as well as improvements in symptom score and rescue salbutamol use. There were no differences between the 2 treatment arms. There were no differences in safety parameters between the 2 treatment arms.

In a 12 week trial of children 4 to 11 years of age [n=203] randomized in a parallel-group study with persistent asthma and who were symptomatic on inhaled corticosteroid, safety was the primary objective. Children received either salmeterol/fluticasone propionate (50/100 micrograms) or fluticasone propionate (100 micrograms) alone twice daily. Two children on salmeterol/fluticasone propionate and 5 children on fluticasone propionate withdrew because of worsening asthma. After 12 weeks no children in either treatment arm had abnormally low 24 hour urinary cortisol excretion. There were no other differences in safety profile between the treatment arms.

#### Fluticasone propionate containing medications in asthma during pregnancy

An observational retrospective epidemiological cohort study utilising electronic health records from the United Kingdom was conducted to evaluate the risk of MCMs following first trimester exposure to inhaled FP alone and salmeterol-FP relative to non-FP containing ICS. No placebo comparator was included in this study.

Within the asthma cohort of 5362 first trimester ICS-exposed pregnancies, 131 diagnosed MCMs were identified; 1612 (30%) were exposed to FP or salmeterol-FP of which 42 diagnosed MCMs were identified. The adjusted odds ratio for MCMs diagnosed by 1 year was 1.1 (95% CI: 0.5 – 2.3) for FP exposed vs non-FP ICS exposed women with moderate asthma and 1.2 (95% CI: 0.7 – 2.0) for women with considerable to severe asthma. No difference in the risk of MCMs was identified following first trimester exposure to FP alone versus salmeterol-FP. Absolute risks of MCM across the asthma severity strata ranged from 2.0 to 2.9 per 100 FP-exposed pregnancies which is comparable to results from a study of 15,840 pregnancies unexposed to asthma therapies in the General Practice Research Database (2.8 MCM events per 100 pregnancies).

## 5.2 Pharmacokinetic properties

For pharmacokinetic purposes each component can be considered separately.

### Salmeterol

Salmeterol acts locally in the lung therefore plasma levels are not an indication of therapeutic effects. In addition there are only limited data available on the pharmacokinetics of salmeterol because of the technical difficulty of assaying the drug in plasma due to the low plasma concentrations at therapeutic doses (approximately 200 picogram /mL or less) achieved after inhaled dosing.

### Fluticasone propionate

The absolute bioavailability of a single dose of inhaled fluticasone propionate in healthy subjects varies between approximately 5 to 11% of the nominal dose depending on the inhalation device used. In patients with asthma or COPD a lesser degree of systemic exposure to inhaled fluticasone propionate has been observed.

Systemic absorption occurs mainly through the lungs and is initially rapid then prolonged. The remainder of the inhaled dose may be swallowed but contributes minimally to systemic exposure due to the low aqueous solubility and pre-systemic metabolism, resulting in oral availability of less than 1%. There is a linear increase in systemic exposure with increasing inhaled dose.

The disposition of fluticasone propionate is characterised by high plasma clearance (1150 mL/min), a large volume of distribution at steady-state (approximately 300 L) and a terminal half-life of approximately 8 hours.

Plasma protein binding is 91%.

Fluticasone propionate is cleared very rapidly from the systemic circulation. The main pathway is metabolism to an inactive carboxylic acid metabolite, by the cytochrome P450 enzyme CYP3A4. Other unidentified metabolites are also found in the faeces.

The renal clearance of fluticasone propionate is negligible. Less than 5% of the dose is excreted in urine, mainly as metabolites. The main part of the dose is excreted in faeces as metabolites and unchanged drug.

### Paediatric population

In a population pharmacokinetic analysis utilising data from 9 controlled clinical trials with different devices (Diskus metered dose inhaler) that included 350 patients with asthma aged 4 to 77 years (174 patients 4 to 11 years of age) higher fluticasone propionate systemic exposure following treatment with salmeterol and fluticasone propionate Diskus 50/100 compared to fluticasone propionate Diskus 100 were seen.

Geometric Mean Ratio [90% CI] for the Salmeterol/fluticasone propionate vs. fluticasone propionate Diskus Comparison in Children and Adolescent/Adult Populations

<b><i>Treatment (test vs. ref)</i></b>	<b><i>Population</i></b>	<b><i>AUC</i></b>	<b><i>C<sub>max</sub></i></b>
<i>Salmeterol/fluticasone propionate Diskus 50/100 fluticasone propionate Diskus 100</i>	<i>Children (4–11yr)</i>	<i>1.20 [1.06 – 1.37]</i>	<i>1.25 [1.11 – 1.41]</i>
<i>Salmeterol/fluticasone propionate Diskus 50/100 fluticasone propionate Diskus 100</i>	<i>Adolescent/Adult (≥12yr)</i>	<i>1.52 [1.08 – 2.13]</i>	<i>1.52 [1.08 – 2.16]</i>

The effect of 21 days of treatment with salmeterol and fluticasone propionate Inhaler 25/50 micrograms (2 inhalations twice daily with or without a spacer) or salmeterol and fluticasone propionate Diskus 50/100 micrograms (1 inhalation twice daily) was evaluated in 31 children aged 4 to 11 years with mild asthma. Systemic exposure to salmeterol was similar for salmeterol and fluticasone propionate Inhaler, salmeterol and fluticasone propionate Inhaler with spacer, and salmeterol and fluticasone propionate Diskus (126 pg hr/mL [95% CI: 70, 225], 103 pg hr/mL [95% CI: 54, 200], and 110 pg hr/mL [95% CI: 55, 219], respectively). Systemic exposure to fluticasone propionate was similar for salmeterol and fluticasone propionate Inhaler with spacer (107 pg hr/mL [95% CI: 45.7, 252.2]) and salmeterol and fluticasone propionate Diskus (138 pg hr/mL [95% CI: 69.3, 273.2]), but lower for salmeterol and fluticasone propionate Inhaler (24 pg hr/mL [95% CI: 9.6, 60.2]).

### **5.3 Preclinical safety data**

The only safety concerns for human use derived from animal studies of salmeterol and fluticasone propionate given separately were effects associated with exaggerated pharmacological actions.

In animal reproduction studies, glucocorticosteroids have been shown to induce malformations (cleft palate, skeletal malformations). However, these animal experimental results do not seem to be relevant for man given recommended doses. Animal studies with salmeterol have shown embryofetal toxicity only at high exposure levels. Following co-administration, increased incidences of transposed umbilical artery and incomplete ossification of occipital bone were found in rats at doses associated with known glucocorticoid-induced abnormalities. Neither salmeterol xinafoate or fluticasone propionate have shown any potential for genetic toxicity.

## **6 PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Lactose monohydrate.

### **6.2 Incompatibilities**

Not applicable.

### **6.3 Shelf life**

2 years.

Use within 1 month of opening the foil pouch.

#### **6.4 Special precautions for storage**

This medicinal product does not require any special storage conditions. Do not refrigerate or freeze.

Do not store the opened product above 25°C.

#### **6.5 Nature and contents of container**

Sereflo Ciphaler are supplied as a disposable rubine red-white plastic inhaler containing a blister strip with 60 blisters. The blister consists of 3 layered forming foil (OPA/ALU/PVC) and a lidding peelable foil (PAPER / PET/ALU/HSL). The inhaler is packaged in a laminated aluminium foil pouch. Laminated aluminium foil pouch consists of 4 layers (PET / PE /Alu Foil / PE).

#### **6.6 Special precautions for disposal**

The inhaler releases a powder which is inhaled into the lungs. A dose indicator on the inhaler indicates the number of doses left. For detailed instructions for use see the Patient Information Leaflet.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

### **7 MARKETING AUTHORISATION HOLDER**

Cipla (EU) Limited  
Dixcart House  
Addlestone Road,  
Bourne Business Park  
Addlestone, Surrey, KT152LE,  
United Kingdom

### **8 MARKETING AUTHORISATION NUMBER(S)**

PLGB 36390/0355

**9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

11/11/2025

**10 DATE OF REVISION OF THE TEXT**

11/11/2025