

Brivaracetam Stirling 10 mg film-coated tablets

Brivaracetam Stirling 25 mg film-coated tablets

Brivaracetam Stirling 50 mg film-coated tablets

Brivaracetam Stirling 75 mg film-coated tablets

Brivaracetam Stirling 100 mg film-coated tablets

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

- 1 What Brivaracetam Stirling film-coated tablets are and what they are used for
- 2 What you need to know before you take Brivaracetam Stirling film-coated tablets
- 3 How to take Brivaracetam Stirling film-coated tablets
- 4 Possible side effects
- 5 How to store Brivaracetam Stirling film-coated tablets
- 6 Contents of the pack and other information

1 WHAT BRIVARACETAM STIRLING FILM-COATED TABLETS ARE AND WHAT THEY ARE USED FOR

What Brivaracetam Stirling film-coated tablets are

Brivaracetam Stirling film-coated tablets contain the active substance brivaracetam. It belongs to a group of medicines called 'anti-epileptics'. These medicines are used to treat epilepsy.

What Brivaracetam Stirling film-coated tablets are used for

- Brivaracetam Stirling film-coated tablets are used in adults, adolescents and children from 2 years of age.
- They are used to treat a type of epilepsy that has partial seizures with or without a secondary generalisation.
- Partial seizures are fits that start by only affecting one side of the brain. These partial seizures can spread and extend to larger areas on both sides of the brain – this is called a 'secondary generalisation'.
- You have been given this medicine to lower the number of fits (seizures) you have.
- Brivaracetam Stirling film-coated tablets are used together with other medicines for epilepsy.

2 WHAT YOU NEED TO KNOW BEFORE YOU TAKE BRIVARACETAM STIRLING FILM-COATED TABLETS

Do not take Brivaracetam Stirling film-coated tablets if:

- you are allergic to brivaracetam, other similar chemical compounds as levetiracetam or piracetam or any of the other ingredients of this medicine (listed in section 6). If you are not sure, talk to your doctor or pharmacist before taking Brivaracetam Stirling film-coated tablets.
- you have ever developed a severe skin rash or skin peeling, blistering and/or mouth sores after taking Brivaracetam Stirling film-coated tablets.

- serious skin reactions including Stevens-Johnson syndrome have been reported in association with Brivaracetam Stirling film-coated tablets treatment. Stop using Brivaracetam Stirling film-coated tablets and seek medical attention immediately if you notice any of the symptoms related to these serious skin reactions described in section 4.

Warnings and precautions

Talk to your doctor or pharmacist before taking Brivaracetam Stirling film-coated tablets if:

- You have thoughts of harming or killing yourself. A small number of people being treated with anti-epileptic medicines such as Brivaracetam Stirling film-coated tablets have had thoughts of harming or killing themselves. If you have any of these thoughts at any time, contact your doctor immediately.
- You have liver problems – your doctor may need to adjust your dose.

Children

Brivaracetam Stirling film-coated tablets are not recommended for use in children under 2 years of age.

Other medicines and Brivaracetam Stirling film-coated tablets

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

In particular, tell your doctor if you are taking any of the following medicines – this is because your doctor may need to adjust your Brivaracetam Stirling film-coated tablets dose:

- Rifampicin – a medicine used to treat bacterial infections.
- St John's wort (also known as Hypericum perforatum) – a herbal medicine used to treat depression and anxiety as well as other conditions.

Brivaracetam Stirling film-coated tablets with alcohol

- Combining this medicine with alcohol is not recommended.
- If you drink alcohol while taking Brivaracetam Stirling film-coated tablets the negative effects of alcohol may be increased.

Pregnancy and breast-feeding

Fertile women should discuss the use of contraceptives with the doctor.

If you are pregnant or breast-feeding, think you may be pregnant or planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

It is not recommended to take Brivaracetam Stirling film-coated tablets if you are pregnant, as the effects of brivaracetam on pregnancy and the unborn baby are not known.

It is not recommended to breast-feed your baby while taking Brivaracetam Stirling film-coated tablets, as brivaracetam passes into breast milk.

Do not stop treatment without talking to your doctor first. Stopping treatment could increase your seizures and harm your baby.

Driving and using machines

- You may feel sleepy, dizzy or tired while taking Brivaracetam Stirling film-coated tablets.
- These effects are more likely at the start of the treatment or after a dose increase.
- Do not drive, cycle or use any tools or machines until you know how the medicine affects you.

Brivaracetam Stirling film-coated tablets contain lactose and sodium

Brivaracetam Stirling film-coated tablets contain:

- lactose (a type of sugar) – if you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.
- sodium – this medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium free'.

3 HOW TO TAKE BRIVARACETAM STIRLING FILM-COATED TABLETS

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure. Other form(s) of this medicine may be more suitable for certain patients e.g. children (if tablets cannot be swallowed in whole, for example); ask your doctor or pharmacist.

You will take Brivaracetam Stirling film-coated tablets together with other medicines for epilepsy.

How much to take

Your doctor will work out the right daily dose for you. Take the daily dose in two equal divided doses, approximately 12 hours apart.

Adolescents and children weighing 50 kg or more, and adults

The recommended dose is from 25 mg to 100 mg taken twice a day. Your doctor may then decide to adjust your dose to find the best dose for you.

Adolescents and children weighing from 20 kg to less than 50 kg

The recommended dose is from 0.5 mg to 2 mg for each kg of bodyweight, taken twice a day. Your doctor may then decide to adjust your dose to find the best dose for you.

Children weighing from 10 kg to less than 20 kg

The recommended dose is from 0.5 mg to 2.5 mg for each kg of bodyweight, taken twice a day. Your child's doctor may then decide to adjust your child's dose to find the best dose for your child.



People with liver problems

If you have problems with your liver:

- As an adolescent or child weighing 50 kg or more, or as an adult, the maximum dose you will take is 75 mg twice a day.
- As an adolescent or child weighing from 20 kg to less than 50 kg, the maximum dose you will take is 1.5 mg for each kg of bodyweight twice a day.
- As a child weighing from 10 kg to less than 20 kg, the maximum dose your child will take is 2 mg for each kg of bodyweight twice a day.

How to take Brivaracetam Stirling film-coated tablets

- Swallow the tablets whole with a glass of liquid.

The medicine may be taken with or without food.

How long to take Brivaracetam Stirling film-coated tablets for

Brivaracetam Stirling film-coated tablets are a long-term treatment – keep taking Brivaracetam Stirling film-coated tablets until your doctor tells you to stop.

If you take more Brivaracetam Stirling film-coated tablets than you should

If you have taken more Brivaracetam Stirling film-coated tablets than you should, talk to your doctor. You may feel dizzy and sleepy. You may also have any of the following symptoms: feeling sick, a feeling of ‘spinning’, problems of keeping your balance, anxiety, feeling very tired, irritability, being aggressive, not being able to sleep, depression, thoughts or attempts of harming or killing yourself.

If you forget to take Brivaracetam Stirling film-coated tablets

- If you miss a dose take it as soon as you remember.
- Then take your next dose at the time you would normally take it.
- Do not take a double dose to make up for a forgotten dose.
- If you are not sure what to do, ask your doctor or pharmacist.

If you stop taking Brivaracetam Stirling film-coated tablets

- Do not stop taking this medicine unless your doctor tells you to. This is because stopping treatment could increase the number of fits you have.
- If your doctor asks you to stop taking this medicine, they will lower your dose gradually. This helps to stop your fits coming back or getting worse.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4 POSSIBLE SIDE EFFECTS

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Very common: may affect more than 1 in 10 people

- feeling sleepy or dizzy

Common: may affect up to 1 in 10 people

- flu
- feeling very tired (fatigue)
- convulsion, a feeling of ‘spinning’ (vertigo)
- feeling and being sick, constipation
- depression, anxiety, not being able to sleep (insomnia), irritability
- infections of the nose and throat (such as the ‘common cold’), cough
- decreased appetite

Uncommon: may affect up to 1 in 100 people

- allergic reactions
- abnormal thinking and/or loss of touch with reality (psychotic disorder), being aggressive, nervous excitement (agitation)
- thoughts or attempts of harming or killing yourself: tell your doctor straight away
- a decrease in white blood cells (called ‘neutropenia’) – shown in blood tests

Not known: frequency cannot be estimated from the available data

- a widespread rash with blisters and peeling skin, particularly around the mouth, nose, eyes and genitals (Stevens-Johnson syndrome)

Additional side effects in children

Common: may affect up to 1 in 10 people

- restlessness and hyperactivity (psychomotor hyperactivity)

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via: Yellow Card Scheme
Website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

5 HOW TO STORE BRIVARACETAM STIRLING FILM-COATED TABLETS

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and blister after EXP. The expiry date refers to the last day of that month.

This medicinal product does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6 CONTENTS OF THE PACK AND OTHER INFORMATION

What Brivaracetam Stirling film-coated tablets contain

The active substance is brivaracetam.
Each film-coated tablet contains 10 mg, 25 mg, 50 mg, 75 mg, or 100 mg brivaracetam.

The other ingredients are:

Core

- Lactose monohydrate, lactose, croscarmellose sodium, magnesium stearate

Coating

- 10 mg film-coated tablets: poly(vinyl alcohol), titanium dioxide (E171), macrogol (3350), talc.
- 25 mg film-coated tablets: poly(vinyl alcohol), titanium dioxide (E171), macrogol (3350), talc, iron oxide black (E172), iron oxide yellow (E172).
- 50 mg film-coated tablets: poly(vinyl alcohol), titanium dioxide (E171), macrogol (3350), talc, iron oxide yellow (E172), iron oxide red (E172).
- 75 mg film-coated tablets: poly(vinyl alcohol), titanium dioxide (E171), macrogol (3350), talc, iron oxide black (E172), iron oxide yellow (E172), iron oxide red (E172).
- 100 mg film-coated tablets: poly(vinyl alcohol), titanium dioxide (E171), macrogol (3350), talc, iron oxide black (E172), iron oxide yellow (E172).

What Brivaracetam Stirling film-coated tablets look like and contents of the pack

- Brivaracetam Stirling 10 mg film-coated tablets are white to off white, round, film-coated tablets of approx. 5.5 mm diameter with ‘10’ debossed on one side and plain on the other side.
- Brivaracetam Stirling 25 mg are grey, oval, film-coated tablets (of approx. 8.9 mm × 5.0 mm) with ‘25’ debossed on one side and plain on the other side.
- Brivaracetam Stirling 50 mg film-coated tablets are yellow, oval, film-coated tablets (of approx. 11.7 mm × 6.6 mm) with score line on one side.
- Brivaracetam Stirling 75 mg film-coated tablets are purple, oval, film-coated tablets (of approx. 13 mm × 7.3 mm) with ‘75’ debossed on one side and plain on the other side.
- Brivaracetam Stirling 100 mg film-coated tablets are green-grey, oval, film-coated tablets (of approx. 14.5 mm × 8.1 mm) with score line on one side.

Brivaracetam Stirling film-coated tablets are packaged in blister packs supplied in cardboard boxes containing either 7, 14, 28, 56, 60, 7 × 1, 14 × 1, 28 × 1, 56 × 1 or 60 × 1 film-coated tablets or in multipacks containing either 168 (3 packs of 56) or 168 × 1 (3 packs of 56 × 1) film-coated tablets.

All packs are available in PVC/PCTFE – Aluminium blisters.

Not all pack sizes may be marketed.

Marketing Authorisation Holder

Stirling Anglian Pharmaceuticals Limited
Hillington Park Innovation Centre
1 Ainslie Road
Hillington Park
Glasgow G52 4RU

Manufacturer

PharmaPath S.A.
28is Oktovriou 1
Agia Varvara, 123 51
Greece

This leaflet was last revised in December 2025