

## SUMMARY OF PRODUCT CHARACTERISTICS

### 1 NAME OF THE MEDICINAL PRODUCT

Captopril 12.5mg Tablets and or Ecopace 12.5mg.

### 2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each Captopril Tablet 12.5mg contains Captopril BP 12.5mg

Excipient with known effect: one tablet contains 32.5 mg lactose.

For the full list of excipients, see section 6.1.

### 3 PHARMACEUTICAL FORM

Tablet.

Captopril 12.5mg tablets are white, round, flat, uncoated tablets with bevelled edge, scored on one side and marked “G12.5” on the other, 6mm in diameter.

Captopril 25mg tablets are white, round, flat, uncoated tablets with bevelled edge, scored on one side and marked “G25” on the other, 8mm in diameter.

Captopril 50mg tablets are white, round, flat, uncoated tablets with bevelled edge, scored on one side and marked “G50” on the other, 10mm in diameter.

The score line is only to facilitate breaking for ease of swallowing and not to divide into equal doses.

#### 4.1 Therapeutic indications

Hypertension: The management of mild to moderate hypertension. In severe hypertension it should be used where standard therapy is ineffective or inappropriate.

Congestive Heart Failure: Captopril is indicated for the treatment of congestive heart failure. The drug should be used together with diuretics, and when appropriate, digitalis and beta-blockers. In patients on doses of over 100 mg daily plus or minus a diuretic, in those with severe renal impairment or those with severe congestive heart failure use of captopril should be under specialist supervision.

Myocardial Infarction:

- *short-term (4 weeks) treatment*: Captopril is indicated in any clinically stable patient within the first 24 hours of an infarction.

- *long-term prevention of symptomatic heart failure:* Captopril is indicated in clinically stable patients with asymptomatic left ventricular dysfunction (ejection fraction < 40%) following myocardial infarction to improve survival, delay the onset of symptomatic heart failure, reduce hospitalisations for heart failure and reduce recurrent myocardial infarction and coronary revascularisation procedures.

Before starting therapy, cardiac function should be determined by radionuclide ventriculography or echocardiography.

Type I Diabetic Nephropathy:

Captopril is indicated in insulin dependent diabetics for the treatment of macroproteinuric diabetic nephropathy (microalbuminuria greater than 30 mg/day) (See Section 5.1). Captopril may prevent the progression of the renal disease and reduce associated clinical events e.g. dialysis, renal transplantation and death.

Captopril can be used alone or in combination with other antihypertensive agents (see sections 4.3, 4.4, 4.5 and 5.1).

## **4.2 Posology and method of administration**

Posology

Dose should be individualised according to patient's profile (see section 4.4) and blood pressure response. The recommended maximum daily dose is 150mg. Captopril may be taken before, during and after meals.

Adults:

Hypertension:

Treatment with captopril should be at the lowest effective dose which should be titrated according to the needs of the patient.

The recommended starting dose is 25-50mg daily in two divided doses. The dose may be increased incrementally, with intervals of at least 2 weeks, to 100-150mg/day in two divided doses as needed to reach target blood pressure. Captopril may be used alone or with other antihypertensive medication such as thiazide diuretics is added (see sections 4.3, 4.4, 4.5 and 5.1). A once-daily dosing regimen may be appropriate when concomitant antihypertensive medication such as thiazide diuretics is added.

In patients with a strongly active renin-angiotensin-aldosterone system (hypovolaemia, renovascular hypertension, cardiac decompensation) it is preferable to commence with a single dose of 6.25mg or 12.5mg. The inauguration of this treatment should preferably take place under close medical supervision. These doses will then be administered at a rate of two per day. The dosage can be gradually increased to 50mg per day in one or two doses and if necessary to 100mg per day in one or two doses.

Congestive Heart failure:

Captopril treatment for congestive heart failure must be started under close medical supervision. The usual starting dose is 6.25mg or 12.5mg BID or TID. Titration to the maintenance dose (75 – 150mg per day) should be based on

patient's response, clinical status and tolerability up to a maximum of 150mg per day in divided doses. The dose should be increased incrementally, with intervals of at least 14 days, to evaluate patient's response.

#### Myocardial infarction:

- *short-term treatment:* Captopril treatment should begin in hospital as soon as possible following the appearance of the signs and/or symptoms in patients with stable haemodynamics. A 6.25mg test dose should be administered, with a 12.5mg dose being administered 2 hours afterwards and a 25mg dose 12 hours later. From the following day, captopril should be administered in a 100mg/day dose, in two daily administrations, for 4 weeks, if warranted by the absence of adverse haemodynamic reactions. At the end of the 4 weeks of treatment, the patient's state should be reassessed before a decision is taken concerning treatment for the postmyocardial infarction stage.

- *chronic treatment:* if captopril treatment has not begun during the first 24 hours of the acute myocardial infarction stage, it is suggested that treatment be instigated between the 3rd and 16th day post-infarction once the necessary treatment conditions have been attained (stable haemodynamics and management of any residual ischaemia). Treatment should be started in hospital under strict surveillance (particularly of blood pressure) until the 75mg dose is reached. The initial dose must be low (see 4.4), particularly if the patient exhibits normal or low blood pressure at the initiation of therapy. Treatment should be initiated with a dose of 6.25mg followed by 12.5mg 3 times daily for 2 days and then 25mg 3 times daily if warranted by the absence of adverse haemodynamic reactions. The recommended dose for effective cardioprotection during long-term treatment is 75 to 150mg daily in two or three doses. In cases of symptomatic hypotension, as in heart failure, the dosage of diuretics and/or other concomitant vasodilators may be reduced in order to attain the steady state dose of captopril. Where necessary, the dose of captopril should be adjusted in accordance with the patient's clinical reactions. Captopril may be used in combination with other treatments for myocardial infarction such as thrombolytic agents, beta-blockers and acetylsalicylic acid.

#### Type I Diabetic nephropathy:

In patients with type I diabetic nephropathy, the recommended daily dose of captopril is 75-100mg in divided doses. Captopril may be used in combination with other antihypertensive agents, i.e. diuretics, beta blockers, centrally acting agents or vasodilators if the reduction in blood pressure is inadequate with captopril alone.

#### Elderly:

As with other antihypertensive agents, consideration should be given to initiating therapy with a lower starting dose (6.25mg BID) in elderly patients who may have reduced renal function and other organ dysfunctions. (see above 'renal impairment' and section 4.4).

The dose should be titrated against the blood pressure response and kept as low as possible to achieve adequate control.

#### Renal impairment:

Since captopril is excreted primarily via the kidneys, dosage should be reduced or the dosage interval should be increased in patients with impaired renal function. When concomitant diuretic therapy is required, a loop diuretic (e.g. furosemide), rather than a thiazide diuretic, is preferred in patients with severe renal impairment. In patients with impaired renal function, the following daily dose may be recommended to avoid accumulation of captopril.

Creatinine clearance (ml/min/1.73 m <sup>2</sup> )	Daily starting dose (mg)	Daily maximum dose (mg)
>40	25-50	150
21-40	25	100
10-20	12.5	75
<10	6.25	37.5

#### Paediatric population

The efficacy and safety of captopril have not been fully established. The use of captopril in children and adolescents should be initiated under close medical supervision. The initial dose of captopril is about 0.3mg/kg body weight. For patients requiring special precautions (children with renal dysfunction, premature infants, new-borns and infants, because their renal function is not the same with older children and adults) the starting dose should be only 0.15mg captopril/kg weight. Generally, captopril is administered to children 3 times a day, but dose and interval of dose should be adapted individually according to patient's response.

#### Method of administration:

For oral use.

### **4.3 Contraindications**

- Hypersensitivity to the active substance, any other ACE inhibitors or to any of the excipients listed in section 6.1.
- Second and third trimesters of pregnancy (see sections 4.4 and 4.6).
- History of angioedema associated with previous ACE inhibitor therapy.
- Hereditary / idiopathic angioneurotic oedema.
- Lactation (see 4.6).
- The concomitant use of captopril with aliskiren-containing products is contraindicated in patients with diabetes mellitus or renal impairment (GFR < 60 ml/min/1.73 m<sup>2</sup>) (see sections 4.5 and 5.1).
- Concomitant use with sacubitril/valsartan therapy. Captopril must not be initiated earlier than 36 hours after the last dose of sacubitril/valsartan (see also sections 4.4 and 4.5).

### **4.4 Special warnings and precautions for use**

*Angioedema:* angioedema of the extremities, face, lips, mucous membranes, tongue, glottis or larynx may occur in patients treated with ACE inhibitors including captopril. This may occur anytime during treatment. However, in rare cases, severe angioedema may develop after long-term treatment with an

ACE inhibitor. In such cases, captopril should be discontinued promptly and appropriate monitoring should be instituted to ensure complete resolution of symptoms prior to dismissing the patient. In those instances where swelling has been confined to the face and lips the condition generally resolved without treatment, although antihistamines have been useful in relieving symptoms. Angioedema involving the tongue, glottis or larynx may be fatal. Where there is involvement of the tongue, glottis or larynx, likely to cause airway obstruction, appropriate therapy, which may include subcutaneous epinephrine solution 1:1000 (0.3 ml to 0.5 ml) and/or measures to ensure a patent airway, should be administered promptly. Emergency therapy should be instituted. The patient should be hospitalised and observed for at least 12 to 24 hours and should not be discharged until complete resolution of symptoms has occurred.

Black patients receiving ACE inhibitors have been reported to have a higher incidence of angioedema compared to non-blacks.

Patients with a history of angioedema unrelated to ACE inhibitor therapy may be at increased risk of angioedema while receiving an ACE inhibitor (see 4.3).

Intestinal angioedema has also been reported rarely in patients treated with ACE inhibitors. These patients presented with abdominal pain (with or without nausea or vomiting); in some cases there was no prior facial angioedema and C-1 esterase levels were normal. The angioedema was diagnosed by procedures including abdominal CT scan, or ultrasound or at surgery and symptoms resolved after stopping the ACE inhibitor. Intestinal angioedema should be included in the differential diagnosis of patients on ACE inhibitors presenting with abdominal pain (see 4.8).

Concomitant use of ACE inhibitors with sacubitril/valsartan is contraindicated due to the increased risk of angioedema. Treatment with sacubitril/valsartan must not be initiated earlier than 36 hours after the last dose of captopril. Treatment with captopril must not be initiated earlier than 36 hours after the last dose of sacubitril/valsartan (see sections 4.3 and 4.5).

*Hypersensitivity/angioedema:* Concomitant use of ACE inhibitors with racecadotril, mTOR inhibitors (e.g. sirolimus, everolimus, temsirolimus and vildagliptin) : Patients taking concomitant mTOR inhibitors (e.g. sirolimus, everolimus, temsirolimus) therapy may be at increased risk for angioedema (e.g. swelling of the airways or tongue, with or without respiratory impairment) (see section 4.5).

Angioneurotic oedema of the extremities, face, lips, mucous membranes, tongue, glottis and/or larynx may occur in patients treated with ACE inhibitors particularly during the first week of treatment. However, in rare cases, severe angioedema may develop after months or years of long-term treatment with an ACE inhibitor. Treatment should be discontinued promptly. Angioedema involving the tongue, glottis or larynx may be fatal. Emergency therapy should be instituted.

Caution should be used when starting racecadotril, mTOR inhibitors (e.g. sirolimus, everolimus, temsirolimus) and vildagliptin in a patient already taking an ACE inhibitor. Serum potassium ACE inhibitors can cause hyperkalemia because they inhibit the release of aldosterone. The effect is usually not significant in patients with normal renal function. However, in patients with impaired renal function and/or in patients taking potassium supplements (including salt substitutes), potassium-sparing diuretics, trimethoprim or co-trimoxazole also known as trimethoprim/sulfamethoxazole and especially aldosterone antagonists or angiotensin receptor blockers, hyperkalemia can occur. Potassium sparing diuretics and angiotensin receptor blockers should be used with caution in patients receiving ACE inhibitors, and serum potassium and renal function should be monitored (see section 4.5).

*Aortic and mitral valve stenosis/Obstructive hypertrophic cardiomyopathy:* ACE inhibitors should be used with caution in patients with left ventricular valvular or outflow tract obstruction and avoided in cases of cardiogenic shock and haemodynamically significant obstruction.

*Renal impairment:*

The incidence of adverse reactions to captopril is principally associated with renal function since the drug is excreted primarily by the kidney. In cases of renal impairment (creatinine clearance < 40 ml/min), the initial dosage of captopril must be adjusted according to the patient's creatinine clearance (see 4.2), and then as a function of the patient's response to treatment. Routine monitoring of potassium and creatinine are part of normal medical practice for these patients.

The dose should not exceed that necessary for adequate control and should be reduced in patients with impaired renal function.

Evaluation of the patient should include assessment of renal function (monitoring of potassium and creatinine) prior to initiation of therapy and at appropriate intervals thereafter. Patients with renal impairment should not normally be treated with captopril.

*Hypotension:* rarely hypotension is observed in uncomplicated hypertensive patients. Symptomatic hypotension is more likely to occur in hypertensive patients who are volume and/or sodium depleted by vigorous diuretic therapy, dietary salt restriction, diarrhoea, vomiting or haemodialysis. Volume and/or sodium depletion should be corrected before the administration of an ACE inhibitor and a lower starting dose should be considered.

Patients with heart failure are at higher risk of hypotension and a lower starting dose is recommended when initiating therapy with an ACE inhibitor. The magnitude of the decrease is greatest early in the course of treatment; this effect stabilises within a week or two, and generally returns to pre-treatment levels, without a decrease in therapeutic efficacy, within two months. Caution should be used whenever the dose of captopril or diuretic is increased in patients with heart failure.

As with any antihypertensive agent, excessive blood pressure lowering in patients with ischaemic cardiovascular or cerebrovascular disease may increase the risk of myocardial infarction or stroke. If hypotension develops, the patient should be placed in a supine position. Volume repletion with intravenous normal saline may be required.

Infants, especially new-borns, may be more susceptible to the adverse haemodynamic effects of captopril. Excessive, prolonged and unpredictable decreases in blood pressure and associated complications, including oliguria and seizures have been reported.

*Renovascular hypertension:* there is an increased risk of hypotension and renal insufficiency when patients with bilateral renal artery stenosis or stenosis of the artery to a single functioning kidney are treated with ACE inhibitors. Loss of renal function may occur with only mild changes in serum creatinine. In these patients, therapy should be initiated under close medical supervision with low doses, careful titration and monitoring of renal function.

*Insulin Autoimmune Syndrome (IAS):*

Cases of Insulin Autoimmune Syndrome (IAS), including severe hypoglycaemic events have been reported during the treatment with captopril (see section 4.8). If IAS is suspected, captopril should be discontinued, and appropriate treatment should be initiated.

*Cough:* cough has been reported with the use of ACE inhibitors. Characteristically, the cough is non-productive, persistent and resolves after discontinuation of therapy.

*Hyperkalaemia:* elevations in serum potassium have been observed in some patients treated with ACE inhibitors, including captopril. Patients at risk for the development of hyperkalaemia include those with renal insufficiency, diabetes mellitus, hypoaldosteronism or those using concomitant potassium-sparing diuretics, potassium supplements or potassium-containing salt substitutes; or those patients taking other drugs associated with increases in serum potassium (e.g. heparin, co-trimoxazole also known as trimethoprim/sulfamethoxazole). If concomitant use of the above mentioned agents is deemed appropriate, regular monitoring of serum potassium is recommended (see section 4.5).

*Lithium:* the combination of lithium and captopril is not recommended due to the potentiation of lithium toxicity (see 4.5).

*Proteinuria:* proteinuria may occur particularly in patients with existing renal function impairment or on relatively high doses of ACE inhibitors.

Total urinary proteins greater than 1 g per day were seen in about 0.7% of patients receiving captopril. The majority of patients had evidence of prior renal disease or had received relatively high doses of captopril (in excess of

150 mg/day), or both. Nephrotic syndrome occurred in about one-fifth of proteinuric patients. In most cases, proteinuria subsided or cleared within six months whether or not captopril was continued. Parameters of renal function, such as BUN and creatinine, were seldom altered in the patients with proteinuria.

Patients with prior renal disease should have urinary protein estimations (dipstick on first morning urine) prior to treatment, and periodically thereafter.

Although membranous glomerulopathy was found in biopsies taken from some proteinuric patients, a causal relationship to captopril has not been established.

*Anaphylactoid reactions during desensitisation:* sustained life-threatening anaphylactoid reactions have been rarely reported for patients undergoing desensitising treatment with hymenoptera venom while receiving another ACE inhibitor. In the same patients, these reactions were avoided when the ACE inhibitor was temporarily withheld, but they reappeared upon inadvertent rechallenge. Therefore, caution should be used in patients treated with ACE inhibitors undergoing such desensitisation procedures.

*Anaphylactoid reactions during high-flux dialysis / lipoprotein apheresis membrane exposure:* anaphylactoid reactions have been reported in patients haemodialysed with high-flux dialysis membranes (e.g. AW 69) or undergoing low-density lipoprotein apheresis with dextran sulphate absorption in patients receiving ACE inhibitors. Therefore, this combination should be avoided. In these patients, consideration should be given to using a different type of dialysis, membrane or a different class of medication.

*Hepatic failure:* rarely, ACE inhibitors have been associated with a syndrome that starts with cholestatic jaundice and progresses to fulminant hepatic necrosis and (sometimes) death. The mechanism of this syndrome is not understood. Patients receiving ACE inhibitors who develop jaundice or marked elevations of hepatic enzymes should discontinue the ACE inhibitor and receive appropriate medical follow-up.

*Neutropenia/Agranulocytosis:* Neutropenia/agranulocytosis thrombocytopenia and anaemia have been reported in patients receiving ACE inhibitors including captopril. In patients with normal renal function and no other complicating factors, neutropenia occurs rarely. Captopril should be used with extreme caution in patients with collagen vascular disease, immunosuppressant therapy, treatment with allopurinol or procainamide, or a combination of these complicating factors, especially if there is pre-existing impaired renal function. Some of these patients developed serious infections which in a few instances did not respond to intensive antibiotic therapy.

If captopril is used in such patients, it is advised that white blood cell count and differential counts should be performed prior to therapy, every 2 weeks during the first three months of captopril therapy, and periodically thereafter.

During treatment, all patients should be instructed to report any sign of infection (e.g. sore throat, fever), when a differential white blood cell count should be performed. Captopril and other concomitant medication (see 4.5) should be withdrawn if neutropenia (neutrophils less than  $1000/\text{mm}^3$ ) is detected or suspected.

In most patients neutrophil counts rapidly returned to normal upon discontinuing captopril.

*Surgery / anaesthesia:* hypotension may occur in patients undergoing major surgery, or during anaesthesia with anaesthetic agents that are known to lower blood pressure. Captopril will block angiotensin II formation secondary to compensatory renin release. If hypotension occurs it may be corrected by volume expansion.

*Diabetic patients:* the glycaemia levels should be closely monitored in diabetic patients previously treated with oral antidiabetic drugs or insulin, namely during the first month of treatment with an ACE inhibitor.

*Renal function in patient with Heart Failure:*

Some patients may develop stable elevations of BUN and serum creatinine  $>20\%$  above normal or baseline upon long-term treatment with captopril. A few patients, generally those with severe pre-existing renal disease, required discontinuation of treatment due to progressively increasing creatinine.

*Risk of hypokalaemia:* the combination of an ACE inhibitor with a thiazide diuretic does not rule out the occurrence of hypokalaemia. Regular monitoring of kalaemia should be performed.

*Lactose:* Patients with rare hereditary problems of galactose intolerance, total lactase deficiency or glucose-galactose malabsorption should not take this medicine.

*Ethnic differences:* as with other angiotensin converting enzyme inhibitors, captopril is apparently less effective in lowering blood pressure in black people than in non-blacks, possibly because of a higher prevalence of low-renin states in the black hypertensive population.

*Pregnancy:* ACE inhibitors should not be initiated during pregnancy. Unless continued ACE inhibitor therapy is considered essential, patients planning pregnancy should be changed to alternative antihypertensive treatments which have an established safety profile for use in pregnancy. When pregnancy is diagnosed, treatment with ACE inhibitors should be stopped immediately, and, if appropriate, alternative therapy should be started (see sections 4.3 and 4.6).

*Dual blockade of the renin-angiotensin-aldosterone system (RAAS):* There is evidence that the concomitant use of ACE-inhibitors, angiotensin II receptor blockers or aliskiren increases the risk of hypotension, hyperkalaemia and decreased renal function (including acute renal failure). Dual blockade of RAAS through the combined use of ACE-inhibitors, angiotensin II receptor

blockers or aliskiren is therefore not recommended (see sections 4.5 and 5.1). If dual blockade therapy is considered absolutely necessary, this should only occur under specialist supervision and subject to frequent close monitoring of renal function, electrolytes and blood pressure. ACE-inhibitors and angiotensin II receptor blockers should not be used concomitantly in patients with diabetic nephropathy.

#### **4.5 Interaction with other medicinal products and other forms of interaction**

*Potassium sparing diuretics or potassium supplements:* ACE inhibitors attenuate diuretic induced potassium loss. Potassium sparing diuretics (e.g. spironolactone, triamterene or amiloride), potassium supplements, or potassium-containing salt substitutes may lead to significant increases in serum potassium. If concomitant use is indicated because of demonstrated hypokalaemia they should be used with caution and with frequent monitoring of serum potassium (see 4.4).

Although serum potassium usually remains within normal limits, hyperkalaemia may occur in some patients treated with captopril. Potassium sparing diuretics (e.g. spironolactone, triamterene, or amiloride), potassium supplements, or potassium containing salt substitutes may lead to significant increases in serum potassium. Care should also be taken when cilazapril is coadministered with other agents that increase serum potassium, such as trimethoprim and co-trimoxazole (trimethoprim/sulfamethoxazole) as trimethoprim is known to act as a potassium sparing diuretic like amiloride. Therefore, the combination of captopril with the above-mentioned drugs is not recommended. If concomitant use is indicated, they should be used with caution and with frequent monitoring of serum potassium.

*Diuretics (thiazide or loop diuretics):* prior treatment with high dose diuretics may result in volume depletion and a risk of hypotension when initiating therapy with captopril (see 4.4). The hypotensive effects can be reduced by discontinuation of the diuretic, by increasing volume or salt intake or by initiating therapy with a low dose of captopril. However, no clinically significant drug interactions have been found in specific studies with hydrochlorothiazide or furosemide.

##### *Ciclosporin:*

Hyperkalaemia may occur during concomitant use of ACE inhibitors with ciclosporin. Monitoring of serum potassium is recommended.

##### *Heparin:*

Hyperkalaemia may occur during concomitant use of ACE inhibitors with heparin. Monitoring of serum potassium is recommended.

*Other antihypertensive agents:* Captopril has been safely co-administered with other commonly used anti-hypertensive agents (e.g., beta-blockers and long-acting calcium channel blockers). Concomitant use of these agents may increase the hypotensive effects of captopril. Treatment with nitroglycerine and other nitrates, or other vasodilators (such as minoxidil), should be used with caution.

*Alpha blocking agents:* concomitant use of alpha blocking agents may increase the antihypertensive effects of captopril and increase the risk of orthostatic hypotension.

*Treatments of acute myocardial infarction:* captopril may be used concomitantly with acetylsalicylic acid (at cardiologic doses), thrombolytics, beta-blockers and/or nitrates in patients with myocardial infarction.

*Tricyclic antidepressants / Antipsychotics:* ACE inhibitors may enhance the hypotensive effects of certain tricyclic antidepressants and antipsychotics (see 4.4). Postural hypotension may occur.

*Non-steroidal anti-inflammatory medicinal products:* it has been described that non-steroidal anti-inflammatory medicinal products (NSAIDs) (such as Indomethacin, Ibuprofen) and ACE inhibitors exert an additive effect on the increase in serum potassium whereas renal function may decrease. These effects are, in principle, reversible. Rarely, acute renal failure may occur, particularly in patients with compromised renal function such as the elderly or dehydrated. Chronic administration of NSAIDs may reduce the antihypertensive effect of an ACE inhibitor.

*Allopurinol, procainamide, cytostatic or immunosuppressive agents:* concomitant administration with ACE inhibitors may lead to an increased risk for leucopenia especially when the latter are used at higher than currently recommended doses.

*Lithium:* reversible increases in serum lithium concentrations and toxicity have been reported during concomitant administration of lithium with ACE inhibitors. Concomitant use of thiazide diuretics may increase the risk of lithium toxicity and enhance the already increased risk of lithium toxicity with ACE inhibitors. Use of captopril with lithium is not recommended, but if the combination proves necessary, careful monitoring of serum lithium levels should be performed (see 4.4)

*Sympathomimetics:* may reduce the antihypertensive effects of ACE inhibitors; patients should be carefully monitored.

*Antidiabetics:* pharmacological studies have shown that ACE inhibitors, including captopril, can potentiate the blood glucose-reducing effects of insulin and oral antidiabetics such as sulphonylurea in diabetics. Should this very rare interaction occur, it may be necessary to reduce the dose of the antidiabetic during simultaneous treatment with ACE inhibitors.

*Medicines increasing the risk of angioedema*

Concomitant use of ACE inhibitors with sacubitril/valsartan is contraindicated as this increases the risk of angioedema (see section 4.3 and 4.4). Concomitant use of ACE inhibitors with racecadotril, *mTOR inhibitors* (e.g. *sirolimus, everolimus, temsirolimus and vildagliptin*): patients taking

concomitant mTOR inhibitors therapy may be at increased risk for angioedema (see section 4.4).

*Co-trimoxazole (trimethoprim/sulfamethoxazole):* patients taking concomitant co-trimoxazole (trimethoprim/sulfamethoxazole) may be at increased risk for hyperkalaemia (see section 4.4).

*Clinical chemistry:* Captopril may cause a false- positive urine test for acetone.

*Angiotensin II receptor blockers or aliskiren:* Clinical trial data has shown that dual blockade of the renin-angiotensin-aldosteronesystem (RAAS) through the combined use of ACE-inhibitors, angiotensin II receptor blockers or aliskiren is associated with a higher frequency of adverse events such as hypotension, hyperkalaemia and decreased renal function (including acute renal failure) compared to the use of a single RAAS-acting agent (see sections 4.3, 4.4 and 5.1).

*Probenecid:*

The renal clearance of captopril is reduced in the presence of probenecid.

*Clonidine:*

It has been suggested that the anti-hypertensive effect of captopril can be delayed when patients treated with clonidine are changed to captopril.

## 4.6 Fertility, pregnancy and lactation

### **Pregnancy:**

The use of ACE inhibitors is not recommended during the first trimester of pregnancy (see section 4.4). The use of ACE inhibitors is contraindicated during the second and third trimesters of pregnancy (see sections 4.3 and 4.4).
--

Epidemiological evidence regarding the risk of teratogenicity following exposure to ACE inhibitors during the first trimester of pregnancy has not been conclusive; however a small increase in risk cannot be excluded.

Unless continued ACE inhibitor therapy is considered essential, if a pregnancy is planned or confirmed, the switch to an alternative treatment, which have an established safety profile for use in pregnancy, When pregnancy is diagnosed, treatment with ACE inhibitors should be stopped immediately, and, if appropriate, alternative therapy should be started.

Exposure to ACE inhibitor therapy during the second and third trimesters is known to induce human foetotoxicity (decreased renal function, oligohydramnios, skull ossification retardation) and neonatal toxicity (renal failure, hypotension, hyperkalaemia). (See section 5.3). Should exposure to ACE inhibitors have occurred from the second trimester of pregnancy, ultrasound check of renal function and skull is recommended. Infants whose

mothers have taken ACE inhibitors should be closely observed for hypotension (see sections 4.3 and 4.4).

**Breast-feeding:**

Limited pharmacokinetic data demonstrate very low concentrations in breast milk (see section 5.2). Although these concentrations seem to be clinically irrelevant, the use of Captopril tablets in breastfeeding is not recommended for preterm infants and for the first few weeks after delivery, because of the hypothetical risk of cardiovascular and renal effects and because there is not enough clinical experience.

In the case of an older infant, the use of Captopril tablets in a breast-feeding mother may be considered if this treatment is necessary for the mother and the child is observed for any adverse effect.

**4.7 Effects on ability to drive and use machines**

As with other antihypertensives, the ability to drive and use machines may be reduced, namely at the start of the treatment, or when posology is modified, and also when used in combination with alcohol, but these effects depend on the individual's susceptibility.

**4.8 Undesirable effects**

Frequency is defined using the following convention: very common ( $\geq 1/10$ ), common ( $> 1/100$  to  $< 1/10$ ), uncommon ( $> 1/1,000$  to  $< 1/100$ ), rare ( $> 1/10,000$  to  $< 1/1,000$ ), very rare ( $< 1/10,000$ ), not known (cannot be estimated from the available data).

Undesirable effects reported for captopril and/or ACE inhibitor therapy include:

**Immune System Disorder:**

Frequency Not known: Insulin autoimmune syndrome

**Blood and lymphatic disorders:**

Very rare: neutropenia/agranulocytosis (see 4.4), pancytopenia particularly in patients with renal dysfunction (see 4.4), anaemia (including aplastic and haemolytic), thrombocytopenia, lymphadenopathy, eosinophilia, auto-immune diseases and/or positive ANA-titres.

**Metabolism and nutrition disorders:**

Uncommon: decreased appetite

Rare: anorexia

Very rare: hyperkalaemia, hyponatremia, hypoglycaemia (see 4.4)

**Psychiatric disorders:**

Common: insomnia

Very rare: confusion state, depression.

**Nervous system disorders:**

Common: dysgeusia, reversible and self-limiting taste impairment, dizziness

Uncommon: headache, paraesthesia  
Rare: somnolence  
Very rare: cerebrovascular incidents, including stroke cerebrovascular insufficiency and syncope.

**Eye disorders:**

Very rare: blurred vision

**Cardiac disorders:**

Uncommon: tachycardia, arrhythmia angina pectoris, palpitations.  
Very rare: cardiac arrest, cardiogenic shock

**Vascular disorders:**

Uncommon: hypotension (see 4.4), Raynaud syndrome, flush, pallor, orthostatic hypotension

**Respiratory, thoracic and mediastinal disorders:**

Common: dry, irritating (non-productive) cough (see 4.4) and dyspnoea  
Very rare: bronchospasm, rhinitis, allergic alveolitis / eosinophilic pneumonia

**Gastrointestinal disorders:**

Common: nausea, vomiting, epigastric discomfort, peptic ulcer, dyspepsia, loss of taste (usually reversible on stopping treatment), gastric irritations, abdominal pain, diarrhoea, constipation, dry mouth.  
Rare: weight loss and loss of appetite, stomatitis/aphthous ulcerations, intestinal angioedema, mouth ulcers  
Very rare: glossitis, , pancreatitis.

**Hepato-biliary disorders:**

Very rare: impaired hepatic function and cholestasis (including jaundice), hepatitis including necrosis, elevated liver enzymes and bilirubin,transaminase increased, blood alkaline phosphatase increased.

**Skin and subcutaneous tissue disorders:**

Common: pruritus with or without a rash, rash, and alopecia.  
Uncommon: angioedema (see 4.4)  
Very rare: urticaria, Stevens Johnson syndrome, erythema multiforme, photosensitivity, erythroderma, pemphigoid reactions and exfoliative dermatitis, psoriasis and psoriasis dermatitis

**Musculoskeletal, connective tissue and bone disorders:**

Very rare: myalgia, arthralgia.

**Renal and urinary disorders:**

Rare: renal impairment, renal failure, polyuria, oliguria, pollakiuria  
Very rare: nephrotic syndrome.

**Reproductive system and breast disorders:**

Very rare: impotence, erectile dysfunction, gynaecomastia.

**General disorders and administration site conditions:**

Uncommon: chest pain, fatigue, malaise, asthenia  
Very rare: fever

**Investigations:**

Very rare: proteinuria, eosinophilia, increase of serum potassium, decrease of serum sodium, elevation of BUN, serum creatinine and serum bilirubin, decreases in haemoglobin, haematocrit, leucocytes, thrombocytes, positive ANA-titre, elevated ESR.

#### **Reporting of suspected adverse reactions**

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via Yellow Card Scheme at:

[www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store.

### **4.9 Overdose**

Symptoms of overdosage are severe hypotension, shock, stupor, bradycardia, electrolyte disturbances and renal failure.

#### **Management:**

Measures to prevent absorption (e.g. gastric lavage, administration of adsorbents and sodium sulphate within 30 minutes after intake) and hasten elimination should be applied if ingestion is recent. If hypotension occurs, the patient should be placed in the shock position and salt and volume supplementations should be given rapidly. Treatment with angiotensin-II should be considered. Bradycardia or extensive vagal reactions should be treated by administering atropine. The use of a pacemaker may be considered.

Captopril may be removed from circulation by haemodialysis. Captopril is not adequately cleared by peritoneal dialysis.

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Agents acting on the renin-angiotensin system, ACE inhibitors, plain, ATC code: C09AA01.

Captopril is a highly specific, competitive inhibitor of angiotensin-I converting enzyme (ACE inhibitors).

The beneficial effects of ACE inhibitors appear to result primarily from the suppression of the plasma renin-angiotensin-aldosterone system. Renin is an endogenous enzyme synthesised by the kidneys and released into the circulation where it converts angiotensinogen to angiotensin-I a relatively inactive decapeptide. Angiotensin-I is then converted by angiotensin converting enzyme, a peptidyl dipeptidase, to angiotensin-II. Angiotensin-II is a potent vasoconstrictor responsible for arterial vasoconstriction and increased blood pressure, as well as for stimulation of the adrenal gland to secrete aldosterone. Inhibition of ACE results in decreased plasma angiotensin-II, which leads to decreased vasopressor activity and to reduced aldosterone secretion. Although the latter decrease is small, small increases in serum potassium concentrations may occur, along with sodium and fluid loss. The

cessation of the negative feedback of angiotensin-II on the renin secretion results in an increase of the plasma renin activity.

Another function of the converting enzyme is to degrade the potent vasodepressive kinin peptide bradykinin to inactive metabolites. Therefore, inhibition of ACE results in an increased activity of circulating and local kallikrein-kinin-system which contributes to peripheral vasodilation by activating the prostaglandin system; it is possible that this mechanism is involved in the hypotensive effect of ACE inhibitors and is responsible for certain adverse reactions.

Reductions of blood pressure are usually maximal 60 to 90 minutes after oral administration of an individual dose of captopril. The duration of effect is dose related. The reduction in blood pressure may be progressive, so to achieve maximal therapeutic effects, several weeks of therapy may be required. The blood pressure lowering effects of captopril and thiazide-type diuretics are additive.

In patients with hypertension, captopril causes a reduction in supine and erect blood pressure, without inducing any compensatory increase in heart rate, nor water and sodium retention.

In haemodynamic investigations, captopril caused a marked reduction in peripheral arterial resistance. In general there were no clinically relevant changes in renal plasma flow or glomerular filtration rate. In most patients, the antihypertensive effect began about 15 to 30 minutes after oral administration of captopril; the peak effect was achieved after 60 to 90 minutes. The maximum reduction in blood pressure of a defined captopril dose was generally visible after three to four weeks.

In the recommended daily dose, the antihypertensive effect persists even during long-term treatment. Temporary withdrawal of captopril does not cause any rapid, excessive increase in blood pressure (rebound). The treatment of hypertension with captopril leads also to a decrease in left ventricular hypertrophy.

Haemodynamic investigations in patients with heart failure, showed that captopril caused a reduction in peripheral systemic resistance and a rise in venous capacity. This resulted in a reduction in pre-load and after-load of the heart (reduction in ventricular filling pressure). In addition, rises in cardiac output, work index and exercise capacity have been observed during treatment with captopril. In a large, placebo-controlled study in patients with left ventricular dysfunction (LVEF  $\leq$ 40%) following myocardial infarction, it was shown that captopril (initiated between the 3rd to the 16th day after infarction) prolonged the survival time and reduced cardiovascular mortality. The latter was manifested as a delay in the development of symptomatic heart failure and a reduction in the necessity for hospitalisation due to heart failure compared to placebo. There was also a reduction in re-infarction and in cardiac revascularisation procedures and/or in the need for additional medication with diuretics and/or digitalis or an increase in their dosage compared to placebo.

A retrospective analysis showed that captopril reduced recurrent infarcts and cardiac revascularisation procedures (neither were target criteria of the study).

Another large, placebo-controlled study in patients with myocardial infarction showed that captopril (given within 24 hours of the event and for a duration of one month) significantly reduced overall mortality after 5 weeks compared to placebo. The favourable effect of captopril on total mortality was still detectable even after one year. No indication of a negative effect in relation to early mortality on the first day of treatment was found.

Captopril cardioprotection effects are observed regardless of the patient's age or gender, location of the infarction and concomitant treatments with proven efficacy during the postinfarction period (thrombolytic agents, beta-blockers and acetylsalicylic acid).

#### *Type I diabetic nephropathy*

In a placebo-controlled, multicentre double blind clinical trial in insulin-dependent (Type I) diabetes with proteinuria, with or without hypertension (simultaneous administration of other antihypertensives to control blood pressure was allowed), captopril significantly reduced (by 51%) the time to doubling of the baseline creatinine concentration compared to placebo; the incidence of terminal renal failure (dialysis, transplantation) or death was also significantly less common under captopril than under placebo (51%). In patients with diabetes and microalbuminuria, treatment with captopril reduced albumin excretion within two years.

The effects of treatment with captopril on the preservation of renal function are in addition to any benefit that may have been derived from the reduction in blood pressure.

Two large randomised, controlled trials (ONTARGET (ONgoing Telmisartan Alone and in combination with Ramipril Global Endpoint Trial) and VA NEPHRON-D (The Veterans Affairs Nephropathy in Diabetes)) have examined the use of the combination of an ACE-inhibitor with an angiotensin II receptor blocker.

ONTARGET was a study conducted in patients with a history of cardiovascular or cerebrovascular disease, or type 2 diabetes mellitus accompanied by evidence of endorgan damage. VA NEPHRON-D was a study in patients with type 2 diabetes mellitus and diabetic nephropathy.

These studies have shown no significant beneficial effect on renal and/or cardiovascular outcomes and mortality, while an increased risk of hyperkalaemia, acute kidney injury and/or hypotension as compared to monotherapy was observed. Given their similar pharmacodynamic properties, these results are also relevant for other ACE-inhibitors and angiotensin II receptor blockers. ACE-inhibitors and angiotensin II receptor blockers should therefore not be used concomitantly in patients with diabetic nephropathy. ALTITUDE (Aliskiren Trial in Type 2 Diabetes Using Cardiovascular and Renal Disease Endpoints) was a study designed to test the benefit of adding

aliskiren to a standard therapy of an ACE-inhibitor or an angiotensin II receptor blocker in patients with type 2 diabetes mellitus and chronic kidney disease, cardiovascular disease, or both. The study was terminated early because of an increased risk of adverse outcomes. Cardiovascular death and stroke were both numerically more frequent in the aliskiren group than in the placebo group and adverse events and serious adverse events of interest (hyperkalaemia, hypotension and renal dysfunction) were more frequently reported in the aliskiren group than in the placebo group.

## **5.2 Pharmacokinetic properties**

### **Biotransformation**

Captopril is an orally active agent that does not require biotransformation for activity.

### **Absorption**

The average minimal absorption is approximately 75%. Peak plasma concentrations are reached within 60-90 minutes. The presence of food in the gastrointestinal tract reduces absorption by about 30-40%. Approximately 25-30% of the circulating drug is bound to plasma proteins.

### **Elimination**

The apparent elimination half-life of unchanged captopril in blood is about 2 hours. Greater than 95% of the absorbed dose is eliminated in the urine within 24 hours; 40-50% is unchanged drug and the remainder are inactive disulphide metabolites (captopril disulphide and captopril cysteine disulphide). Impaired renal function could result in drug accumulation. Therefore, in patients with impaired renal function the dose should be reduced and/or dosage interval prolonged (see 4.2).

Studies in animals indicate that captopril does not cross the blood-brain barrier to any significant extent.

### **Lactation:**

In the report of twelve women taking oral captopril 100 mg 3 times daily, the average peak milk level was 4.7 µg/L and occurred 3.8 hours after the dose. Based on these data, the maximum daily dosage that a nursing infant would receive is less than 0.002% of the maternal daily dosage.

## **5.3 Preclinical safety data**

Animal studies performed during organogenesis with captopril have not shown any teratogenic effect but captopril has produced foetal toxicity in several species, including foetal mortality during late pregnancy, growth retardation and postnatal mortality in the rat. Preclinical data reveal no other specific hazard for humans based on conventional studies of safety pharmacology, repeated dose toxicology, genotoxicity and carcinogenicity.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Lactose monohydrate

Maize starch

Cellulose, microcrystalline

Stearic acid

### **6.2 Incompatibilities**

Not applicable.

### **6.3 Shelf life**

3 years.

### **6.4 Special precautions for storage**

Store in the original package in order to protect from light. Do not store above 25 °C.

### **6.5 Nature and contents of container**

Captopril 12.5mg, 25mg and 50mg tablets are available in PVC/PVDC/Alu blisters of 14 tablets, in packs of 56 tablets.

### **6.6 Special precautions for disposal**

No special precautions.

## **7 MARKETING AUTHORISATION HOLDER**

Crescent Pharma Limited  
Key House, Sarum Hill,  
Basingstoke, RG21 8SR  
United Kingdom

**8     MARKETING AUTHORISATION NUMBER(S)**

PL 20416/0307

**9     DATE OF FIRST AUTHORISATION/RENEWAL OF THE  
AUTHORISATION**

22/11/1996

**10    DATE OF REVISION OF THE TEXT**

28/04/2026