

## **SUMMARY OF PRODUCT CHARACTERISTICS**

### **1 NAME OF THE MEDICINAL PRODUCT**

Cuenca 80 mg film-coated tablets

### **2 QUALITATIVE AND QUANTITATIVE COMPOSITION**

One film-coated tablet contains 80 mg of valsartan.

Cuenca 80 mg film-coated tablets:

Excipients with known effect:

sorbitol..... 9.25 mg

lactose monohydrate.. 1.1 mg

sodium..... 0.32 mg (0.01 mmol)

### **3 PHARMACEUTICAL FORM**

Film-coated tablet.

Cuenca 80 mg film-coated tablets: cylindrical, coated, scored on one side, pink film-coated tablets.

The tablet can be divided into equal doses.

### **4 CLINICAL PARTICULARS**

#### **4.1 Therapeutic indications**

##### Hypertension

Treatment of essential hypertension in adults, and hypertension in children and adolescents 6 to 18 years of age.

##### Recent myocardial infarction

Treatment of clinically stable adult patients with symptomatic heart failure or asymptomatic left ventricular systolic dysfunction after a recent (12 hours-10 days) myocardial infarction (see sections 4.4 and 5.1).

##### Heart failure

Treatment of symptomatic heart failure in adult patients when Angiotensin Converting Enzyme (ACE) inhibitors cannot be used, or as add-on therapy to ACE inhibitors when beta blockers cannot be used (see sections 4.4 and 5.1).

## 4.2 Posology and method of administration

### Posology

#### Hypertension

The recommended starting dose of Cuenca is 80 mg once daily. The antihypertensive effect is substantially present within 2 weeks, and maximal effects are attained within 4 weeks. In some patients whose blood pressure is not adequately controlled, the dose can be increased to 160 mg and to a maximum of 320 mg.

Cuenca may also be administered with other antihypertensive agents. The addition of a diuretic such as hydrochlorothiazide will decrease blood pressure even further in these patients.

#### Recent myocardial infarction

In clinically stable patients, therapy may be initiated as early as 12 hours after a myocardial infarction. After an initial dose of 20 mg twice daily, valsartan should be titrated to 40 mg, 80 mg, and 160 mg twice daily over the next few weeks. The starting dose is provided by the 40 mg divisible tablet.

The target maximum dose is 160 mg twice daily. In general, it is recommended that patients achieve a dose level of 80 mg twice daily by two weeks after treatment initiation and that the target maximum dose, 160 mg twice daily, be achieved by three months, based on the patient's tolerability. If symptomatic hypotension or renal dysfunction occur, consideration should be given to a dose reduction.

Valsartan may be used in patients treated with other post-myocardial infarction therapies, e.g. thrombolytics, acetylsalicylic acid, beta blockers, statins, and diuretics. The combination with ACE inhibitors is not recommended (see sections 4.4 and 5.1).

Evaluation of post-myocardial infarction patients should always include assessment of renal function.

#### Heart failure

The recommended starting dose of Cuenca is 40 mg twice daily. Uptitration to 80 mg and 160 mg twice daily should be done at intervals of at least two weeks to the highest dose, as tolerated by the patient.

Consideration should be given to reducing the dose of concomitant diuretics. The maximum daily dose administered in clinical trials is 320 mg in divided doses.

Valsartan may be administered with other heart failure therapies. However, the triple combination of an ACE inhibitor, a beta blocker and valsartan is not recommended (see sections 4.4 and 5.1).

Evaluation of patients with heart failure should always include assessment of renal function.

### Additional information on special populations

#### Elderly

No dose adjustment is required in elderly patients.

#### Renal impairment

No dose adjustment is required for adult patients with a creatinine clearance >10 ml/min (see sections 4.4 and 5.2). Concomitant use of Cuenca with aliskiren is contraindicated in patients with renal impairment (GFR < 60 mL/min/1.73 m<sup>2</sup>) (see section 4.3).

### Diabetes Mellitus

Concomitant use of Cuenca with aliskiren is contraindicated in patients with diabetes mellitus (see section 4.3).

### Hepatic impairment

Cuenca is contraindicated in patients with severe hepatic impairment, biliary cirrhosis and in patients with cholestasis (see sections 4.3, 4.4 and 5.2). In patients with mild to moderate hepatic impairment without cholestasis, the dose of valsartan should not exceed 80 mg.

### Paediatric population

#### Paediatric hypertension

##### *Children and adolescents 6 to 18 years of age*

The initial dose is 40 mg once daily for children weighing below 35 kg and 80 mg once daily for those weighing 35 kg or more. The dose should be adjusted based on blood pressure response. For maximum doses studied in clinical trials please refer to the table below.

Doses higher than those listed have not been studied and are therefore not recommended.

Weight	Maximum dose studied in clinical trials
≥18 kg to <35 kg	80 mg
≥35 kg to <80 kg	160 mg
≥80 kg to ≤ 160 kg	320 mg

##### *Children less than 6 years of age*

Available data are described in sections 4.8, 5.1 and 5.2. However safety and efficacy of Cuenca in children aged 1 to 6 years have not been established.

#### Use in paediatric patients aged 6 to 18 years with renal impairment

Use in paediatric patients with a creatinine clearance <30 ml/min and paediatric patients undergoing dialysis has not been studied, therefore valsartan is not recommended in these patients. No dose adjustment is required for paediatric patients with a creatinine clearance >30 ml/min. Renal function and serum potassium should be closely monitored (see sections 4.4 and 5.2).

#### Use in paediatric patients aged 6 to 18 years with hepatic impairment

As in adults, Cuenca is contraindicated in paediatric patients with severe hepatic impairment, biliary cirrhosis and in patients with cholestasis (see sections 4.3, 4.4 and 5.2). There is limited clinical experience with Cuenca in paediatric patients with mild to moderate hepatic impairment. The dose of valsartan should not exceed 80 mg in these patients.

#### Paediatric heart failure and recent myocardial infarction

Cuenca is not recommended for the treatment of heart failure or recent myocardial infarction in children and adolescents below the age of 18 years due to the lack of data on safety and efficacy.

#### Method of administration

Cuenca may be taken independently of a meal and should be administered with water.

### 4.3 Contraindications

- Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.
- Severe hepatic impairment, biliary cirrhosis and cholestasis.
- Second and third trimester of pregnancy (see sections 4.4 and 4.6).
- Concomitant use of angiotensin receptor antagonists (ARBs) - including Cuenca - or of angiotensin-converting-enzyme inhibitors (ACEIs) with aliskiren in patients with diabetes mellitus or renal impairment (GFR < 60 mL/min/1.73m<sup>2</sup>) (see sections 4.4 and 4.5).

### 4.4 Special warnings and precautions for use

#### Hyperkalaemia

Concomitant use with potassium supplements, potassium-sparing diuretics, salt substitutes containing potassium, or other agents that may increase potassium levels (heparin, etc.) is not recommended.

Monitoring of potassium should be undertaken as appropriate.

#### Impaired renal function

There is currently no experience on the safe use in patients with a creatinine clearance <10 ml/min and patients undergoing dialysis, therefore valsartan should be used with caution in these patients. No dose adjustment is required for adult patients with creatinine clearance >10 ml/min (see sections 4.2 and 5.2).

#### Hepatic impairment

In patients with mild to moderate hepatic impairment without cholestasis, Cuenca should be used with caution (see sections 4.2 and 5.2).

#### Sodium- and/or volume-depleted patients

In severely sodium-depleted and/or volume-depleted patients, such as those receiving high doses of diuretics, symptomatic hypotension may occur in rare cases after initiation of therapy with Cuenca. Sodium and/or volume depletion should be corrected before starting treatment with Cuenca, for example by reducing the diuretic dose.

#### Renal artery stenosis

In patients with bilateral renal artery stenosis or stenosis to a solitary kidney, the safe use of Cuenca has not been established.

Short-term administration of Cuenca to twelve patients with renovascular hypertension secondary to unilateral renal artery stenosis did not induce any significant changes in renal haemodynamics, serum creatinine, or blood urea nitrogen (BUN). However, other agents that affect the renin-angiotensin system may increase blood urea and serum creatinine in patients with unilateral renal artery stenosis, therefore monitoring of renal function is recommended when patients are treated with valsartan.

#### Kidney transplantation

There is currently no experience on the safe use of Cuenca in patients who have recently undergone kidney transplantation.

#### Primary hyperaldosteronism

Patients with primary hyperaldosteronism should not be treated with Cuenca as their renin-angiotensin system is not activated.

### Aortic and mitral valve stenosis, obstructive hypertrophic cardiomyopathy

As with all other vasodilators, special caution is indicated in patients suffering from aortic or mitral stenosis, or hypertrophic obstructive cardiomyopathy (HOCM).

### Pregnancy

Angiotensin II Receptor Antagonists (AIIRAs) should not be initiated during pregnancy. Unless continued AIIRAs therapy is considered essential, patients planning pregnancy should be changed to alternative anti-hypertensive treatments which have an established safety profile for use in pregnancy. When pregnancy is diagnosed, treatment with AIIRAs should be stopped immediately, and, if appropriate, alternative therapy should be started (see sections 4.3 and 4.6).

### Recent myocardial infarction

The combination of captopril and valsartan has shown no additional clinical benefit, instead the risk for adverse events increased compared to treatment with the respective therapies (see sections 4.2 and 5.1). Therefore, the combination of valsartan with an ACE inhibitor is not recommended.

Caution should be observed when initiating therapy in post-myocardial infarction patients. Evaluation of post-myocardial infarction patients should always include assessment of renal function (see section 4.2). Use of Cuenca in post-myocardial infarction patients commonly results in some reduction in blood pressure, but discontinuation of therapy because of continuing symptomatic hypotension is not usually necessary provided dosing instructions are followed (see section 4.2).

### Heart Failure

In patients with heart failure, the triple combination of an ACE inhibitor, a beta blocker and Cuenca has not shown any clinical benefit (see section 5.1). This combination apparently increases the risk for adverse events and is therefore not recommended.

Caution should be observed when initiating therapy in patients with heart failure. Evaluation of patients with heart failure should always include assessment of renal function (see section 4.2).

Use of Cuenca in patients with heart failure commonly results in some reduction in blood pressure, but discontinuation of therapy because of continuing symptomatic hypotension is not usually necessary provided dosing instructions are followed (see section 4.2).

In patients whose renal function may depend on the activity of the renin-angiotensin system (e.g. patients with severe congestive heart failure), treatment with angiotensin converting enzyme inhibitors has been associated with oliguria and/or progressive azotaemia and in rare cases with acute renal failure and/or death. As valsartan is an angiotensin II antagonist, it cannot be excluded that the use of Cuenca may be associated with impairment of the renal function.

### History of angioedema

Angioedema, including swelling of the larynx and glottis, causing airway obstruction and/or swelling of the face, lips, pharynx, and/or tongue has been reported in patients treated with valsartan; some of these patients previously experienced angioedema with other drugs including ACE inhibitors. <invented name> should be immediately discontinued in patients who develop angioedema, and <invented name> should not be re-administered (see section 4.8).

### Dual Blockade of the Renin-Angiotensin-Aldosterone System (RAAS)

Hypotension, syncope, stroke, hyperkalaemia, and changes in renal function (including acute renal failure) have been reported in susceptible individuals, especially if combining medicinal products that affect this system. Dual blockade of the renin-angiotensin-aldosterone system by combining aliskiren with an angiotensin converting enzyme inhibitor (ACEI) or an angiotensin II receptor blocker (ARB) is therefore not recommended.

The use of aliskiren in combination with Cuenca is contraindicated in patients with diabetes mellitus or renal impairment (GFR < 60 ml/min/1.73 m<sup>2</sup>) (see section 4.3).

### Paediatric population

#### Impaired renal function

Use in paediatric patients with a creatinine clearance <30 ml/min and paediatric patients undergoing dialysis has not been studied, therefore valsartan is not recommended in these patients. No dose adjustment is required for paediatric patients with a creatinine clearance >30 ml/min (see sections 4.2 and 5.2). Renal function and serum potassium should be closely monitored during treatment with valsartan. This applies particularly when valsartan is given in the presence of other conditions (fever, dehydration) likely to impair renal function.

The concomitant use of ARBs - including Cuenca - or of ACEIs with aliskiren is contraindicated in patients with renal impairment (GFR < 60 mL/min/1.73m<sup>2</sup>) (see sections 4.3 and 4.5).

#### Impaired hepatic function

As in adults, Cuenca is contraindicated in paediatric patients with severe hepatic impairment, biliary cirrhosis and in patients with cholestasis (see sections 4.3 and 5.2). There is limited clinical experience with Cuenca in paediatric patients with mild to moderate hepatic impairment. The dose of valsartan should not exceed 80 mg in these patients.

#### Warning about excipients:

This medicine contains sorbitol. Patients with hereditary problems of fructose intolerance should not take this medicine.

This medicine contains lactose. Patients with hereditary problems of galactose intolerance, the Lapp lactase deficiency (insufficiency observed in certain populations of Lapland) or glucose-galactose malabsorption should not take this medicine.

This medicine contains less than 23 mg of sodium per dose; that is, essentially “sodium free”.

## **4.5 Interaction with other medicinal products and other forms of interaction**

### Dual blockade of the Renin-Angiotensin- System (RAS) with ARBs, ACEIs, or aliskiren

Caution is required while co-administering ARBs, including Cuenca, with other agents blocking the RAAS such as ACEIs or aliskiren (see section 4.4).

Concomitant use of angiotensin receptor antagonists (ARBs) - including Cuenca - or of angiotensin-converting-enzyme inhibitors (ACEIs) with aliskiren in patients with diabetes mellitus or renal impairment (GFR < 60 ml/min/1.73m<sup>2</sup>) is contraindicated (see section 4.3).

### Concomitant use not recommended

#### *Lithium*

Reversible increases in serum lithium concentrations and toxicity have been reported during concurrent use of ACE inhibitors. Due to the lack of experience with concomitant

use of valsartan and lithium, this combination is not recommended. If the combination proves necessary, careful monitoring of serum lithium levels is recommended.

*Potassium-sparing diuretics, potassium supplements, salt substitutes containing potassium and other substances that may increase potassium levels*

If a medicinal product that affects potassium levels is considered necessary in combination with valsartan, monitoring of potassium plasma levels is advised.

#### Caution required with concomitant use

*Non-steroidal anti-inflammatory medicines (NSAIDs), including selective COX-2 inhibitors, acetylsalicylic acid >3 g/day, and non-selective NSAIDs*

When angiotensin II antagonists are administered simultaneously with NSAIDs, attenuation of the antihypertensive effect may occur. Furthermore, concomitant use of angiotensin II antagonists and NSAIDs may lead to an increased risk of worsening of renal function and an increase in serum potassium. Therefore, monitoring of renal function at the beginning of the treatment is recommended, as well as adequate hydration of the patient.

#### Transporters

In vitro data indicates that valsartan is a substrate of the hepatic uptake transporter OATP1B1/OATP1B3 and the hepatic efflux transporter MRP2. The clinical relevance of this finding is unknown. Co-administration of inhibitors of the uptake transporter (eg. rifampin, ciclosporin) or efflux transporter (eg. ritonavir) may increase the systemic exposure to valsartan. Exercise appropriate care when initiating or ending concomitant treatment with such drugs.

#### *Others*

In drug interaction studies with valsartan, no interactions of clinical significance have been found with valsartan or any of the following substances: cimetidine, warfarin, furosemide, digoxin, atenolol, indometacin, hydrochlorothiazide, amlodipine, glibenclamide.

#### Paediatric population

In hypertension in children and adolescents, where underlying renal abnormalities are common, caution is recommended with the concomitant use of valsartan and other substances that inhibit the renin angiotensin aldosterone system which may increase serum potassium. Renal function and serum potassium should be closely monitored.

## **4.6 Fertility, pregnancy and lactation**

### Pregnancy

The use of Angiotensin II Receptor Antagonists (AIIRAs) is not recommended during the first trimester of pregnancy (see section 4.4). The use of AIIRAs is contra-indicated during the second and third trimester of pregnancy (see sections 4.3 and 4.4).

Epidemiological evidence regarding the risk of teratogenicity following exposure to ACE inhibitors during the first trimester of pregnancy has not been conclusive; however, a small increase in risk cannot be excluded. Whilst there is no controlled epidemiological data on the risk with AIIRAs, similar risks may exist for this class of drugs. Unless continued AIIRA therapy is considered essential, patients planning pregnancy should be changed to alternative anti-hypertensive treatments which have an established safety profile for use in pregnancy. When pregnancy is diagnosed, treatment with AIIRAs should be stopped immediately, and, if appropriate, alternative therapy should be started. AIIRAs

therapy exposure during the second and third trimesters is known to induce human fetotoxicity (decreased renal function, oligohydramnios, skull ossification retardation) and neonatal toxicity (renal failure, hypotension, hyperkalemia); see also section 5.3 "Preclinical safety data".

Should exposure to AIIRAs have occurred from the second trimester of pregnancy, ultrasound check of renal function and skull is recommended.

Infants whose mothers have taken AIIRAs should be closely observed for hypotension (see also sections 4.3 and 4.4).

Breastfeeding Because no information is available regarding the use of valsartan during breastfeeding, Cuenca is not recommended and alternative treatments with better established safety profiles during breast-feeding are preferable, especially while nursing a newborn or preterm infant.

#### Fertility

Valsartan had no adverse effects on the reproductive performance of male or female rats at oral doses up to 200 mg/kg/day. This dose is 6 times the maximum recommended human dose on a  $\text{mg}/\text{m}^2$  basis (calculations assume an oral dose of 320 mg/day and a 60-kg patient).

### **4.7 Effects on ability to drive and use machines**

No studies on the effects on the ability to drive have been performed. When driving vehicles or operating machines it should be taken into account that occasionally dizziness or weariness may occur.

### **4.8 Undesirable effects**

In controlled clinical studies in adult patients with hypertension, the overall incidence of adverse reactions (ADRs) was comparable with placebo and is consistent with the pharmacology of valsartan. The incidence of ADRs did not appear to be related to dose or treatment duration and also showed no association with gender, age or race.

The ADRs reported from clinical studies, post-marketing experience and laboratory findings are listed below according to system organ class.

Adverse reactions are ranked by frequency, the most frequent first, using the following convention:

Very common ( $\geq 1/10$ ),

Common ( $\geq 1/100$  to  $< 1/10$ )

Uncommon ( $\geq 1/1,000$  to  $< 1/100$ )

Rare ( $\geq 1/10,000$  to  $< 1/1,000$ )

Very rare ( $< 1/10,000$ )

Not known (cannot be estimated from the available data)

Within each frequency grouping, adverse reactions are ranked in order of decreasing seriousness.

For all the ADRs reported from post-marketing experience and laboratory findings, it is not possible to apply any ADR frequency and therefore they are mentioned with a "not known" frequency.

- Hypertension

<b>Blood and lymphatic system disorders</b>	
Not known	Decrease in haemoglobin, Decrease in haematocrit, Neutropenia, Thrombocytopenia
<b>Immune system disorders</b>	
Not known	Hypersensitivity including serum sickness
<b>Metabolism and nutrition disorders</b>	
Not known	Increase of serum potassium, Hyponatraemia
<b>Ear and labyrinth system disorders</b>	
Uncommon	Vertigo
<b>Vascular disorders</b>	
Not known	Vasculitis
<b>Respiratory, thoracic and mediastinal disorders</b>	
Uncommon	Cough
<b>Gastrointestinal disorders</b>	
Uncommon	Abdominal pain
<b>Hepato-biliary disorders</b>	
Not known	Elevation of liver function values including increase of serum bilirubin
<b>Skin and subcutaneous tissue disorders</b>	
Not known	Angioedema, Rash, Pruritus
<b>Musculoskeletal and connective tissue disorders</b>	
Not known	Myalgia
<b>Renal and urinary disorders</b>	
Not known	Renal failure and impairment, Elevation of serum creatinine
<b>General disorders and administration site conditions</b>	
Uncommon	Fatigue

### Paediatric population

#### Hypertension

The antihypertensive effect of valsartan has been evaluated in two randomised, double-blind clinical studies in 561 paediatric patients from 6 to 18 years of age. With the exception of isolated gastrointestinal disorders (like abdominal pain, nausea, vomiting) and dizziness, no relevant differences in terms of type, frequency and severity of adverse reactions were identified between the safety profile for paediatric patients aged 6 to 18 years and that previously reported for adult patients.

Neurocognitive and developmental assessment of paediatric patients aged 6 to 16 years of age revealed no overall clinically relevant adverse impact after treatment with Cuenca for up to one year.

In a double-blind randomized study in 90 children aged 1 to 6 years, which was followed by a one-year open-label extension, two deaths and isolated cases of marked liver transaminases elevations were observed. These cases occurred in a population who had significant comorbidities. A causal relationship to Cuenca has not been established. In a second study in which 75 children aged 1 to 6 years were randomised, no significant liver transaminase elevations or death occurred with valsartan treatment.

Hyperkalaemia was more frequently observed in children and adolescents aged 6 to 18

years with underlying chronic kidney disease. The safety profile seen in controlled-clinical studies in adult patients with post-myocardial infarction and/or heart failure varies from the overall safety profile seen in hypertensive patients. This may relate to the patients underlying disease. ADRs that occurred in adult patients with post-myocardial infarction and/or heart failure patients are listed below.

- Post-myocardial infarction and/or heart failure (studied in adult patients only)

<b>Blood and lymphatic system disorders</b>	
Not known	Thrombocytopenia
<b>Immune system disorders</b>	
Not known	Hypersensitivity including serum sickness
<b>Metabolism and nutrition disorders</b>	
Uncommon	Hyperkalaemia
Not known	Increase of serum potassium, Hyponatraemia
<b>Nervous system disorders</b>	
Common	Dizziness, Postural dizziness
Uncommon	Syncope, Headache
<b>Ear and labyrinth system disorders</b>	
Uncommon	Vertigo
<b>Cardiac disorders</b>	
Uncommon	Cardiac failure
<b>Vascular disorders</b>	
Common	Hypotension, Orthostatic hypotension
Not known	Vasculitis
<b>Respiratory, thoracic and mediastinal disorders</b>	
Uncommon	Cough
<b>Gastrointestinal disorders</b>	
Uncommon	Nausea, Diarrhoea
<b>Hepato-biliary disorders</b>	
Not known	Elevation of liver function values
<b>Skin and subcutaneous tissue disorders</b>	
Uncommon	Angioedema
Not known	Rash, Pruritis
<b>Musculoskeletal and connective tissue disorders</b>	
Not known	Myalgia
<b>Renal and urinary disorders</b>	
Common	Renal failure and impairment
Uncommon	Acute renal failure, Elevation of serum creatinine
Not known	Increase in Blood Urea Nitrogen
<b>General disorders and administration site conditions</b>	
Uncommon	Asthenia, Fatigue

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in Appendix V\*.

## 4.9 Overdose

Symptoms

Overdose with Cuenca may result in marked hypotension, which could lead to depressed level of consciousness, circulatory collapse and/or shock.

#### Therapy

The therapeutic measures depend on the time of ingestion and the type and severity of the symptoms; stabilisation of the circulatory condition is of prime importance. If hypotension occurs, the patient should be placed in a supine position and blood volume correction should be undertaken.

Valsartan is unlikely to be removed by haemodialysis.

## **5 PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Angiotensin II Antagonists, plain, ATC code: C09CA03

Valsartan is an orally active, potent, and specific angiotensin II (Ang II) receptor antagonist. It acts selectively on the AT<sub>1</sub> receptor subtype, which is responsible for the known actions of angiotensin II. The increased plasma levels of Ang II following AT<sub>1</sub> receptor blockade with valsartan may stimulate the unblocked AT<sub>2</sub> receptor, which appears to counterbalance the effect of the AT<sub>1</sub> receptor. Valsartan does not exhibit any partial agonist activity at the AT<sub>1</sub> receptor and has much (about 20,000 fold) greater affinity for the AT<sub>1</sub> receptor than for the AT<sub>2</sub> receptor. Valsartan is not known to bind to or block other hormone receptors or ion channels known to be important in cardiovascular regulation.

Valsartan does not inhibit ACE (also known as kininase II) which converts Ang I to Ang II and degrades bradykinin. Since there is no effect on ACE and no potentiation of bradykinin or substance P, angiotensin II antagonists are unlikely to be associated with coughing. In clinical trials where valsartan was compared with an ACE inhibitor, the incidence of dry cough was significantly ( $p < 0.05$ ) less in patients treated with valsartan than in those treated with an ACE inhibitor (2.6% versus 7.9% respectively). In a clinical trial of patients with a history of dry cough during ACE inhibitor therapy, 19.5% of trial subjects receiving valsartan and 19.0% of those receiving a thiazide diuretic experienced cough compared to 68.5% of those treated with an ACE inhibitor ( $p < 0.05$ ).

#### Hypertension

Administration of Cuenca to patients with hypertension results in reduction of blood pressure without affecting pulse rate.

In most patients, after administration of a single oral dose, onset of antihypertensive activity occurs within 2 hours, and the peak reduction of blood pressure is achieved within 4-6 hours. The antihypertensive effect persists over 24 hours after dosing. During repeated dosing, the antihypertensive effect is substantially present within 2 weeks, and maximal effects are attained within 4 weeks and persist during long-term therapy.

Combined with hydrochlorothiazide, a significant additional reduction in blood pressure is achieved.

Abrupt withdrawal of Cuenca has not been associated with rebound hypertension or other adverse clinical events.

In hypertensive patients with type 2 diabetes and microalbuminuria, valsartan has been shown to reduce the urinary excretion of albumin. The MARVAL (Micro Albuminuria Reduction with Valsartan) study assessed the reduction in urinary albumin excretion

(UAE) with valsartan (80-160 mg/od) versus amlodipine (5-10 mg/od), in 332 type 2 diabetic patients (mean age: 58 years; 265 men) with microalbuminuria (valsartan: 58 µg/min; amlodipine: 55.4 µg/min), normal or high blood pressure and with preserved renal function (blood creatinine <120 µmol/l). At 24 weeks, UAE was reduced ( $p < 0.001$ ) by 42% (-24.2 µg/min; 95% CI: -40.4 to -19.1) with valsartan and approximately 3% (-1.7 µg/min; 95% CI: -5.6 to 14.9) with amlodipine despite similar rates of blood pressure reduction in both groups.

The Cuenca Reduction of Proteinuria (DROP) study further examined the efficacy of valsartan in reducing UAE in 391 hypertensive patients (BP=150/88 mmHg) with type 2 diabetes, albuminuria (mean=102 µg/min; 20-700 µg/min) and preserved renal function (mean serum creatinine = 80 µmol/l). Patients were randomized to one of 3 doses of valsartan (160, 320 and 640 mg/od) and treated for 30 weeks.

The purpose of the study was to determine the optimal dose of valsartan for reducing UAE in hypertensive patients with type 2 diabetes. At 30 weeks, the percentage change in UAE was significantly reduced by 36% from baseline with valsartan 160 mg (95%CI: 22 to 47%), and by 44% with valsartan 320 mg (95%CI: 31 to 54%). It was concluded that 160-320 mg of valsartan produced clinically relevant reductions in UAE in hypertensive patients with type 2 diabetes.

#### Recent myocardial infarction

The VALsartan In Acute myocardial iNfarcTion trial (VALIANT) was a randomised, controlled, multinational, double-blind study in 14,703 patients with acute myocardial infarction and signs, symptoms or radiological evidence of congestive heart failure and/or evidence of left ventricular systolic dysfunction (manifested as an ejection fraction  $\leq 40\%$  by radionuclide ventriculography or  $\leq 35\%$  by echocardiography or ventricular contrast angiography). Patients were randomised within 12 hours to 10 days after the onset of myocardial infarction symptoms to valsartan, captopril, or the combination of both. The mean treatment duration was two years. The primary endpoint was time to all-cause mortality.

Valsartan was as effective as captopril in reducing all-cause mortality after myocardial infarction. All-cause mortality was similar in the valsartan (19.9%), captopril (19.5%), and valsartan + captopril (19.3%) groups. Combining valsartan with captopril did not add further benefit over captopril alone. There was no difference between valsartan and captopril in all-cause mortality based on age, gender, race, baseline therapies or underlying disease. Valsartan was also effective in prolonging the time to and reducing cardiovascular mortality, hospitalisation for heart failure, recurrent myocardial infarction, resuscitated cardiac arrest, and non-fatal stroke (secondary composite endpoint).

The safety profile of valsartan was consistent with the clinical course of patients treated in the post-myocardial infarction setting. Regarding renal function, doubling of serum creatinine was observed in 4.2% of valsartan-treated patients, 4.8% of valsartan+captopril-treated patients, and 3.4% of captopril-treated patients. Discontinuations due to various types of renal dysfunction occurred in 1.1% of valsartan-treated patients, 1.3% in valsartan+captopril patients, and 0.8% of captopril patients. An assessment of renal function should be included in the evaluation of patients post-myocardial infarction.

There was no difference in all-cause mortality, cardiovascular mortality or morbidity when beta blockers were administered together with the combination of valsartan + captopril, valsartan alone, or captopril alone. Irrespective of treatment, mortality was lower in the group of patients treated with a beta blocker, suggesting that the known beta blocker benefit in this population was maintained in this trial.

#### Heart failure

Val-HeFT was a randomised, controlled, multinational clinical trial of valsartan compared with placebo on morbidity and mortality in 5,010 NYHA class II (62%), III (36%) and

IV (2%) heart failure patients receiving usual therapy with LVEF <40% and left ventricular internal diastolic diameter (LVIDD) >2.9 cm/m<sup>2</sup>. Baseline therapy included ACE inhibitors (93%), diuretics (86%), digoxin (67%) and beta blockers (36%). The mean duration of follow-up was nearly two years. The mean daily dose of Cuenca in Val-HeFT was 254 mg. The study had two primary endpoints: all cause mortality (time to death) and composite mortality and heart failure morbidity (time to first morbid event) defined as death, sudden death with resuscitation, hospitalisation for heart failure, or administration of intravenous inotropic or vasodilator agents for four hours or more without hospitalisation.

All cause mortality was similar (p=NS) in the valsartan (19.7%) and placebo (19.4%) groups. The primary benefit was a 27.5% (95% CI: 17 to 37%) reduction in risk for time to first heart failure hospitalisation (13.9% vs. 18.5%). Results appearing to favour placebo (composite mortality and morbidity was 21.9% in placebo vs. 25.4% in valsartan group) were observed for those patients receiving the triple combination of an ACE inhibitor, a beta blocker and valsartan.

In a subgroup of patients not receiving an ACE inhibitor (n=366), the morbidity benefits were greatest. In this subgroup all-cause mortality was significantly reduced with valsartan compared to placebo by 33% (95% CI: -6% to 58%) (17.3% valsartan vs. 27.1% placebo) and the composite mortality and morbidity risk was significantly reduced by 44% (24.9% valsartan vs. 42.5% placebo).

In patients receiving an ACE inhibitor without a beta-blocker, all cause mortality was similar (p=NS) in the valsartan (21.8%) and placebo (22.5%) groups. Composite mortality and morbidity risk was significantly reduced by 18.3% (95% CI: 8% to 28%) with valsartan compared with placebo (31.0% vs. 36.3%).

In the overall Val-HeFT population, valsartan treated patients showed significant improvement in NYHA class, and heart failure signs and symptoms, including dyspnoea, fatigue, oedema and rales compared to placebo. Patients treated with valsartan had a better quality of life as demonstrated by change in the Minnesota Living with Heart Failure Quality of Life score from baseline at endpoint than placebo. Ejection fraction in valsartan treated patients was significantly increased and LVIDD significantly reduced from baseline at endpoint compared to placebo.

### Paediatric population

#### Hypertension

The antihypertensive effect of valsartan have been evaluated in four randomized, double-blind clinical studies in 561 paediatric patients from 6 to 18 years of age and 165 paediatric patients 1 to 6 years of age. Renal and urinary disorders, and obesity were the most common underlying medical conditions potentially contributing to hypertension in the children enrolled in these studies.

#### Clinical experience in children at or above 6 years of age

In a clinical study involving 261 hypertensive paediatric patients 6 to 16 years of age, patients who weighed <35 kg received 10, 40 or 80 mg of valsartan tablets daily (low, medium and high doses), and patients who weighed ≥35 kg received 20, 80, and 160 mg of valsartan tablets daily (low, medium and high doses). At the end of 2 weeks, valsartan reduced both systolic and diastolic blood pressure in a dose-dependent manner.

Overall, the three dose levels of valsartan (low, medium and high) significantly reduced systolic blood pressure by 8, 10, 12 mm Hg from the baseline, respectively. Patients were re-randomized to either continue receiving the same dose of valsartan or were switched to

placebo. In patients who continued to receive the medium and high doses of valsartan, systolic blood pressure at trough was -4 and -7 mm Hg lower than patients who received the placebo treatment. In patients receiving the low dose of valsartan, systolic blood pressure at trough was similar to that of patients who received the placebo treatment. Overall, the dose-dependent antihypertensive effect of valsartan was consistent across all the demographic subgroups.

In another clinical study involving 300 hypertensive paediatric patients 6 to 18 years of age, eligible patients were randomized to receive valsartan or enalapril tablets for 12 weeks. Children weighing between  $\geq 18$  kg and  $< 35$  kg received valsartan 80 mg or enalapril 10 mg; those between  $\geq 35$  kg and  $< 80$  kg received valsartan 160 mg or enalapril 20 mg; those  $\geq 80$  kg received valsartan 320 mg or enalapril 40 mg. Reductions in systolic blood pressure were comparable in patients receiving valsartan (15 mmHg) and enalapril (14 mm Hg) (non-inferiority p-value  $< 0.0001$ ). Consistent results were observed for diastolic blood pressure with reductions of 9.1 mmHg and 8.5 mmHg with valsartan and enalapril, respectively.

#### Clinical experience in children less than 6 years of age

Two clinical studies were conducted in patients aged 1 to 6 years with 90 and 75 patients, respectively. No children below the age of 1 year were enrolled in these studies. In the first study, the efficacy of valsartan was confirmed compared to placebo but a dose-response could not be demonstrated. In the second study, higher doses of valsartan were associated with greater BP reductions, but the dose response trend did not achieve statistical significance and the treatment difference compared to placebo was not significant.

Because of these inconsistencies, valsartan is not recommended in this age group (see section 4.8).

The European Medicines Agency has waived the obligation to submit the results of studies with Cuenca in all subsets of the paediatric population in heart failure and heart failure after recent myocardial infarction. See section 4.2 for information on paediatric use.

## **5.2 Pharmacokinetic properties**

### Absorption:

Following oral administration of valsartan alone, peak plasma concentrations of valsartan are reached in 2– 4 hours with tablets and 1–2 hours with solution formulation. Mean absolute bioavailability is 23% and 39% with tablets and solution formulation, respectively. Food decreases exposure (as measured by AUC) to valsartan by about 40% and peak plasma concentration ( $C_{max}$ ) by about 50%, although from about 8 h post dosing plasma valsartan concentrations are similar for the fed and fasted groups. This reduction in AUC is not, however, accompanied by a clinically significant reduction in the therapeutic effect, and valsartan can therefore be given either with or without food.

### Distribution:

The steady-state volume of distribution of valsartan after intravenous administration is about 17 litres, indicating that valsartan does not distribute into tissues extensively. Valsartan is highly bound to serum proteins (94–97%), mainly serum albumin.

### Biotransformation:

Valsartan is not biotransformed to a high extent as only about 20% of dose is recovered as metabolites. A hydroxy metabolite has been identified in plasma at low concentrations (less than 10% of the valsartan AUC). This metabolite is pharmacologically inactive.

#### Elimination:

Valsartan shows multiexponential decay kinetics ( $t_{1/2\alpha} < 1$  h and  $t_{1/2\beta}$  about 9 h). Valsartan is primarily eliminated by biliary excretion in faeces (about 83% of dose) and renally in urine (about 13% of dose), mainly as unchanged drug. Following intravenous administration, plasma clearance of valsartan is about

2 l/h and its renal clearance is 0.62 l/h (about 30% of total clearance). The half-life of valsartan is 6 hours.

#### In heart failure patients:

The average time to peak concentration and elimination half-life of valsartan in heart failure patients are similar to that observed in healthy volunteers. AUC and  $C_{max}$  values of valsartan are almost proportional with increasing dose over the clinical dosing range (40 to 160 mg twice a day). The average accumulation factor is about 1.7. The apparent clearance of valsartan following oral administration is approximately 4.5 l/h. Age does not affect the apparent clearance in heart failure patients.

#### Special populations

##### Elderly

A somewhat higher systemic exposure to valsartan was observed in some elderly subjects than in young subjects; however, this has not been shown to have any clinical significance.

##### Impaired renal function

As expected for a compound where renal clearance accounts for only 30% of total plasma clearance, no correlation was seen between renal function and systemic exposure to valsartan. Dose adjustment is therefore not required in patients with renal impairment (creatinine clearance  $>10$  ml/min). There is currently no experience on the safe use in patients with a creatinine clearance  $<10$  ml/min and patients undergoing dialysis, therefore valsartan should be used with caution in these patients (see sections 4.2 and 4.4).

Valsartan is highly bound to plasma protein and is unlikely to be removed by dialysis.

##### Hepatic impairment

Approximately 70% of the dose absorbed is eliminated in the bile, essentially in the unchanged form. Valsartan does not undergo any noteworthy biotransformation. A doubling of exposure (AUC) was observed in patients with mild to moderate hepatic impairment compared to healthy subjects. However, no correlation was observed between plasma valsartan concentration versus degree of hepatic dysfunction. Cuenca has not been studied in patients with severe hepatic dysfunction (see sections 4.2, 4.3 and 4.4).

##### Paediatric population

In a study of 26 paediatric hypertensive patients (aged 1 to 16 years) given a single dose of a suspension of valsartan (mean: 0.9 to 2 mg/kg, with a maximum dose of 80 mg), the clearance (litres/h/kg) of valsartan was comparable across the age range of 1 to 16 years and similar to that of adults receiving the same formulation.

#### Impaired renal function

Use in paediatric patients with a creatinine clearance <30 ml/min and paediatric patients undergoing dialysis has not been studied, therefore valsartan is not recommended in these patients. No dose adjustment is required for paediatric patients with a creatinine clearance >30 ml/min. Renal function and serum potassium should be closely monitored (see sections 4.2 and 4.4).

### **5.3 Preclinical safety data**

Non-clinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, repeated dose toxicity, genotoxicity, carcinogenic potential. In rats, maternally toxic doses (600 mg/kg/day) during the last days of gestation and lactation led to lower survival, lower weight gain and delayed development (pinna detachment and ear-canal opening) in the offspring (see section 4.6). These doses in rats (600 mg/kg/day) are approximately 18 times the maximum recommended human dose on a  $\text{mg}/\text{m}^2$  basis (calculations assume an oral dose of 320 mg/day and a 60-kg patient). In non-clinical safety studies, high doses of valsartan (200 to 600 mg/kg body weight) caused in rats a reduction of red blood cell parameters (erythrocytes, haemoglobin, haematocrit) and evidence of changes in renal haemodynamics (slightly raised plasma urea, and renal tubular hyperplasia and basophilia in males). These doses in rats (200 and 600 mg/kg/day) are approximately 6 and 18 times the maximum recommended human dose on a  $\text{mg}/\text{m}^2$  basis (calculations assume an oral dose of 320 mg/day and a 60-kg patient). In marmosets at similar doses, the changes were similar though more severe, particularly in the kidney where the changes developed to a nephropathy which included raised urea and creatinine. Hypertrophy of the renal juxtaglomerular cells was also seen in both species. All changes were considered to be caused by the pharmacological action of valsartan which produces prolonged hypotension, particularly in marmosets. For therapeutic doses of valsartan in humans, the hypertrophy of the renal juxtaglomerular cells does not seem to have any relevance.

#### Paediatric population

Daily oral dosing of neonatal/juvenile rats (from a postnatal day 7 to postnatal day 70) with valsartan at doses as low as 1 mg/kg/day (about 10-35% of the maximum recommended paediatric dose of 4 mg/kg/day on systemic exposure basis) produced persistent, irreversible kidney damage. These effects above mentioned represent an expected exaggerated pharmacological effect of angiotensin converting enzyme inhibitors and angiotensin II type 1 blockers; such effects are observed if rats are treated during the first 13 days of life. This period coincides with 36 weeks of gestation in humans, which could occasionally extend up to 44 weeks after conception in humans. The rats in the juvenile valsartan study were dosed up to day 70, and effects on renal maturation (postnatal 4-6 weeks) cannot be excluded. Functional renal maturation is an ongoing process within the first year of life in humans. Consequently, a clinical relevance in children <1 year of age cannot be excluded, while preclinical data do not indicate a safety concern for children older than 1 year.

## **6 PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

#### *Tablet core*

Cellulose, microcrystalline (E 460)

Silica, colloidal anhydrous (E 551)

Sorbitol (E-420)

Magnesium carbonate (E 504)

Maize starch, pregelatinised

Povidone K-25 (E 1201)

Sodium stearyl fumarate

Sodium lauryl sulphate

Crospovidone Type A (E 1202)

#### *Film coating*

Lactose monohydrate

Hypromellose (E 464)

Titanium dioxide (E 171)

Macrogol 4000

### **6.2 Incompatibilities**

Not applicable

### **6.3 Shelf life**

2 years

### **6.4 Special precautions for storage**

Do not store above 30°C. Store in the original package in order to protect from moisture

### **6.5 Nature and contents of container**

PVC/PE/PVDC/aluminium blister.

Pack sizes: 7, 14, 28, 56, 98, 280 film-coated tablets

Not all pack sizes may be marketed.

**6.6 Special precautions for disposal**

No special requirements.

Any unused product or waste should be disposed of in accordance with local requirements.

**7 MARKETING AUTHORISATION HOLDER**

Laboratorios Liconsa, S.A.

Gran Vía Carlos III, 98, 7th floor

08028 Barcelona, SPAIN

**8 MARKETING AUTHORISATION NUMBER(S)**

PL 23218/0043

**9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

31/01/2013

**10 DATE OF REVISION OF THE TEXT**

31/01/2013