

Celecol® 200 mg Film-coated Tablets

(celiprolol hydrochloride)

Your medicine is known by the above name, but will be referred to as Celecol throughout this leaflet. Celecol is also available as the 400 mg strength.

Patient Information Leaflet

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

- 1) What Celecol is and what it is used for
- 2) What you need to know before you take Celecol
- 3) How to take Celecol
- 4) Possible side effects
- 5) How to store Celecol
- 6) Contents of the pack and other information

1) What Celecol is and what it is used for

What Celecol is

The name of your medicine is Celecol 200 mg film-coated tablets (called Celecol throughout this leaflet). Celecol contains a medicine called celiprolol hydrochloride. This belongs to a group of medicines called beta-blockers.

How Celecol works

It works by slowing your heart rate or lowering your blood pressure.

What Celecol is used for

It is used to treat high blood pressure (hypertension).

2) What you need to know before you take Celecol

Do not take Celecol:

- * if you are allergic to celiprolol or any of the other ingredients of this medicine (listed in section 6).
Signs of an allergic reaction include: a rash, swallowing or breathing problems, swelling of your lips, face, throat or tongue.
- * if you have or have had asthma or wheezing.
- * if you have a very slow heartbeat (bradycardia).
- * if you have heart problems – where you have difficulty breathing and swollen ankles.
- * if you have kidney problems.
- * if you have a tumour of the adrenal gland (phaeochromocytoma) and your blood pressure has not yet been treated with a type of medicine known as an alpha blocker, such as doxazosin or terazosin.
- * if you have low blood pressure (hypotension).
- * if you have abnormal levels of acid in your blood (metabolic acidosis).
- * if you have been diagnosed by your doctor with late stages of a peripheral arterial disease (such as Raynaud's disease). Symptoms include persistent cramping pain in legs even at rest, and development of non-healing ulcers over fingers and/or toes.

Do not take this medicine if any of the above apply to you. If you are not sure, talk to your doctor or pharmacist before taking Celecol.

Warnings and precautions

Talk to your doctor or pharmacist before taking Celecol:

- if you have any heart problems.
- if you have breathing problems due to long term lung problems (called bronchitis or emphysema).
- if you have diabetes.
- if you have liver problems.
- if you have a tumour of the adrenal gland (phaeochromocytoma) and your blood pressure is not yet controlled by a type of medicine known as an alpha blocker, such as doxazosin or terazosin.
- if you have an overactive thyroid gland, as celiprolol can hide some of the symptoms associated with this condition.
- if you have a history of severe allergies, including anaphylactic reactions.
- if you have ever suffered from psoriasis.
- if you suffer from a type of chest pain called Prinzmetal's angina.
- if you suffer from poor circulation (such as Raynaud's disease).
Symptoms include fingers and/ or toes changing colour when cold and tingling or painful on warming up. Your doctor may closely monitor you while taking this medicine.

If you are not sure if any of the above apply to you, talk to your doctor or pharmacist before taking Celecol.

Operations or anaesthetics

Tell your doctor or dentist you are taking Celecol if you are going to have an anaesthetic or an operation (including dental surgery).

Other medicines and Celecol

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines, including medicines obtained without a prescription and herbal medicines. This is because Celecol can affect the way some other medicines work. Also, some other medicines can affect the way Celecol works.

In particular tell your doctor if you are taking any of the following:

- Insulin or tablets to treat diabetes.

- Medicines, including some cough and cold medicines, containing ephedrine, pseudoephedrine or adrenaline.
 - Medicines to treat depression called MAOIs (monoamine oxidase inhibitors) or have taken them in the last 2 weeks. MAOIs are medicines such as meclobemide, phenelzine or tranylcypromine.
 - Medicines used for mental problems (antipsychotic medicines) – such as chlorpromazine or other phenothiazines.
 - Medicines used for asthma (such as theophylline).
 - Medicines for chest pain (angina) – such as verapamil, nifedipine and diltiazem.
 - Medicines used for heart problems – such as lidocaine (also used as a local anaesthetic) amiodarone, digitalis, diltiazem, disopyramide, nifedipine, quinidine and verapamil (also used for severe allergies).
 - Diltiazem as it may lead to increased risk of depression.
 - Non-Steroidal Anti-inflammatory Drugs (NSAIDs) – used to relieve pain such as ibuprofen and indometacin.
 - Medicines used to lower blood pressure – such as chlorthalidone, hydrochlorothiazide and clonidine.
 - Medicines used for the prevention of malaria such as mefloquine.
 - Some antibiotics or antifungal medicines e.g. rifampicin, erythromycin, clarithromycin, ketoconazole, itraconazole.
 - Medicines used to alter the way your immune system works e.g. fingolimod, ciclosporin.
 - Medicines obtained without a prescription e.g. St John's Wort.
- If you are not sure, talk to your doctor or pharmacist before taking Celecol.

Celecol with food and drink

- Celecol should be taken preferably first thing in the morning, 30 minutes before food or two hours after a meal.
- Citrus fruit juices should be avoided when taking this medicine as they can decrease the absorption of celiprolol.

Pregnancy and breast-feeding

If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine.

If your doctor cannot find a safer alternative you may be given Celecol. If you take Celecol during pregnancy, particularly in the last 3 months, you should tell your midwife. This is because your baby may be affected when it is born. Any effects usually begin on the first day after birth and can include lower than normal levels of glucose in the blood (hypoglycaemia), slow heartbeat, heart failure or having trouble breathing. If your baby has any of these side effects when it is born and you are worried, tell your doctor or midwife.

You should not breast-feed if you are taking Celecol. This is because small amounts may pass into the mother's milk. If you are breast-feeding or planning to breast-feed, ask your doctor or pharmacist for advice before taking this medicine.

Ask your doctor or pharmacist for advice before taking any medicine if you are pregnant or breast-feeding.

Driving and using machines

You may feel tired or dizzy whilst taking Celecol. If this happens, do not drive or use any tools or machines.

Drug tests in sport

Celecol may give a positive result if you have a blood or urine test to check for the presence of drugs.

Information about one of the ingredients of this medicine

This medicine contains less than 1 mmol sodium (23 mg) per film-coated tablet, that is to say essentially 'sodium-free'.

3) How to take Celecol

Always take this medicine exactly as your doctor has told you. The dose of Celecol will depend on your needs and the illness being treated. Check with your doctor or pharmacist if you are not sure.

Taking this medicine

- Swallow the tablets whole with a drink of water.
- Take your medicine at the same time each day, preferably first thing in the morning, 30 minutes before food, or 2 hours after a meal.

How much to take

Adults

- The usual starting dose is 200 mg once daily.
- The maximum dose is 400 mg once daily.

Children

Celecol should not be given to children.

Elderly patients and patients with kidney problems

Your doctor may decide to lower your dose of Celecol if necessary.

If you take more Celecol than you should

Tell your doctor or go to the nearest hospital casualty department straight away. Remember to take with you any tablets that are left and the pack.

This is so the doctor knows what you have taken.

The following effects may happen: feeling dizzy or weak, difficulty breathing or wheezing.

If you forget to take Celecol

If you forget to take a dose, take it as soon as you remember, unless it is time for your next dose. **Do not** take a double dose to make up for a forgotten dose.

If you stop taking Celectol

Keep taking your medicine until your doctor tells you to stop. Do not stop taking Celectol just because you feel better. If you stop taking Celectol, your illness may get worse. When your doctor says that you can stop taking Celectol, you need to do this gradually. Your doctor will help you do this. If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4) Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Stop taking Celectol and see a doctor or go to a hospital straight away if:

- you get swelling of the hands, feet, ankles, face, lips or throat which may cause difficulty in swallowing or breathing. You could also notice an itchy, lumpy rash (hives) or nettle rash (urticaria). This may mean you are having an **allergic reaction** to Celectol.
- you get flu-like symptoms, swollen or painful joints, increased sensitivity to sunlight, hair loss, abdominal pain, anaemia. This may mean you have a condition called Systemic Lupus Erythematosus (SLE). These symptoms usually go away after Celectol has been stopped.

Tell your doctor as soon as possible if you have any of the following side effects:

Common side effects (may affect up to 1 in 10 people)

- Feeling shaky (tremor), tired, drowsy or weak
- Stomach pain
- Depression
- Hot flushes
- Skin rashes

Uncommon side effects (may affect up to 1 in 100 people)

- Slow heart rate or very fast, uneven or forceful heartbeat (palpitations)
- Feeling dizzy or weak. These could be signs of low blood pressure.
- Difficulty breathing or shortness of breath
- Difficulty sleeping (insomnia)

Not known (frequency cannot be estimated from available data)

- Pain in your chest and feeling weak or dizzy. These could be signs of a heart problem.
- Slow or irregular heart rate
- Changes in behaviour and feeling that you are losing touch with reality, or are unable to think or judge things clearly. You may also feel more aggressive than usual. These could be signs of a serious mental health problem (psychoses).
- Seeing or hearing things that are not there (hallucinations), confusion, nightmares
- Problems with your eyesight such as blurred vision
- Dryness or redness of the eye
- Diarrhoea
- Appearance of scaly skin rash (psoriasis-like rash) or worsening of existing psoriasis
- Low blood sugar levels (hypoglycaemia). You may feel a sense of nervousness, shaky or sweaty.
- High blood sugar levels (hyperglycaemia). You may be passing large amounts of urine, have excessive thirst and a dry mouth or skin. You may also get frequent infections such as thrush.
- Hair loss (alopecia)

Tell your doctor or pharmacist if any of the following side effects get serious or last longer than a few days:

Common side effects (may affect 1 in 10 people)

- Feeling sick (nausea) or being sick (vomiting), dry mouth
- Feeling dizzy or light-headed, headache
- Sweating more than usual
- Worsening of existing circulation disorders of the extremities. Symptoms may include a cold or numb feeling in your fingers or toes (Raynaud's disease or syndrome), or cramping pain in the legs (claudication).
- Unusual skin sensations such as numbness, tingling, pricking, burning or creeping on the skin
- Difficulty in getting or maintaining an erection, or in ejaculating (impotence)

Uncommon side effects (may affect up to 1 in 100 people)

- Muscle spasms

Not known (frequency cannot be estimated from available data)

- Fainting
- Breathing problems such as asthma or productive cough
- Lack of interest in sexual activities (libido)
- Pain in your joints (arthralgia)

Blood tests

- The results of some blood tests may show higher levels of liver enzymes than normal.
- Blood tests may show abnormal levels of glucose in the blood.
- Blood tests may show abnormal levels of autoimmune factors.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the **Google Play** or **Apple App Store**. By reporting side effects you can help provide more information on the safety of this medicine.

5) How to store Celectol

- **Keep this medicine out of the sight and reach of children.**
- Do not take this medicine after the expiry date which is stated on the carton and blister label after 'EXP'. The expiry date refers to the last day of that month.
- Do not store above 25°C.
- If the medicine becomes discoloured or show any other signs of deterioration, you should seek the advice of your pharmacist who will tell you what to do.
- Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6) Contents of the pack and other information

What Celectol contain

The active substance is celiprolol hydrochloride. Each film-coated tablet contains 200 mg of celiprolol hydrochloride.

The other ingredients are mannitol (E421), microcrystalline cellulose, carboxymethylcellulose sodium and magnesium stearate.

Film-coating:

Opadry Y-1-7000 (white) which contains hypromellose (E464), titanium dioxide (E171) and macrogol 400.

Opadry YS-1R-7006 (clear) which contains hypromellose (E464), macrogol 400 and macrogol 6000.

What Celectol looks like and contents of the pack

Celectol 200 mg film-coated tablets are white, film-coated, biconvex, heart shaped and engraved with the Celectol logo on one side, and 200 and a breakline on the other side.

Celectol film-coated tablets are supplied in blister packs of 28 tablets.

PL 46420/0605 Celectol 200 mg Film-coated Tablets **POM**

Who makes and repackages your medicine?

Your medicine is manufactured by Opella Healthcare International SAS, 56, Route de Choisy, 60200 Compiègne, France.

Procured from within the EU and repackaged by the Product Licence holder: Suerte Pharma Ltd, 4/5 Northolt Trading Estate, Belvue Road, Northolt, Middlesex, UB5 5QS.

Leaflet Date: 27.10.2025

Celectol is a registered trademark of Cheplapharm Arzneimittel GmbH, Germany.

For any information about this medicine, or to request for a copy of this leaflet in Braille, large print or audio, please call the PL holder: Suerte Pharma Ltd on 020 8839 3000.

Celiprolol Hydrochloride 200 mg Film-coated Tablets

Your medicine is known by the above name, but will be referred to as Celiprolol Hydrochloride Tablets throughout this leaflet. Celiprolol Hydrochloride Tablets is also available as the 400 mg strength.

Patient Information Leaflet

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

- 1) What Celiprolol Hydrochloride Tablets is and what it is used for
- 2) What you need to know before you take Celiprolol Hydrochloride Tablets
- 3) How to take Celiprolol Hydrochloride Tablets
- 4) Possible side effects
- 5) How to store Celiprolol Hydrochloride Tablets
- 6) Contents of the pack and other information

1) What Celiprolol Hydrochloride Tablets is and what it is used for

What Celiprolol Hydrochloride Tablets is

The name of your medicine is Celiprolol Hydrochloride Tablets 200 mg film-coated tablets (called Celiprolol Hydrochloride Tablets throughout this leaflet). Celiprolol Hydrochloride Tablets contains a medicine called celiprolol hydrochloride. This belongs to a group of medicines called beta-blockers.

How Celiprolol Hydrochloride Tablets works

It works by slowing your heart rate or lowering your blood pressure.

What Celiprolol Hydrochloride Tablets is used for

It is used to treat high blood pressure (hypertension).

2) What you need to know before you take Celiprolol Hydrochloride Tablets

Do not take Celiprolol Hydrochloride Tablets:

- × if you are allergic to celiprolol or any of the other ingredients of this medicine (listed in section 6).
Signs of an allergic reaction include: a rash, swallowing or breathing problems, swelling of your lips, face, throat or tongue.
- × if you have or have had asthma or wheezing.
- × if you have a very slow heartbeat (bradycardia).
- × if you have heart problems – where you have difficulty breathing and swollen ankles.
- × if you have kidney problems.
- × if you have a tumour of the adrenal gland (phaeochromocytoma) and your blood pressure has not yet been treated with a type of medicine known as an alpha blocker, such as doxazosin or terazosin.
- × if you have low blood pressure (hypotension).
- × if you have abnormal levels of acid in your blood (metabolic acidosis).
- × if you have been diagnosed by your doctor with late stages of a peripheral arterial disease (such as Raynaud's disease). Symptoms include persistent cramping pain in legs even at rest, and development of non-healing ulcers over fingers and/or toes.

Do not take this medicine if any of the above apply to you. If you are not sure, talk to your doctor or pharmacist before taking Celiprolol Hydrochloride Tablets.

Warnings and precautions

Talk to your doctor or pharmacist before taking Celiprolol Hydrochloride Tablets:

- if you have any heart problems.
- if you have breathing problems due to long term lung problems (called bronchitis or emphysema).
- if you have diabetes.
- if you have liver problems.
- if you have a tumour of the adrenal gland (phaeochromocytoma) and your blood pressure is not yet controlled by a type of medicine known as an alpha blocker, such as doxazosin or terazosin.
- if you have an overactive thyroid gland, as celiprolol can hide some of the symptoms associated with this condition.
- if you have a history of severe allergies, including anaphylactic reactions.
- if you have ever suffered from psoriasis.
- if you suffer from a type of chest pain called Prinzmetal's angina.
- if you suffer from poor circulation (such as Raynaud's disease).
Symptoms include fingers and/ or toes changing colour when cold and tingling or painful on warming up. Your doctor may closely monitor you while taking this medicine.

If you are not sure if any of the above apply to you, talk to your doctor or pharmacist before taking Celiprolol Hydrochloride Tablets.

Operations or anaesthetics

Tell your doctor or dentist you are taking Celiprolol Hydrochloride Tablets if you are going to have an anaesthetic or an operation (including dental surgery).

Other medicines and Celiprolol Hydrochloride Tablets

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines, including medicines obtained without a prescription and herbal medicines. This is because Celiprolol Hydrochloride Tablets can affect the way some other medicines work. Also, some other medicines can affect the way Celiprolol Hydrochloride Tablets works. In particular tell your doctor if you are taking any of the following:

- Insulin or tablets to treat diabetes.
- Medicines, including some cough and cold medicines, containing ephedrine, pseudoephedrine or adrenaline.
- Medicines to treat depression called MAOIs (monoamine oxidase inhibitors) or have taken them in the last 2 weeks. MAOIs are medicines such as meclobemide, phenelzine or tranylcypromine.
- Medicines used for mental problems (antipsychotic medicines) – such as chlorpromazine or other phenothiazines.
- Medicines used for asthma (such as theophylline).
- Medicines for chest pain (angina) – such as verapamil, nifedipine and diltiazem.
- Medicines used for heart problems – such as lidocaine (also used as a local anaesthetic) amiodarone, digitalis, diltiazem, disopyramide, nifedipine, quinidine and verapamil (also used for severe allergies).
- Diltiazem as it may lead to increased risk of depression.
- Non-Steroidal Anti-inflammatory Drugs (NSAIDs) – used to relieve pain such as ibuprofen and indometacin.
- Medicines used to lower blood pressure – such as chlorthalidone, hydrochlorothiazide and clonidine.
- Medicines used for the prevention of malaria such as mefloquine.
- Some antibiotics or antifungal medicines e.g. rifampicin, erythromycin, clarithromycin, ketoconazole, itraconazole.
- Medicines used to alter the way your immune system works e.g. fingolimod, ciclosporin.
- Medicines obtained without a prescription e.g. St John's Wort.

If you are not sure, talk to your doctor or pharmacist before taking Celiprolol Hydrochloride Tablets.

Celiprolol Hydrochloride Tablets with food and drink

- Celiprolol Hydrochloride Tablets should be taken preferably first thing in the morning, 30 minutes before food or two hours after a meal.
- Citrus fruit juices should be avoided when taking this medicine as they can decrease the absorption of celiprolol.

Pregnancy and breast-feeding

If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. If your doctor cannot find a safer alternative you may be given Celiprolol Hydrochloride Tablets. If you take Celiprolol Hydrochloride Tablets during pregnancy, particularly in the last 3 months, you should tell your midwife. This is because your baby may be affected when it is born. Any effects usually begin on the first day after birth and can include lower than normal levels of glucose in the blood (hypoglycaemia), slow heartbeat, heart failure or having trouble breathing. If your baby has any of these side effects when it is born and you are worried, tell your doctor or midwife. You should not breast-feed if you are taking Celiprolol Hydrochloride Tablets. This is because small amounts may pass into the mother's milk. If you are breast-feeding or planning to breast-feed, ask your doctor or pharmacist for advice before taking this medicine. Ask your doctor or pharmacist for advice before taking any medicine if you are pregnant or breast-feeding.

Driving and using machines

You may feel tired or dizzy whilst taking Celiprolol Hydrochloride Tablets. If this happens, do not drive or use any tools or machines.

Drug tests in sport

Celiprolol Hydrochloride Tablets may give a positive result if you have a blood or urine test to check for the presence of drugs.

Information about one of the ingredients of this medicine

This medicine contains less than 1 mmol sodium (23 mg) per film-coated tablet, that is to say essentially 'sodium-free'.

3) How to take Celiprolol Hydrochloride Tablets

Always take this medicine exactly as your doctor has told you. The dose of Celiprolol Hydrochloride Tablets will depend on your needs and the illness being treated. Check with your doctor or pharmacist if you are not sure.

Taking this medicine

- Swallow the tablets whole with a drink of water.
- Take your medicine at the same time each day, preferably first thing in the morning, 30 minutes before food, or 2 hours after a meal.

How much to take

Adults

- The usual starting dose is 200 mg once daily.
- The maximum dose is 400 mg once daily.

Children

Celiprolol Hydrochloride Tablets should not be given to children.

Elderly patients and patients with kidney problems

Your doctor may decide to lower your dose of Celiprolol Hydrochloride Tablets if necessary.

If you take more Celiprolol Hydrochloride Tablets than you should

Tell your doctor or go to the nearest hospital casualty department straight away. Remember to take with you any tablets that are left and the pack. This is so the doctor knows what you have taken. The following effects may happen: feeling dizzy or weak, difficulty breathing or wheezing.

If you forget to take Celiprolol Hydrochloride Tablets

If you forget to take a dose, take it as soon as you remember, unless it is time for your next dose. **Do not** take a double dose to make up for a forgotten dose.

If you stop taking Celiprolol Hydrochloride Tablets

Keep taking your medicine until your doctor tells you to stop. Do not stop taking Celiprolol Hydrochloride Tablets just because you feel better. If you stop taking Celiprolol Hydrochloride Tablets, your illness may get worse. When your doctor says that you can stop taking Celiprolol Hydrochloride Tablets, you need to do this gradually. Your doctor will help you do this. If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4) Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Stop taking Celiprolol Hydrochloride Tablets and see a doctor or go to a hospital straight away if:

- you get swelling of the hands, feet, ankles, face, lips or throat which may cause difficulty in swallowing or breathing. You could also notice an itchy, lumpy rash (hives) or nettle rash (urticaria). This may mean you are having an **allergic reaction** to Celiprolol Hydrochloride Tablets.
- you get flu-like symptoms, swollen or painful joints, increased sensitivity to sunlight, hair loss, abdominal pain, anaemia. This may mean you have a condition called Systemic Lupus Erythematosus (SLE). These symptoms usually go away after Celiprolol Hydrochloride Tablets has been stopped.

Tell your doctor as soon as possible if you have any of the following side effects:

Common side effects (may affect up to 1 in 10 people)

- Feeling shaky (tremor), tired, drowsy or weak
- Stomach pain
- Depression
- Hot flushes
- Skin rashes

Uncommon side effects (may affect up to 1 in 100 people)

- Slow heart rate or very fast, uneven or forceful heartbeat (palpitations)
- Feeling dizzy or weak. These could be signs of low blood pressure.
- Difficulty breathing or shortness of breath
- Difficulty sleeping (insomnia)

Not known (frequency cannot be estimated from available data)

- Pain in your chest and feeling weak or dizzy. These could be signs of a heart problem.
- Slow or irregular heart rate
- Changes in behaviour and feeling that you are losing touch with reality, or are unable to think or judge things clearly. You may also feel more aggressive than usual. These could be signs of a serious mental health problem (psychoses).
- Seeing or hearing things that are not there (hallucinations), confusion, nightmares
- Problems with your eyesight such as blurred vision
- Dryness or redness of the eye
- Diarrhoea
- Appearance of scaly skin rash (psoriasis-like rash) or worsening of existing psoriasis
- Low blood sugar levels (hypoglycaemia). You may feel a sense of nervousness, shaky or sweaty.
- High blood sugar levels (hyperglycaemia). You may be passing large amounts of urine, have excessive thirst and a dry mouth or skin. You may also get frequent infections such as thrush.
- Hair loss (alopecia)

Tell your doctor or pharmacist if any of the following side effects get serious or last longer than a few days:

Common side effects (may affect 1 in 10 people)

- Feeling sick (nausea) or being sick (vomiting), dry mouth
- Feeling dizzy or light-headed, headache
- Sweating more than usual
- Worsening of existing circulation disorders of the extremities. Symptoms may include a cold or numb feeling in your fingers or toes (Raynaud's disease or syndrome), or cramping pain in the legs (claudication).
- Unusual skin sensations such as numbness, tingling, pricking, burning or creeping on the skin
- Difficulty in getting or maintaining an erection, or in ejaculating (impotence)

Uncommon side effects (may affect up to 1 in 100 people)

- Muscle spasms

Not known (frequency cannot be estimated from available data)

- Fainting
- Breathing problems such as asthma or productive cough
- Lack of interest in sexual activities (libido)
- Pain in your joints (arthralgia)

Blood tests

- The results of some blood tests may show higher levels of liver enzymes than normal.
- Blood tests may show abnormal levels of glucose in the blood.
- Blood tests may show abnormal levels of autoimmune factors.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the **Google Play** or **Apple App Store**. By reporting side effects you can help provide more information on the safety of this medicine.

5) How to store Celiprolol Hydrochloride Tablets

- **Keep this medicine out of the sight and reach of children.**
- Do not take this medicine after the expiry date which is stated on the carton and blister label after 'EXP'. The expiry date refers to the last day of that month.
- Do not store above 25°C.
- If the medicine becomes discoloured or show any other signs of deterioration, you should seek the advice of your pharmacist who will tell you what to do.
- Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6) Contents of the pack and other information

What Celiprolol Hydrochloride Tablets contain

The active substance is celiprolol hydrochloride. Each film-coated tablet contains 200 mg of celiprolol hydrochloride.

The other ingredients are mannitol (E421), microcrystalline cellulose, carboxymethylcellulose sodium and magnesium stearate.

Film-coating:

Opadry Y-1-7000 (white) which contains hypromellose (E464), titanium dioxide (E171) and macrogol 400.

Opadry YS-1R-7006 (clear) which contains hypromellose (E464), macrogol 400 and macrogol 6000.

What Celiprolol Hydrochloride Tablets looks like and contents of the pack

Celiprolol Hydrochloride 200 mg film-coated tablets are white, film-coated, biconvex, heart shaped and engraved with the Celectol logo on one side, and 200 and a breakline on the other side.

Celiprolol Hydrochloride film-coated tablets are supplied in blister packs of 28 tablets.

PL 46420/0605 Celiprolol Hydrochloride 200 mg Film-coated Tablets

POM

Who makes and repackages your medicine?

Your medicine is manufactured by Opella Healthcare International SAS, 56, Route de Choisy, 60200 Compiègne, France.

Procured from within the EU and repackaged by the Product Licence holder: Suerte Pharma Ltd, 4/5 Northolt Trading Estate, Belvue Road, Northolt, Middlesex, UB5 5QS.

Leaflet Date: 27.10.2025

For any information about this medicine, or to request for a copy of this leaflet in Braille, large print or audio, please call the PL holder: Suerte Pharma Ltd on 020 8839 3000.