

Package leaflet: Information for the patient

Malarone®

250mg/100mg film-coated tablets

(atovaquone/proguanil hydrochloride)

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

The name of your medicine is Malarone 250mg/100mg film-coated tablets but will be referred to as Malarone throughout this leaflet.

What is in this leaflet

1. What Malarone is and what it is used for
2. What you need to know before you take Malarone
3. How to take Malarone
4. Possible side effects
5. How to store Malarone
6. Contents of the pack and other information

1. What Malarone is and what it is used for

Malarone belongs to a group of medicines called *antimalarials*. It contains two active ingredients, atovaquone and proguanil hydrochloride.

What Malarone is used for

Malarone has two uses:

- to prevent malaria
- to treat malaria

Dosage instructions for each use are in Section 3, ‘How to take Malarone’.

Malaria is spread by the bite of an infected mosquito, which passes the malaria parasite (*Plasmodium falciparum*) into the bloodstream. Malarone prevents malaria by killing this parasite. For people who are already infected with malaria, Malarone also kills these parasites.

Protect yourself from catching malaria

People of any age can get malaria. It is a serious disease, but is preventable.

As well as taking Malarone, it is very important that you also take steps to avoid being bitten by mosquitoes.

- Use insect repellent on exposed areas of the skin
- Wear light coloured clothing that covers most of the body, especially after sunset as this is the time when mosquitoes are most active
- Sleep in a screened room or under a mosquito net impregnated with insecticide
- Close windows and doors at sunset, if they are not screened
- Consider using an insecticide (mats, spray, plug-ins) to clear a room of insects or to deter mosquitoes from entering the room.

→ If you need further advice, talk to your doctor or pharmacist.

It is still possible to get malaria after taking the necessary precautions.

Some types of malaria infection take a long time to cause symptoms, so the illness may not start until several days, weeks or even months after returning from abroad.

→ See a doctor immediately if you get symptoms such as high temperature, headache, shivering and tiredness after returning home.

2. What you need to know before you take Malarone

Do not take Malarone:

- if you are allergic to atovaquone, proguanil hydrochloride or any of the ingredients of this medicine listed in section 6.
- for preventing malaria, if you have severe kidney disease.

→ Tell your doctor if either of these apply to you.

Take special care with Malarone
Talk to your doctor or pharmacist before taking Malarone if:

- you have severe kidney disease
- your child is being treated for Malaria and weighs less than 11kg. There is another tablet strength to treat children who weigh less than 11kg (see section 3).

→ Tell your doctor or pharmacist if any of these applies to you.

Other medicines and Malarone

Tell your doctor or pharmacist if you are taking, have recently taken, or might take any other medicines, including medicines you've bought without a prescription.

Some medicines can affect the way Malarone works, or Malarone itself can strengthen or weaken the effectiveness of other medicines taken at the same time. These include:

- metoclopramide, used to treat nausea and vomiting
- the antibiotics, tetracycline, rifampicin and rifabutin
- efavirenz or certain highly active protease-inhibitors used to treat HIV
- warfarin and other medicines that stop blood clotting
- etoposide used to treat cancer.

→ Tell your doctor if you are taking any of these. Your doctor may decide that Malarone isn't suitable for you, or that you need extra check ups while you're taking it.

→ Remember to tell your doctor if you start taking any other medicines while you're taking Malarone.

Malarone with food and drink

Take Malarone with food or a milky drink, where possible. This will increase the amount of Malarone your body can absorb, and make your treatment more effective.

Pregnancy and breast-feeding

If you are pregnant, do not take Malarone unless your doctor recommends it.

→ Ask your doctor or pharmacist for advice before taking Malarone

Do not breast-feed while taking Malarone, as the ingredients of Malarone may pass into breast milk and may harm your baby.

Driving and using machines

If you feel dizzy, do not drive. Malarone makes some people feel dizzy. If this happens to you, do not drive, use machines or take part in activities where you may put yourself or others at risk.

Malarone contains sodium

This medicine contains less than 1mmol sodium (23mg) per tablet, that is to say essentially 'sodium free'.

3. How to take Malarone

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

Take Malarone with food or a milky drink, where possible.

It is best to take Malarone at the same time each day.

If you are sick (vomit)

For preventing malaria:

- if you are sick (vomit) within 1 hour of taking your Malarone tablet, take another dose straight away
- it is important to take the full course of Malarone. If you have to take extra tablets due to sickness, you may need another prescription.
- if you have been vomiting, it is especially important to use extra protection, such as repellents and bednets. Malarone may not be as effective, as the amount absorbed will be reduced.

For treating malaria:

- if you have vomiting and diarrhoea tell your doctor, you will need regular blood tests. Malarone will not be as effective, as the amount absorbed will be reduced. The tests will check whether the malaria parasite is being cleared from your blood.

To prevent malaria

The recommended usual dose for adults is 1 tablet once a day, taken as below.

Not recommended for preventing malaria in children, or in adults who weigh less than 40kgs.

Malarone paediatric tablets are recommended for preventing malaria in adults and children who weigh less than 40kgs.

To prevent malaria in adults:

- start taking Malarone 1 to 2 days before travelling to an area which has malaria
- continue taking it every day during your stay
- continue taking it for another 7 days after your return to a malaria-free area.

To treat malaria

The recommended dose for adults is 4 tablets once a day for 3 days.

For children the dose depends on their bodyweight:

- 11-20kg – 1 tablet once a day for 3 days
- 21-30kg – 2 tablets once a day for 3 days
- 31-40kg – 3 tablets once a day for 3 days
- over 40kg – dose as for adults.

Not recommended for treating malaria in children who weigh less than 11kgs.

For children who weigh less than 11kgs talk to your doctor. There may be a different type of Malarone tablet available in your country.

If you take more Malarone than you should

Contact a doctor or pharmacist for advice. If possible show them the Malarone pack.

If you forget to take Malarone

It is very important that you take the full course of Malarone.

If you forget to take a dose, don't worry. Just take your next dose as soon as you remember. Then continue your treatment as before.

Don't take extra tablets to make up for a missed dose. Just take your next dose at the usual time.

Don't stop taking Malarone without advice

Keep taking Malarone for 7 days after you return to a malaria-free area. Take the full course of Malarone for maximum protection. Stopping early puts you at risk of getting malaria, as it takes 7 days to ensure that any parasites that may be in your blood following a bite from an infected mosquito are killed.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Look out for the following severe reactions. They have occurred in a small number of people, but their exact frequency is unknown.

Severe allergic reactions - signs include:

- rash and itching
- sudden wheezing, tightness of the chest or throat, or difficulty breathing
- swollen eyelids, face, lips, tongue or other part of the body.

→ Contact a doctor immediately if you get any of these symptoms. Stop taking Malarone.

Severe skin reactions

- skin rash, which may blister and looks like small targets (central dark spots, surrounded by paler area with a dark ring around the edge) (erythema multiforme)
- severe widespread rash with blisters and peeling skin, particularly occurring around the mouth, nose, eyes and genitals (Stevens-Johnson syndrome).

→ If you notice any of these symptoms contact a doctor urgently.

Most of the other side effects reported have been mild and have not lasted very long.

Very common side effects

- These may affect more than 1 in 10 people:
- headache
 - feeling sick and being sick (nausea and vomiting)
 - stomach pain
 - diarrhoea.

Common side effects

- These may affect up to 1 in 10 people:
- dizziness
 - sleeping problems (insomnia)
 - strange dreams
 - depression
 - loss of appetite
 - fever
 - rash which may be itchy
 - cough

Common side effects, which may show up in your blood tests are:

- reduced numbers of red blood cells (anaemia) which can cause tiredness, headaches and shortness of breath
- reduced numbers of white blood cells (neutropenia) which may make you more likely to catch infections
- low levels of sodium in the blood (hyponatraemia)
- an increase in liver enzymes.

Uncommon side effects

These may affect up to 1 in 100 people:

- anxiety
- an unusual awareness of abnormal beating of the heart (palpitations)
- swelling and redness of the mouth
- hair loss
- itchy, bumpy rash (hives).

Uncommon side effects that may show up in your blood tests:

- an increase in amylase (an enzyme produced in the pancreas).

Rare side effects

These may affect up to 1 in 1,000 people:

- seeing or hearing things that are not there (hallucinations)

Other side effects

Other side effects have occurred in a small number of people but their exact frequency is unknown.

- Inflammation of the liver (hepatitis)
 - blockage of the bile ducts (cholestatis)
 - increase in heart rate (tachycardia)
 - inflammation of the blood vessels (vasculitis) which may be visible as red or purple raised spots on the skin but can affect other parts of the body
 - fits (seizures)
 - panic attacks, crying
 - nightmares
 - severe mental health problem in which the person loses contact with reality and is unable to think and judge clearly
 - indigestion
 - mouth ulcers
 - blisters
 - peeling skin
 - increased sensitivity of the skin to sunlight.
- Other side effects that may show up in your blood tests:
- A decrease in all types of blood cells (pancytopenia).

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store.

By reporting side effects, you can help provide more information on the safety of this medicine.

5. How to store Malarone

Keep out of the sight and reach of children.

Malarone does not require any special storage conditions.

Do not take the tablets after the expiry date which is stated on the carton and blister labels after 'Exp'. The expiry date refers to the last day of that month.

If the tablet becomes discoloured or shows any signs of deterioration, seek the advice of your pharmacist.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

6. Contents of the pack and other information

What Malarone contains

The active ingredients in the Malarone are atovaquone and proguanil hydrochloride.

Each film-coated tablet contains 250mg atovaquone and 100mg proguanil hydrochloride.

The other ingredients are:
Tablet core: poloxamer 188, microcrystalline cellulose (E460), hydroxypropyl cellulose (E463), povidone K30 (E1201), sodium starch glycolate (Type A), magnesium stearate (E470B).

Tablet coating: hypromellose (E464), titanium dioxide (E171), red iron oxide (E172), macrogol 400 and polyethylene glycol 8000 (see section 2).
→ Tell your doctor without taking Malarone if you might be allergic to any of these ingredients.

What Malarone looks like and contents of the pack

Malarone is a pink, round, biconvex, film-coated tablet engraved 'GX CM3' on one side and plain on the reverse. Malarone is available in blister packs of 12 tablets.

Manufactured by: Glaxo Wellcome S.A., Avenida de Extremadura, 3, 09400 Aranda de Duero, Burgos, Spain.

Procured from within the EU and repackaged by the Product Licence holder: B&S Healthcare, Unit 4, Bradfield Road, Ruislip, Middlesex, HA4 0NU, UK.

Malarone® 250mg/100mg film-coated tablets; PL 18799/3141

POM

Leaflet date: 28.12.2022

Malarone is the registered trademark of Glaxo Group Limited.

Blind or partially sighted?
Is this leaflet hard to see or read?
Call 0208 515 3763 to obtain the leaflet in a format suitable for you.