

Package leaflet: Information for the user

**AMOXICILLIN 125 mg/5 ml
ORAL SUSPENSION SUGAR FREE BP
AMOXICILLIN 250 mg/5 ml
ORAL SUSPENSION SUGAR FREE BP**

Amoxicillin (as amoxicillin trihydrate)

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you (or for your child) only. Do not pass it on to others. It may harm them even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Amoxicillin Oral Suspension Sugar Free is and what it is used for
2. What you need to know before you take Amoxicillin Oral Suspension Sugar Free
3. How to take Amoxicillin Oral Suspension Sugar Free
4. Possible side effects
5. How to store Amoxicillin Oral Suspension Sugar Free
6. Contents of the pack and other information

1. WHAT AMOXICILLIN ORAL SUSPENSION SUGAR FREE IS AND WHAT IT IS USED FOR

What Amoxicillin Oral Suspension Sugar Free is

Amoxicillin Oral Suspension Sugar Free is an antibiotic. The active ingredient is amoxicillin. This belongs to a group of medicines called 'penicillins'.

What Amoxicillin Oral Suspension Sugar Free is used for

Amoxicillin Oral Suspension Sugar Free is used to treat infections caused by bacteria in different parts of the body. Amoxicillin Oral Suspension Sugar Free may also be used in combination with other medicines to treat stomach ulcers.

2. WHAT YOU NEED TO KNOW BEFORE YOU TAKE AMOXICILLIN ORAL SUSPENSION SUGAR FREE

Do not take Amoxicillin Oral Suspension Sugar Free:

- if you are allergic to amoxicillin, penicillin or any of the other ingredients of this medicine (listed in section 6).
- if you have ever had an allergic reaction to any antibiotic. This can include a skin rash or swelling of the face or throat.

Do not take Amoxicillin Oral Suspension Sugar Free if any of the above apply. If you are not sure, talk to your doctor or pharmacist before taking Amoxicillin Oral Suspension Sugar Free.

Warnings and Precautions

Talk to your doctor or pharmacist before taking Amoxicillin Oral Suspension Sugar Free if you:

- have kidney problems,
- have glandular fever (fever, sore throat, swollen glands and extreme tiredness)
- are not urinating regularly

If you are not sure if any of the above apply to you, talk to your doctor or pharmacist before taking Amoxicillin Oral Suspension Sugar Free.

Conditions you need to look out for

Amoxicillin can make some existing conditions worse, or cause serious side effects, such as severe allergic reactions, chest pain, repetitive vomiting within 1 to 4 hours of Amoxicillin administration, serious skin reactions or severe diarrhoea (pseudomembranous colitis). You must look out for certain symptoms while you are receiving Amoxicillin to help reduce the risk of any problems (see section 4).

Blood and urine tests

If you are having:

- Urine tests (glucose) or blood tests for liver function
- Oestriol tests (used during pregnancy to check the baby is developing normally)

Tell your doctor or pharmacist that you are taking Amoxicillin Oral Suspension Sugar Free. This is because Amoxicillin Oral Suspension Sugar Free can affect the results of these tests.

Other medicines and Amoxicillin Oral Suspension Sugar Free

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

- If you are taking allopurinol (used for gout) with Amoxicillin Oral Suspension Sugar Free, it may be more likely that you will have an allergic skin reaction.
- If you are taking probenecid (used for gout), concomitant use of probenecid may reduce the excretion of this medicine and is not recommended.
- If you are taking medicines to help stop blood clots (such as warfarin), you may need extra blood tests.
- If you are taking other antibiotics (such as tetracycline) Amoxicillin Oral Suspension Sugar Free may be less effective.
- If you are taking methotrexate (used for the treatment of cancer and severe psoriasis) Amoxicillin Oral Suspension Sugar Free may reduce the excretion of methotrexate causing a potential increase in side effects.
- Oral typhoid vaccine (may not work if taken with Amoxicillin Oral Suspension Sugar Free).

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

Driving and using machines

Amoxicillin Oral Suspension Sugar Free can have side effects and the symptoms (such as allergic reactions, dizziness and convulsions) may make you unfit to drive. Do not drive or operate machinery unless you are feeling well.

Amoxicillin 125 mg/5 ml and 250 mg/5 ml Oral Suspension Sugar Free BP contains sorbitol, sodium and sodium benzoate

- This medicine contains 800 mg sorbitol in each 5ml dose.
- This medicine contains less than 1 mmol sodium (23 mg) per 5ml, that is to say essentially 'sodium-free'.
- Sodium Benzoate may increase jaundice (yellowing of the skin and eyes) in newborn babies (up to 4 weeks old).

3. HOW TO TAKE AMOXICILLIN ORAL SUSPENSION SUGAR FREE

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

- Shake bottle well before each dose
- Space the doses evenly during the day, at least 4 hours apart.

The usual dose is:

Children weighing less than 40kg

All doses are worked out depending on your child's body weight in kilograms.

- Your doctor will advise you how much Amoxicillin Oral Suspension Sugar Free you should give to your baby or child.
- The usual dose is 40mg to 90mg for each kilogram of body weight a day given in two or three divided doses.
- The maximum recommended dose is 100 mg for each kilogram of body weight a day.

Adults, elderly patients and children weighing 40kg or more

This suspension is not usually prescribed for adults and children weighing more than 40 kg. Ask your doctor or pharmacist for advice.

Kidney problems

If you have kidney problems the dose might be lower than the usual dose.

If you take more Amoxicillin Oral Suspension Sugar Free than you should

If you have taken too much Amoxicillin Oral Suspension Sugar Free, signs might be an upset stomach (feeling sick, being sick or diarrhoea) or crystals in the urine, which may be seen as cloudy urine, or problems urinating. Talk to your doctor as soon as possible. Take the medicine to show the doctor.

If you forget to take Amoxicillin Oral Suspension Sugar Free

- If you forget to take a dose, take it as soon as you remember.
- Do not take the next dose too soon, wait about 4 hours before taking the next dose.
- Do not take a double dose to make up for a forgotten dose.

How long should you take Amoxicillin Oral Suspension Sugar Free for?

- Keep taking Amoxicillin Oral Suspension Sugar Free for as long as your doctor has told you to, even if you feel better. You need every dose to help fight the infection. If some bacteria survive they can cause the infection to come back.
- Once you finish treatment, if you still feel unwell you should go back to see the doctor

Thrush (a yeast infection of moist areas of the body which can cause soreness, itching and white discharge) may develop if Amoxicillin Oral Suspension Sugar Free is used for a long time. If this occurs tell your doctor.

If you take Amoxicillin Oral Suspension Sugar Free for a long time, your doctor may perform additional tests to check your kidneys, liver and blood are working normally.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. POSSIBLE SIDE EFFECTS

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Stop taking Amoxicillin Oral Suspension Sugar Free and see a doctor straight away, if**you notice any of the following serious side effects – you may need urgent medical treatment:**

The following reactions have been reported but the frequency is unknown (the frequency cannot be estimated from the available data):

- chest pain in the context of allergic reactions, which may be a symptom of allergy triggered cardiac infarction (Kounis syndrome)
- Drug-induced enterocolitis syndrome (DIES): DIES has been reported mainly in children receiving amoxicillin. It is a certain kind of allergic reaction with the leading symptom of repetitive vomiting (1-4 hours after drug intake). Further symptoms could comprise abdominal pain, lethargy, diarrhoea, and low blood pressure.
- Crystals in urine leading to acute renal injury
- Rash with blisters arranged in a circle with central crusting or like a string of pearls (linear IgA disease)
- Inflammation of the membranes that surround the brain and spinal cord (aseptic meningitis)

The following are very rare (may affect up to 1 in 10,000 people)

- allergic reactions, the signs may include: skin itching or rash, swelling of the face, lips, tongue, body or breathing difficulties. These can be serious and occasionally deaths have occurred
- rash or pinpoint flat red round spots under the skin surface or bruising of the skin. This is due to inflammation of blood vessel walls due to an allergic reaction. It can be associated with joint pain (arthritis) and kidney problems
- a delayed allergic reaction can occur usually 7 to 12 days after having Amoxicillin Oral Suspension Sugar Free, some signs include: rashes, fever, joint pains and enlargement of the lymph nodes especially under the arms
- a skin reaction known as 'erythema multiforme' where you may develop: itchy reddish purple patches on the skin especially on the palms of the hands or soles of the feet, 'hive-like' raised swollen areas on the skin, tender areas on the surfaces of the mouth, eyes and genitals. You may have a fever and be very tired
- other severe skin reactions can include: changes in skin colour, bumps under the skin, blistering, pustules, peeling, redness, pain, itching, scaling. These may be associated with fever, headaches and body aches
- flu-like symptoms with a rash, fever, swollen glands, and abnormal blood test results (including increased white blood cells (eosinophilia) and liver enzymes) (Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS)).
- a red rash commonly seen on both sides of buttocks, upper inner thighs, armpits, neck [Symmetrical Drug-Related Intertriginous and Flexural Exanthema (SDRIFE)]
- fever, chills, a sore throat or other signs of an infection, or if you bruise easily. These may be signs of a problem with your blood cells
- the *Jarisch-Herxheimer reaction* which occurs during treatment with Amoxicillin Oral Suspension Sugar Free for Lyme disease and causes fever, chills, headache, muscle pain and skin rash.
- inflammation of the large bowel (colon) with diarrhoea (sometimes containing blood), pain and fever
- serious liver side effects may occur. They are mainly associated with people having treatment over a long period, males and the elderly. You must tell your doctor urgently if you get:
 - o severe diarrhoea with bleeding
 - o blisters, redness or bruising of the skin
 - o darker urine or paler stools
 - o yellowing of the skin or the whites of the eyes (jaundice). See also anaemia below which might result in jaundice

These can happen when having the medicine for or up to several weeks after.

If any of the above happens stop taking the medicine and see your doctor straight away.**Sometimes you may get less severe skin reactions such as:**

- a mildly itchy rash (round, pink-red patches), 'hive-like' swollen areas on forearms, legs, palms, hands or feet. This is uncommon (may affect up to 1 in 100 people).

If you have any of these talk to your doctor as Amoxicillin Oral Suspension Sugar Free will need to be stopped.

The other possible side effects are:

- **Common** (may affect up to 1 in 10 people)
 - skin rash
 - feeling sick (nausea)
 - diarrhoea.

- **Uncommon** (may affect up to 1 in 100 people)
 - being sick (vomiting).

- **Very rare** (may affect up to 1 in 10,000 people)
 - thrush (a yeast infection of the vagina, mouth or skin folds), you can get treatment for thrush from your doctor or pharmacist.

- kidney problems
- fits (convulsions), seen in patients on high doses or with kidney problems
- dizziness
- hyperactivity
- teeth may appear stained, usually returning to normal with brushing (this has been reported in children).
- the tongue may change to yellow, brown or black and it may have a hairy appearance
- an excessive breakdown of red blood cells causing a type of anaemia. Signs include: tiredness, headaches, shortness of breath, dizziness, looking pale and yellowing of the skin and the whites of the eyes

- low number of white blood cells
- low number of cells involved with blood clotting
- the blood may take longer to clot than it normally would. You may notice this if you have a nosebleed or cut yourself.
- inflammation of the protective membrane surrounding the brain (aseptic meningitis).

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the yellow card scheme at www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store.

By reporting side effects you can help provide more information on the safety of this medicine.

5. HOW TO STORE AMOXICILLIN ORAL SUSPENSION SUGAR FREE

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the label. The expiry date refers to the last day of that month.

Dry powder

Do not store above 25°C.

Store in original package. Keep the bottle tightly closed.

Reconstituted suspension

Once made up, store up to 14 days at 2°C-8°C in a refrigerator.

Do not use this medicine if there are visible signs of deterioration.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. CONTENTS OF THE PACK AND OTHER INFORMATION

What Amoxicillin Oral Suspension Sugar Free BP contains

- The active substance in each suspension is 125 mg or 250 mg of amoxicillin as amoxicillin trihydrate.
- The other ingredients are: Sodium Benzoate (E211), Disodium Edetate, Sodium Citrate, Orange Bramble Flavour, Quinoline Yellow (E104), Citric Acid Monohydrate, Colloidal Anhydrous Silica, Xanthan Gum, Sorbitol (E420) and Saccharin Sodium.

What Amoxicillin Oral Suspension Sugar Free BP looks like and contents of the pack

The suspensions are pale yellow with a characteristic odour. Amoxicillin Oral Suspensions Sugar Free BP are available in bottles of 100ml. Not all pack sizes may be marketed.

Marketing Authorisation Holder

Kent Pharma UK Limited, 2nd Floor, Connect 38, 1 Dover Place, Ashford, Kent, England, TN23 1FB.

Manufacturer

Athlone Laboratories Limited, Ballymurray, Co. Roscommon, Ireland.
Kent Pharmaceuticals Limited, Repton Road, Measham, DE12 7DT, U.K.

PL 51463/0174

PL 51463/0175

This leaflet was last revised in January 2026.

General advice regarding the use of antibiotics

Antibiotics are used to treat infections caused by bacteria. They have no effect against infections caused by viruses.

Sometimes an infection caused by bacteria does not respond to a course of an antibiotic. One of the commonest reasons for this to occur is because the bacteria causing the infection are resistant to the antibiotic that is being taken. This means that they can survive and even multiply despite the antibiotic. Bacteria can become resistant to antibiotics for many reasons. Using antibiotics carefully can help to reduce the chance of bacteria becoming resistant to them.

When your doctor prescribes a course of an antibiotic it is intended to treat only your current illness. Paying attention to the following advice will help prevent the emergence of resistant bacteria that could stop the antibiotic working.

1. It is very important that you take the antibiotic at the right dose, at the right times and for the right number of days. Read the instructions on the label and if you do not understand anything ask your doctor or pharmacist to explain.
2. You should not take an antibiotic unless it has been prescribed specifically for you and you should use it only to treat the infection for which it was prescribed.
3. You should not take antibiotics that have been prescribed for other people even if they had an infection that was similar to yours.
4. You should not give antibiotics that were prescribed for you to other people.
5. If you have any antibiotic left over when you have taken the course as directed by your doctor you should take the remainder to a pharmacy for appropriate disposal.

Customer: Kent **Originated by:** DOB (AWG)

Product: Amoxicillin JNT Oral Suspension SF

Dimensions: 148mm x 450mm

Item Code: CP.AMO.JNT.SuSF.100.Sp.AT.V11P1 (P513/15)


Proof No and Date: 1 15/01/2026

Revision: Brand Review

ARTWORK FOR SUBMISSION

Supersedes: CP.AMO.JNT.SuSF.100.Sp.AT.V10P1 P513/14

Colours



Pantone Blue
280