


Base Label



Proof: 2		Artwork Information Panel			Almirall		
<b>Description:</b> PIL Robaxin 750mg tabs GB Mock up		<b>Printing Colours</b>		<b>Non Printing Colours</b>		<b>Inline control code</b>	
<b>Regulatory text:</b> N/A		P Black	30% P 2718	Die cut	Braille		
<b>Item Number:</b> Mock up		<b>Material size:</b> 150 x 560 mm		Colour reference in PMS Bridge Equate with CMYK			
<b>Designer:</b> Thinkinpressi		<b>Date:</b> 03.07.2017					
<b>Fonts:</b> Fruitger LT Com light, Fruitger LT Com light italic, Fruitger LT Com Bold, Fruitger LT Com Black, Fruitger LT Com Black Condensed							
<b>Manufacturer's Technical check:</b> Manufacturer confirms the printability and technical compliance of this packaging material for correct use during packaging process. <input type="checkbox"/>							
						Date and Signature	



Front

1.

2.

3.

4.

<p><b>Children:</b> These tablets should not be given to children</p> <p><b>Liver impairment:</b> you may need a longer interval between taking the tablets if you have liver disease. You should always follow your doctor's instructions carefully.</p> <p>These tablets are for short-term treatment. Take them for as long as your doctor has recommended, or until you no longer need them to ease the pain. Return any unused tablets to your pharmacist.</p> <p><b>If you take more Robaxin-750 than you should</b> <b>Contact your doctor immediately,</b> go or take the person to the Accident and Emergency Department of your local hospital. Always take the bottle of Robaxin-750 with you, even if it is empty.</p> <p><b>If you forget to take Robaxin-750</b> Take your normal dose when you remember. Always leave at least 4 hours before taking any more tablets. DO NOT TAKE A DOUBLE DOSE OF TABLETS TO MAKE UP FOR A MISSED DOSE.</p>	<p><b>4. Possible side effects</b></p> <p>Like all medicines, this medicine can cause side effects, although not everybody gets them.</p> <p>If you notice any of the following rare symptoms, you should <b>STOP</b> taking your tablets and tell your doctor immediately:</p> <ul style="list-style-type: none"> <li>• confusion, convulsions, fits, and rash or swelling of the face or neck and shortness of breath which may be a sign of an allergy.</li> </ul> <p>If you notice any of these symptoms, you should contact your doctor:</p> <ul style="list-style-type: none"> <li>• yellowing of the skin and/or whites of the eyes.</li> </ul> <p>The levels of your white blood cells may also be reduced which can lead to recurrent:</p> <ul style="list-style-type: none"> <li>• infections</li> <li>• mouth ulcers.</li> </ul> <p>If any of the following side effects are troublesome, you should tell your doctor:</p> <ul style="list-style-type: none"> <li>• drowsiness, difficulty in sleeping,</li> <li>• restlessness, anxiety or worry,</li> <li>• dizziness,</li> </ul>	<ul style="list-style-type: none"> <li>• memory loss,</li> <li>• blurred or double vision, rapid or unusual eye movements,</li> <li>• reddening of the eyes together with nasal stuffiness,</li> <li>• feeling sick or vomiting,</li> <li>• heartburn,</li> <li>• taste disturbance,</li> <li>• tremor or shakiness, mild muscle incoordination,</li> <li>• headache,</li> <li>• fever,</li> <li>• itching,</li> <li>• a slow heartbeat,</li> <li>• flushing,</li> <li>• fainting or feeling faint,</li> <li>• low blood pressure,</li> </ul> <p><b>Reporting of side effects</b></p> <p>If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme, Website: <a href="http://www.mhra.gov.uk/yellowcard">http://www.mhra.gov.uk/yellowcard</a>.</p>	<p>By reporting side effects you can also provide more information on the safety of this medicine.</p> <p><b>5. How to store Robaxin-750</b> <b>Keep this medicine out of the sight and reach of children.</b></p> <p>Do not take this medicine after the expiry date which is stated on the label of the container after EXP.</p> <p>Take any unused medicine to your pharmacist to be disposed of.</p> <p><b>6. Contents of the pack and other information</b> <b>What Robaxin-750 contains</b></p> <ul style="list-style-type: none"> <li>- The active substance is methocarbamol (750mg)</li> <li>- The other ingredients are: maize starch, alginate acid, povidone, sodium lauryl sulphate, magnesium stearate, talc, and gelatine.</li> </ul> <p>The tablet coating contains hypromellose (E464), microcrystalline cellulose (E460), polyoxyl 8 stearate, propylene glycol and titanium dioxide (E171).</p>	<p><b>What Robaxin-750 looks like and contents of the pack</b></p> <p>Robaxin-750 tablets are white, capsule-shaped tablets and are supplied in bottles of 100 tablets.</p> <p><b>Marketing Authorisation Holder and Manufacturer</b> <b>Marketing Authorisation Holder:</b> Almirall, S.A. Ronda General Mitre, 151 08022 Barcelona - Spain</p> <p><b>Manufacturer**:</b> Hamol Limited, Nottingham, NG90 2DB or Industrias Farmacéuticas Almirall, S.A., Ctra. Nacional II, Km. 593, 08740 Sant Andreu de la Barca, Barcelona, Spain [**only the manufacturer releasing the batch will appear on the printed leaflet]</p> <p><b>This leaflet was last revised in: July 2017</b></p>
--	--	---	--	---

5.

6.

7.

8.

9.