

# SUMMARY OF PRODUCT CHARACTERISTICS

## Product Summary

### 1. Name of the Medicinal Product

Cutivate 0.05% w/w Cream

### 2. Qualitative and Quantitative Composition

Fluticasone Propionate (micronised) HSE 0.05% w/w.

Excipients with known effect:

Cetostearyl Alcohol

Propylene Glycol

Imidurea

For the full list of excipients, see section 6.1.

### 3. PHARMACEUTICAL FORM

Cream

## 4 CLINICAL PARTICULARS

### 4.1 Therapeutic indications

#### TREATMENT OF INFLAMMATORY DERMATOSES

##### Adults:

Fluticasone propionate cream is a potent topical corticosteroid indicated for the relief of the inflammatory and pruritic manifestations of corticosteroid-responsive dermatoses; these include the following:

- Atopic dermatitis
- Nummular dermatitis (discoid eczemas)

- Prurigo nodularis
- Psoriasis (excluding widespread plaque psoriasis)
- Lichen simplex chronicus (neurodermatitis) and lichen planus
- Seborrhoeic dermatitis
- Irritant or allergic contact dermatitis
- Discoid lupus erythematosus
- An adjunct to systemic steroid therapy in generalised erythroderma
- Insect bite reactions
- Miliaria (prickly heat)

#### **Children:**

For children and infants aged three months and over who are unresponsive to lower potency corticosteroids, Cutivate cream is indicated for the relief of the inflammatory and pruritic manifestations of atopic dermatitis under the supervision of a specialist. Expert opinion should be sought prior to the use of Cutivate cream in other corticosteroid-responsive dermatoses in children.

## **4.2 Posology and Method of Administration**

Route of administration: Cutaneous

Adults, elderly, children and infants aged 3 months and over

Cream

Crems are especially appropriate for moist or weeping surfaces.

### **TREATMENT OF INFLAMMATORY DERMATOSES**

Apply thinly and gently rub in using only enough to cover the entire affected area once or twice a day for up to 4 weeks until improvement occurs, then reduce the frequency of application or change the treatment to a less potent preparation. Allow adequate time for absorption after each application before applying an emollient.

Therapy with topical corticosteroids should be gradually discontinued once control is achieved and an emollient continued as maintenance therapy.

Rebound of pre-existing dermatoses can occur with abrupt discontinuation of topical steroids especially with potent preparations.

Duration of treatment for adults and elderly

If the condition worsens or does not improve within four weeks, treatment and diagnosis should be re-evaluated.

Children over 3 months

Children are more likely to develop local and systemic side effects of topical corticosteroids and, in general, require shorter courses and less potent agents than adults.

Care should be taken when using fluticasone propionate to ensure the amount applied is the minimum that provides therapeutic benefit.

Duration of treatment for children and Infants

When Cutivate is used in the treatment of children, if there is no improvement within 7 – 14 days, treatment should be withdrawn and the child re-evaluated. Once the condition has been controlled (usually within 7-14 days), frequency of application should be reduced to the lowest effective dose for the shortest possible time. Continuous daily treatment for longer than 4 weeks is not recommended

Elderly

Clinical studies have not identified differences in responses between the elderly and younger patients. The greater frequency of decreased hepatic or renal function in the elderly may delay elimination if systemic absorption occurs. Therefore the minimum quantity should be used for the shortest duration to achieve the desired clinical benefit.

Renal / Hepatic Impairment

In case of systemic absorption (when application is over a large surface area for a prolonged period) metabolism and elimination may be delayed therefore increasing the risk of systemic toxicity. Therefore the minimum quantity should be used for the shortest duration to achieve the desired clinical benefit.

### **4.3 Contraindications**

Hypersensitivity to the active substance or any of the excipients listed in section 6.1.

The following conditions should not be treated with fluticasone propionate:

- Untreated cutaneous infections
- Rosacea
- Acne vulgaris

- Perioral dermatitis
- Perianal and genital pruritus
- Pruritus without inflammation
- Dermatoses in infants under three months of age, including dermatitis and nappy rash.

#### 4.4 Special warnings and precautions for use

Fluticasone propionate should be used with caution in patients with a history of local hypersensitivity to other corticosteroids. Local hypersensitivity reactions (*see section 4.8*) may resemble symptoms of the condition under treatment.

Manifestations of hypercortisolism (Cushing's Syndrome) and reversible hypothalamic-pituitary-adrenal (HPA) axis suppression, leading to glucocorticosteroid insufficiency, can occur in some individuals as a result of increased systemic absorption of topical steroids. If either of the above are observed, withdraw the drug gradually by reducing the frequency of application, or by substituting a less potent corticosteroid. Abrupt withdrawal of treatment may result in glucocorticosteroid insufficiency (*see section 4.8*).

Risk factors for increased systemic effects are:

- Potency and formulation of topical steroid
- Duration of exposure
- Application to a large surface area
- Use on occluded areas of skin (e.g. on intertriginous areas or under occlusive dressings (in infants the nappy may act as an occlusive dressing))
- Increasing hydration of the stratum corneum
- Use on thin skin areas such as the face
- Use on broken skin or other conditions where the skin barrier may be impaired
- In comparison with adults, children and infants may absorb proportionally larger amounts of topical corticosteroids and thus be more susceptible to systemic adverse effects. This is because children have an immature skin barrier and a greater surface area to body weight ratio compared with adults.

##### **Children**

In infants and children under 12 years of age, long-term continuous topical corticosteroid therapy should be avoided where possible, as adrenal suppression is more likely to occur.

##### **Use in psoriasis**

Topical steroids should be used with caution in psoriasis as rebound relapses, development of tolerance, risk of generalised pustular psoriasis and development of local or systemic toxicity due to impaired barrier function of the skin have been reported in some cases. If used in psoriasis, careful patient supervision is important.

##### **Application to the face**

Prolonged application to the face is undesirable as this area is more susceptible to atrophic changes.

#### **Application to the eyelids**

If applied to the eyelids, care is needed to ensure that the preparation does not enter the eye, as cataract and glaucoma might result from repeated exposure.

#### **Visual disturbance**

Visual disturbance may be reported with systemic and topical corticosteroid use. If a patient presents with symptoms such as blurred vision or other visual disturbances, the patient should be considered for referral to an ophthalmologist for evaluation of possible causes which may include cataract, glaucoma or rare diseases such as central serous chorioretinopathy (CSCR) which have been reported after use of systemic and topical corticosteroids.

#### **Concomitant infection**

Appropriate antimicrobial therapy should be used whenever treating inflammatory lesions which have become infected. Any spread of infection requires withdrawal of topical corticosteroid therapy and administration of appropriate antimicrobial therapy.

#### **Infection risk with occlusion**

Bacterial infection is encouraged by the warm, moist conditions within skin folds or caused by occlusive dressings. When using occlusive dressings, the skin should be cleansed before a fresh dressing is applied.

#### **Chronic leg ulcers**

Topical corticosteroids are sometimes used to treat the dermatitis around chronic leg ulcers. However, this use may be associated with a higher occurrence of local hypersensitivity reactions and an increased risk of local infection.

Overt suppression of the HPA-axis (morning plasma cortisol less than 5 micrograms/dL) is very unlikely to result from therapeutic use of fluticasone propionate cream or ointment unless treating more than 50% of an adult's body surface and applying more than 20 g per day.

Fluticasone propionate cream contains the excipient imidurea which releases traces of formaldehyde as a breakdown product. Formaldehyde may cause allergic sensitisation or irritation upon contact with the skin.

Fluticasone propionate cream contains the excipient cetostearyl alcohol which may cause local skin reactions (e.g. local dermatitis).

Fluticasone propionate cream contains paraffin. Instruct patients not to smoke or go near naked flames due to the risk of severe burns. Fabric (clothing, bedding, dressings etc) that has been in contact with this product burns more easily and is a serious fire hazard. Washing clothing and bedding may reduce product build-up but not totally remove it.

Fluticasone propionate cream contains 100 mg of the excipient propylene glycol in each gram which may cause local skin irritation.

#### **Topical steroid withdrawal syndrome**

Long term use of topical steroids can result in the development of rebound flares after stopping treatment (topical steroid withdrawal syndrome). A severe form of rebound flare can develop which takes the form of a dermatitis with intense redness, stinging

and burning that can spread beyond the initial treatment area. It is more likely to occur when delicate skin sites such as the face and flexures are treated. Should there be a reoccurrence of the condition within days to weeks after successful treatment a withdrawal reaction should be suspected. Reapplication should be with caution and specialist advice is recommended in these cases or other treatment options should be considered.

The label will state strong steroid.

#### **4.5 Interaction with other medicinal products and other forms of interaction**

Co-administered drugs that can inhibit CYP3A4 (e.g. ritonavir, itraconazole) have been shown to inhibit the metabolism of corticosteroids leading to increased systemic exposure. The extent to which this interaction is clinically relevant depends on the dose and route of administration of the corticosteroids and the potency of the CYP3A4 inhibitor.

#### **4.6 Fertility, pregnancy and lactation**

##### Pregnancy

There are limited data from the use of fluticasone propionate in pregnant women.

Topical administration of corticosteroids to pregnant animals can cause abnormalities of foetal development (*see section 5.3*).

The relevance of this finding to humans has not been established; however, administration of fluticasone propionate during pregnancy should only be considered if the expected benefit to the mother is greater than any possible risk to the foetus. The minimum quantity should be used for the minimum duration.

##### Breast-feeding

The safe use of topical corticosteroids during lactation has not been established.

It is not known whether the topical administration of corticosteroids could result in sufficient systemic absorption to produce detectable amounts in breast milk.

When measurable plasma levels were obtained in lactating laboratory rats following subcutaneous administration there was evidence of fluticasone propionate in the milk.

Administration of fluticasone propionate during lactation should only be considered if the expected benefit to the mother outweighs the risk to the infant.

If used during lactation, fluticasone propionate should not be applied to the breasts to avoid accidental ingestion by the infant.

### Fertility

There are no data in humans to evaluate the effect of topical corticosteroids on fertility (*see section 5.3*).

## **4.7 Effect on Ability to Drive and Use Machines**

There have been no studies to investigate the effect of fluticasone propionate on driving performance or the ability to operate machinery. A detrimental effect on such activities would not be anticipated from the adverse reaction profile of topical fluticasone propionate.

## **4.8 Undesirable effects**

Adverse drug reactions (ADRs) are listed below by MedDRA system organ class and by frequency. Frequencies are defined as: very common ( $\geq 1/10$ ), common ( $\geq 1/100$  to  $< 1/10$ ), uncommon ( $\geq 1/1,000$  to  $< 1/100$ ), rare ( $\geq 1/10,000$  to  $< 1/1,000$ ) and very rare ( $< 1/10,000$ ), including isolated reports.

### **Post-Marketing Data**

#### **Infections and infestations**

Very rare: opportunistic infection

#### **Immune system disorders**

Very rare: Hypersensitivity.

#### **Endocrine disorders**

Very rare:

Hypothalamic-pituitary adrenal (HPA) axis suppression:

- Increased weight / obesity
- Delayed weight gain/growth retardation in children
- Cushingoid features (e.g. moon face, central obesity)
- Decreased endogenous cortisol levels
- Hyperglycaemia/glucosuria
- Hypertension
- Osteoporosis
- Cataract

- Glaucoma

### **Skin and subcutaneous tissue disorders**

Common: Pruritus.  
Uncommon: Local skin burning.

Very rare: Skin thinning, atrophy, striae, telangiectasias, pigmentation changes hypertrichosis, allergic contact dermatitis, exacerbation of underlying symptoms, pustular psoriasis, erythema, rash, urticaria

Not known Withdrawal reactions - redness of the skin which may extend to areas beyond the initial affected area, burning or stinging sensation, itch, skin peeling, oozing pustules. (see section 4.4)

### **Eye disorders**

Not known: Vision, blurred (see also section 4.4)

### **Reporting of suspected adverse reactions**

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store.

## **4.9 Overdose**

### Symptoms and Signs

Topically applied fluticasone propionate may be absorbed in sufficient amounts to produce systemic effects. Acute overdosage is very unlikely to occur, however, in the case of chronic overdosage or misuse, the features of hypercortisolism may appear. (*See section 4.8*)

### Treatment

In the event of overdose, fluticasone propionate should be withdrawn gradually by reducing the frequency of application, or by substituting a less potent corticosteroid because of the risk of glucocorticosteroid insufficiency. Further management should be as clinically indicated or as recommended by the national poisons centre, where available.

## **5 PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

**ATC Code: D07AC Corticosteroid, potent (Group III)**



Topical corticosteroids, have anti-inflammatory, antipruritic, and vasoconstrictive properties.

They act as anti-inflammatory agents via multiple mechanisms to inhibit late phase allergic reactions including decreasing the density of mast cells, decreasing chemotaxis and activation of eosinophils, decreasing cytokine production by lymphocytes, monocytes, mast cells and eosinophils, and inhibiting the metabolism of arachidonic acid.

Fluticasone propionate is a glucocorticoid with high topical anti-inflammatory potency but low HPA-axis suppressive activity after dermal administration. It therefore has a therapeutic index which is greater than most of the commonly available steroids.

It shows high systemic glucocorticoid potency after subcutaneous administration but very weak oral activity, probably due to metabolic inactivation. *In vitro* studies show a strong affinity for and agonist activity at, human glucocorticoid receptors.

Fluticasone propionate has no unexpected hormonal effects, and no overt, marked effects upon the central and peripheral nervous systems, the gastrointestinal system, or the cardiovascular or respiratory systems.

## **5.2. Pharmacokinetic Properties**

Pharmacokinetic data for the rat and dog indicate rapid elimination and extensive metabolic clearance. Bioavailability is very low after topical or oral administration, due to limited absorption through the skin or from the gastrointestinal tract, and because of extensive first-pass metabolism. Distribution studies have shown that only minute traces of orally administered compound reach the systemic circulation, and that any systemically-available radiolabel is rapidly eliminated in the bile and excreted in the faeces.

Fluticasone propionate does not persist in any tissue, and does not bind to melanin. The major route of metabolism is hydrolysis of the S-fluoromethyl carbothioate group, to yield a carboxylic acid (GR36264), which has very weak glucocorticoid or anti-inflammatory activity. In all test animal species, the route of excretion of radioactivity is independent of the route of administration of radiolabelled fluticasone propionate. Excretion is predominantly faecal and is essentially complete within 48 hours.

In man too, metabolic clearance is extensive, and elimination is consequently rapid. Thus drug entering the systemic circulation via the skin, will be rapidly inactivated. Oral bioavailability approaches zero, due to poor absorption and extensive first-pass metabolism. Therefore systemic exposure to any ingestion of the topical formulation will be low.

## **5.3. Pre-clinical Safety Data**

Reproductive studies suggest that administration of corticosteroids to pregnant animals can result in abnormalities of foetal development including cleft palate/lip. However, in humans, there is no convincing evidence of congenital abnormalities, such as cleft palate or lip.

Studies of safety pharmacology, repeated dose toxicity, genotoxicity, carcinogenic potential, fertility and general reproductive performance revealed no special hazard for humans, other than that anticipated for a potent steroid.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of Excipients**

Liquid Paraffin  
Cetostearyl Alcohol  
Isopropyl Myristate  
Cetomacrogol 1000  
Propylene Glycol  
Imidurea  
Sodium Phosphate  
Citric Acid Monohydrate  
Purified Water

### **6.2 Incompatibilities**

None reported.

### **6.3 Shelf-Life**

24 months.

### **6.4 Special Precautions for Storage**

Store below 30°C.

### **6.5 Nature and Content of Container**

15g, 30g, 50g and 100g collapsible internally-lacquered, blind-end aluminium tubes, with latex bands and closed with polypropylene caps.

Not all pack sizes may be marketed.

#### **6.6 Special precautions for disposal and other handling**

No special instructions.

### **7 MARKETING AUTHORISATION HOLDER**

Glaxo Wellcome UK Ltd,  
trading as GlaxoSmithKline UK,  
GSK Medicines Research Centre  
Gunnels Wood Road  
Stevenage  
Hertfordshire  
SG1 2NY  
UK

### **8. MARKETING AUTHORISATION NUMBER**

PL 10949/0013

### **9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

17/02/1993 / 30/05/2008

### **10 DATE OF REVISION OF THE TEXT**

06/08/2024