





1. Shake the bottle well and remove the dust cap.
2. Blow your nose gently.
3. Close one nostril as shown and put the nozzle in the other nostril. Tilt your head forward slightly and keep the bottle upright. Hold the bottle as shown (Picture 3).
4. Start to breathe in slowly through your nose. While you are breathing in, squirt a spray of fine mist into your nostril by pressing down firmly on the collar with your fingers.
5. Breathe out through your mouth. Repeat step 4 in the same nostril for a second spray.
6. Remove the nozzle from this nostril and breathe out through your mouth (Picture 4)
7. Repeat steps 3 to 6 for the other nostril (Picture 5).
8. After using the spray, wipe the nozzle carefully with a clean tissue or handkerchief and replace the dust cap.



1. Take the dust cap off.
2. Pull upwards on the white collar to remove the nozzle (Picture 6)
3. Soak the nozzle and dust cap in warm water for a few minutes and then rinse under a running tap.
4. Shake off the excess water and allow to dry in a warm place- not too hot!
5. Re-fit the nozzle.
6. If necessary, press the pump down several times until you get a fine spray



Like all medicines, Beconase Hayfever can have side effects, but not everybody gets them:

Very rare side effects (may affect up to 1 in 10,000 people) reported with Beconase Hayfever include:

- Allergic reactions. These may include developing a rash, swelling of the mouth or face or having difficulty breathing

Rare side effects (may affect up to 1 in 1,000 people) reported with Beconase Hayfever include:

- Eye problems such as pain or blurred vision
- Nasal problems such as pain and/or persistent bleeding
- You may notice an unpleasant taste or smell
- Nose bleeds or irritation in the nose or throat

Frequency not known:

- Blurred vision.

It is important to follow the dose instructions shown in Section 3. Using Beconase Hayfever at higher doses or for longer than 3 months at a time can cause a number of changes in your body such as catching infections more easily.

Reporting of side effects: If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store.

By reporting side effects you can help provide more information on the safety of this medicine.

Keep out of the sight and reach of children.
Do not use this medicine after the 'EXP' date shown on the pack.
Keep the container in the outer carton to protect from light.
Do not store above 25°C (77°F). Do not store in a fridge.
Use within three months of first use.

What Beconase Hayfever contains

Active ingredient Each spray contains Beclometasone Dipropionate 50 micrograms.

Other ingredients Microcrystalline cellulose, sodium carboxymethylcellulose, dextrose, polysorbate 80, purified water, benzalkonium chloride and phenylethyl alcohol.

The plastic bottle contains 180 aqueous nasal sprays.

The marketing authorization holder is
Omega Pharma Ltd., Wrafton, Branton, Devon, EX33 2DL, UK and all enquiries should be sent to this address.

The manufacturers are
Richard Bittner AG, Ossischerstraße 7, 9560 Feldkirchen, Austria.

This leaflet was last revised in May 2023
The information in this leaflet only applies to Beconase Hayfever.



Pharmacode 401 -->

90	File title:	CR08016_LF03499J_Beconase Hayfever 180 Sprays PIL_UK v05
	Item No.:	8100002876
	OPI No.:	LF03499J
	Replacement No.:	8100002836
	Last edit:	23rd May 2023
	Brand:	Beconase
80	Variant:	Hayfever 180 Sprays
	Component:	Leaflet
	Market/s:	UK
70	Dimensions:	304 x 180mm
	Spec. Ref.:	...
	Substrate:	Offset White Paper
60	PharmaCode:	401
	EAN Code:	N/A
	Other Code/Spec.:	...
50	Manufacturing Site:	Richard Bittner
	Printer:	...
	No. of Colours:	1



OneAgency.co

512 Metal Box Factory,
30 Great Guildford Street,
London, SE1 0HS, UK
+44 (0)20 3982 8471
london@oneagency.co