

## SUMMARY OF PRODUCT CHARACTERISTICS

### 1 NAME OF THE MEDICINAL PRODUCT

Zolpidem Tartrate 10mg Film-coated Tablets

### 2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each Film-coated tablet contains 10mg zolpidem tartrate.

Excipient with known effect:

Each Film-coated tablet contains 84.60mg lactose monohydrate.

For the full list of excipients, see section 6.1.

### 3 PHARMACEUTICAL FORM

Film-coated tablets.

White to off-white, capsule shaped, Film-coated tablets having a break line on the one side and plain on the other side. The tablet can be divided into equal doses.

### 4 CLINICAL PARTICULARS

#### 4.1 Therapeutic indications

Zolpidem is indicated for the short-term treatment of insomnia in adults in situations where the insomnia is debilitating or is causing severe distress for the patient.

#### 4.2 Posology and method of administration

##### Posology

The treatment should be taken in a single intake and not be re-administered during the same night.

The recommended daily dose for adults is 10mg to be taken immediately at bedtime. The lowest effective daily dose of zolpidem tartrate should be used and must not exceed 10mg.

Treatment should be as short as possible. It should not exceed four weeks including the period of tapering off. In certain cases, extension beyond the maximum treatment period may be necessary; if so,

extension beyond the maximum treatment period should not take place without re-evaluation of the patient's status since the risk of abuse and dependence increases with the duration of treatment (see section 4.4).

Prior to starting treatment with zolpidem tartrate, a discussion should be held with patients to put in place a strategy for ending treatment with zolpidem tartrate in order to minimise the risk of dependence, addiction and drug withdrawal syndrome (see section 4.4).

Treatment should be given for the shortest possible duration.

### Special populations

#### *Paediatric population*

Zolpidem tartrate is not recommended for use in children and adolescents below 18 years of age, due to a lack of data to support use in this age group. The available evidence from placebo-controlled clinical trials is presented in section 5.1.

#### *Elderly*

Elderly or debilitated patients may be especially sensitive to the effects of zolpidem tartrate, therefore a 5mg dose is recommended. These recommended doses should not be exceeded.

#### *Hepatic impairment*

As clearance and metabolism of zolpidem tartrate is reduced in hepatic impairment, dosage should begin at 5mg in these patients with particular caution being exercised in elderly patients. In adults (under 65 years) dosage may be increased to 10mg only where the clinical response is inadequate, and the drug is well tolerated.

Zolpidem tartrate must not be used in patients with severe hepatic impairment as it may contribute to encephalopathy (see section 4.3).

### Method of administration

Oral administration

## **4.3 Contraindications**

Zolpidem tartrate is contraindicated in patients with

- Hypersensitivity to zolpidem tartrate or any of the excipients listed in section 6.1
- Obstructive sleep apnoea
- Myasthenia gravis
- Severe hepatic insufficiency
- Acute and/or severe respiratory depression

In the absence of data, zolpidem tartrate should not be prescribed for children or patients with psychotic illness.

#### 4.4 Special warning and precautions for use

The cause of insomnia should be identified wherever possible, and the underlying factors treated before a hypnotic is prescribed. The failure of insomnia to remit after a 7-14 day course of treatment may indicate the presence of a primary psychiatric or physical disorder, and the patient should be carefully re-evaluated at regular intervals.

##### Next-day psychomotor impairment

Like other sedative/hypnotic drugs, zolpidem has CNS-depressant effects. The risk of next-day psychomotor impairment, including impaired driving ability, is increased if

- Zolpidem tartrate is taken within less than 8 hours before performing activities that require mental alertness (see section 4.7)
- A dose higher than the recommended dose is taken
- Zolpidem tartrate is co-administered with other CNS depressants or with other drugs that increase the blood levels of zolpidem, or with alcohol or illicit drugs (see section 4.5)

Zolpidem tartrate should be taken in a single intake immediately at bedtime and not be re-administered during the same night.

##### Specific patient groups

###### *Respiratory insufficiency*

As hypnotics have the capacity to depress respiratory drive, precautions should be observed if zolpidem tartrate is prescribed to patients with compromised respiratory function.

###### *Hepatic insufficiency*

See section 4.2.

###### *Elderly*

See section 4.2 for dose recommendations.

###### *Risk from concomitant use of opioids*

Concomitant use of zolpidem tartrate and opioids may result in sedation, respiratory depression, coma and death. Because of these risks, concomitant prescribing of sedative medicines such as benzodiazepines or related drugs such as zolpidem tartrate with opioids should be reserved for patients for whom alternative treatment options are not possible.

If a decision is made to prescribe zolpidem tartrate concomitantly with opioids, the lowest effective dose should be used, and the duration of treatment should be as short as possible (see also general dose recommendation in section 4.2).

The patients should be followed closely for signs and symptoms of respiratory depression and sedation. In this respect, it is strongly recommended to inform patients and others in their environment to be aware of these symptoms (see section 4.5).

#### *Use in patients with a history of drug or alcohol abuse*

Extreme caution should be exercised when prescribing for patients with a history of drug or alcohol abuse. These patients should be under careful surveillance when receiving zolpidem tartrate or any other hypnotic, since they are at risk of habituation and psychological dependence.

#### *Psychotic illness*

Hypnotics such as zolpidem tartrate are not recommended for the primary treatment of psychotic illness.

#### *Suicidal ideation/suicide attempt/suicide and depression*

Some epidemiological studies suggest an increased incidence of suicidal ideation, suicide attempt and suicide in patients with or without depression and treated with benzodiazepines and other hypnotics including zolpidem tartrate. However, a causal relationship has not been established. As with other sedative/hypnotic drugs, zolpidem tartrate should be administered with caution in patients exhibiting symptoms of depression. Suicidal tendencies may be present therefore the least amount of zolpidem tartrate that is feasible should be supplied to these patients to avoid the possibility of intentional overdose by the patient. Pre-existing depression may be unmasked during use of zolpidem tartrate. Since insomnia may be a symptom of depression, the patient should be re-evaluated if insomnia persists.

General information relating to effects seen following administration of benzodiazepines and other hypnotic agents which should be taken into account by the prescribing physician are described below.

#### *Drug dependence, tolerance and potential for abuse*

Drug addiction comprises behavioural, cognitive and physiological phenomena that may include a strong desire to take the drug, difficulties in controlling drug use and possible tolerance or physical dependence. Physical dependence is a state that develops as a result of physiological adaptation in response to repeated drug use, which manifests as withdrawal signs and symptoms after abrupt discontinuation or a significant dose reduction of a drug. Addiction and dependence are related but distinct presentations and in discussing these themes, terminology that apportion blame to the individual should be avoided.

For all patients, prolonged use of this product may lead to drug dependence and addiction but can occur with short-term use at recommended therapeutic doses. The risks are increased in individuals with current or past history of substance misuse disorder (including alcohol misuse) or mental health disorder (e.g., major depression).

Additional support and monitoring may be necessary when prescribing for patients at risk of drug misuse.

A comprehensive patient history should be taken to document concomitant medications, including over-the-counter medicines and medicines obtained on-line, and past and present medical and psychiatric conditions.

Patients may find that treatment is less effective with chronic use and express a need to increase the dose to obtain the same level of symptom control as initially experienced. Patients may also supplement their treatment with additional medications to achieve the same

effect. These could be signs that the patient is developing tolerance. The risks of developing tolerance should be explained to the patient.

Overuse or misuse may result in overdose and/or death. It is important that patients only use medicines that are prescribed for them at the dose they have been prescribed and do not give this medicine to anyone else.

Patients should be closely monitored for signs of misuse, abuse, or addiction.

The clinical need for treatment with zolpidem tartrate should be reviewed regularly, with frequent assessments of patients being undertaken during the course of their treatment.

#### *Drug withdrawal syndrome*

Prior to starting treatment with zolpidem tartrate, a discussion should be held with patients to explain the risk of dependence, addiction, and drug withdrawal syndrome. A withdrawal strategy for ending treatment with zolpidem tartrate should also be put in place with the patient before starting treatment (there may be exceptions to this in specific clinical situations such as symptom management in end of life palliative care).

Drug withdrawal syndrome may occur upon abrupt cessation of therapy or dose reduction. When a patient no longer requires therapy, it is advisable to taper the dose gradually to minimise symptoms of withdrawal. Tapering from a high dose may take in excess of weeks or months. Patients should be informed of this when the medication is first prescribed.

The reduction schedule for a patient should be tailored to the individual and should be modified to allow intolerable withdrawal symptoms to improve before making the next reduction. If using a published withdrawal schedule, apply it flexibly to accommodate the person's preferences, changes to their circumstances and the response to dose reductions.

Suggest a slow stepwise rate of reduction proportionate to the existing dose, so that decrements become smaller as the dose is lowered, unless clinical risk is such that rapid withdrawal is needed.

#### *Rebound insomnia*

A transient syndrome whereby the symptoms that led to treatment with a benzodiazepine or benzodiazepine-like agent recur in an enhanced form may occur on withdrawal of hypnotic treatment. It may be accompanied by other reactions including mood changes, anxiety, and restlessness.

It is important that the patient should be aware of the possibility of rebound phenomena, thereby minimising anxiety over such symptoms should they occur when the medicinal product is discontinued. Since the risk of withdrawal phenomena or rebound has been shown to be greater after abrupt discontinuation of treatment, it is recommended that the dosage is decreased gradually where clinically appropriate.

There are indications that, in the case of benzodiazepines and benzodiazepine-like agents with a short duration of action, withdrawal phenomena can become manifest within the dosage interval, especially when the dosage is high.

#### *Amnesia*

Benzodiazepines or benzodiazepine-like agents such as zolpidem tartrate may induce anterograde amnesia. The condition occurs most often several hours after ingesting the product and therefore to reduce the risk, patients should ensure that they will be able to have an uninterrupted sleep of 8 hours (see section 4.8).

#### *Other psychiatric and “paradoxical” reactions*

Other psychiatric and paradoxical reactions like restlessness, exacerbated insomnia, agitation, irritability, aggression, delusion, anger, nightmares, hallucinations, psychosis, abnormal behaviour and other adverse behavioural effects are known to occur when using benzodiazepines or benzodiazepine-like agents. Should this occur, use of the product should be discontinued. These reactions are more likely to occur in the elderly.

#### *Somnambulism and associated behaviours*

Sleep-walking and other associated behaviours such as “sleep driving”, preparing and eating food, making phone calls, or having sex, with amnesia for the event, have been reported in patients who had taken zolpidem tartrate and were not fully awake. The use of alcohol and other CNS-depressants with zolpidem tartrate appears to increase the risk of such behaviour, as does the use of zolpidem tartrate at doses exceeding the maximum recommended dose. Discontinuation of zolpidem should be strongly considered for patients who report such behaviour (for example, sleep driving), due to the risk to the patient and others.

#### *Severe injuries*

Due to its pharmacological properties, zolpidem tartrate can cause drowsiness and a decreased level of consciousness, which may lead to falls and consequently to severe injuries, see section 4.8.

#### Excipients

Zolpidem tartrate contains lactose monohydrate. Patients with rare hereditary problems of galactose intolerance, total lactase deficiency, or glucose-galactose malabsorption should not take this medicine.

This medicine contains less than 1 mmol sodium (23mg) per tablet, that is to say essentially ‘sodium-free’.

## **4.5 Interaction with other medicinal products and other forms of interaction**

#### Not recommended

##### *Concomitant intake with alcohol*

The sedative effect may be enhanced when the product is used in combination with alcohol. This affects the ability to drive or use machines.

#### Take into account

##### *Combination with CNS depressants*

Enhancement of the central depressive effect may occur in cases of concomitant use with antipsychotics (neuroleptics), hypnotics, anxiolytics/sedatives, antidepressant agents, narcotic analgesics, antiepileptic drugs, anaesthetics and sedative antihistamines. Therefore, concomitant use of zolpidem tartrate with these drugs may increase

drowsiness and next-day psychomotor impairment, including impaired driving ability (see sections 4.4 and 4.7). Also, isolated cases of visual hallucinations were reported in patients taking zolpidem tartrate with antidepressants including bupropion, desipramine, fluoxetine, sertraline and venlafaxine.

Co-administration of fluvoxamine may increase blood levels of zolpidem tartrate, concurrent use is not recommended.

In the case of narcotic analgesics, enhancement of euphoria may also occur, leading to an increase in psychological dependence.

### *Opioids*

The concomitant use of sedative medicines such as benzodiazepines or related drugs such as zolpidem tartrate with opioids increases the risk of sedation, respiratory depression, coma and death because of additive CNS depressant effect. The dosage and duration of concomitant use should be limited (see section 4.4).

### *CYP450 inhibitors and inducers*

Co-administration of ciprofloxacin may increase blood levels of zolpidem, concurrent use is not recommended.

Compounds which inhibit certain hepatic enzymes (particularly cytochrome P450) may enhance the activity of benzodiazepines and benzodiazepine-like agents.

Zolpidem tartrate is metabolised via several hepatic cytochrome P450 enzymes, the main enzyme being CYP3A4 with the contribution of CYP1A2. The pharmacodynamic effect of zolpidem tartrate is decreased when it is administered with a CYP3A4 inducer such as rifampicin and St. John's Wort. St. John's Wort has been shown to have a pharmacokinetic interaction with zolpidem tartrate. Mean  $C_{max}$  and AUC were decreased (33.7 and 30.0% lower, respectively) for zolpidem administered with St. John's Wort compared to zolpidem administered alone. Co-administration of St. John's Wort may decrease blood levels of zolpidem tartrate, concurrent use is not recommended.

However, when zolpidem tartrate was administered with itraconazole (a CYP3A4 inhibitor) its pharmacokinetics and pharmacodynamics were not significantly modified. The clinical relevance of these results is unknown. Co-administration of zolpidem tartrate with ketoconazole (200mg twice daily), a potent CYP3A4 inhibitor, prolonged zolpidem tartrate elimination half-life, increased total AUC, and decreased apparent oral clearance when compared to zolpidem tartrate plus placebo. The total AUC for zolpidem tartrate when co-administered with ketoconazole, increased by a factor of 1.83 when compared to zolpidem tartrate alone. A routine dosage adjustment of zolpidem tartrate is not considered necessary, but patients should be advised that use of zolpidem tartrate with ketoconazole may enhance the sedative effects.

Since CYP3A4 plays an important role in zolpidem metabolism, possible interactions with drugs that are substrates or inducers of CYP3A4 should be considered.

#### *Other drugs*

When zolpidem tartrate was administered with ranitidine, no significant pharmacokinetic interactions were observed.

## **4.6 Fertility, pregnancy and lactation**

### Pregnancy

The use of zolpidem tartrate is not recommended during pregnancy.

Animal studies do not indicate direct or indirect harmful effects with respect to reproductive toxicity.

Zolpidem tartrate crosses the placenta.

A large amount of data on pregnant women (more than 1000 pregnancy outcomes) collected from cohort studies has not demonstrated evidence of the occurrence of malformations following exposure to benzodiazepines or benzodiazepine-like substances during the first trimester of pregnancy. However, certain case-control studies reported an increased incidence of cleft lip and palate associated with the use of benzodiazepines during pregnancy.

Cases of reduced foetal movement and foetal heart rate variability have been described after administration of benzodiazepines or benzodiazepine-like substances during the second and/or third trimester of pregnancy.

Administration of zolpidem tartrate during the late phase of pregnancy or during labour has been associated with effects on the neonate, such as hypothermia, hypotonia, feeding difficulties (“floppy infant syndrome”), and respiratory depression, due to the pharmacological action of the product. Cases of severe neonatal respiratory depression have been reported.

Moreover, infants born to mothers who took sedative/hypnotic agents chronically during the latter stages of pregnancy may have developed physical dependence and may be at risk of developing withdrawal symptoms in the postnatal period. Appropriate monitoring of the newborn in the postnatal period is recommended.

If zolpidem tartrate is prescribed to a woman of childbearing potential, she should be warned to contact her physician about stopping the product if she intends to become or suspects that she is pregnant.

### Breast-feeding

Small quantities of zolpidem tartrate appear in breast milk. The use of zolpidem tartrate in nursing mothers is therefore not recommended.

## **4.7 Effects on ability to drive and use machines**

Zolpidem tartrate has major influence on the ability to drive and use machines.

Vehicle drivers and machine operators should be warned that, as with other hypnotics, there may be a possible risk of drowsiness, prolonged reaction time, dizziness, sleepiness, blurred/double vision and reduced alertness and impaired driving the morning after therapy (see section 4.8). In order to minimise this risk, a resting period of at least 8 hours is recommended between taking zolpidem tartrate and driving, using machinery and working at heights.

Driving ability impairment and behaviours such as 'sleep-driving' have occurred with zolpidem tartrate alone at therapeutic doses.

Furthermore, co-administration of zolpidem tartrate with alcohol and other CNS depressants increases the risk of such behaviours (see sections 4.4 and 4.5). Patients should be warned not to use alcohol or other psychoactive substances when taking zolpidem tartrate.

#### **4.8 Undesirable effects on ability to drive and use machines**

The following CIOMS frequency rating is used, when applicable:

*Very common*  $\geq 10\%$

*Common*  $\geq 1$  and  $< 10\%$

*Uncommon*  $\geq 0.1$  and  $< 1\%$

*Rare*  $\geq 0.01$  and  $< 0.1\%$

*Very rare*  $< 0.01\%$

*Not known: cannot be estimated based on available data.*

There is evidence of a dose-relationship for adverse effects associated with zolpidem tartrate use, particularly for certain CNS and gastrointestinal events. As recommended in section 4.2 they should in theory be less if zolpidem tartrate is taken immediately before retiring, or in bed. They occur most frequently in elderly patients.

##### Immune system disorders

*Not known:* angioneurotic oedema

##### Psychiatric disorders

*Common:* hallucination, agitation, nightmare, depression (see section 4.4)

*Uncommon:* confusional state, irritability, restlessness, aggression, somnambulism (see section 4.4), euphoric mood

*Rare:* libido disorder

*Very rare:* delusion, drug dependence (withdrawal symptoms, or rebound effects may occur after treatment discontinuation, see section 4.4)

*Not known:* anger, psychosis, abnormal behaviour

Most of these psychiatric undesirable effects are related to paradoxical reactions.

##### Nervous system disorders

*Common:* somnolence, headache, dizziness, exacerbated insomnia, cognitive disorders such as anterograde amnesia (amnesic effects may be associated with inappropriate behaviour)

*Uncommon:* paraesthesia, tremor, disturbance in attention, speech disorder

*Rare:* depressed level of consciousness

#### Eye disorders

*Uncommon:* diplopia, vision blurred

*Very rare:* visual impairment

#### Skin and subcutaneous tissue disorders

*Uncommon:* rash, pruritus, hyperhidrosis

*Rare:* urticaria

#### Musculoskeletal and connective tissue disorders

*Common:* back pain

*Uncommon:* myalgia, muscle spasms, muscular weakness

#### Infections and infestations

*Common:* upper respiratory tract infection, lower respiratory tract infection

#### General disorders and administration site conditions

*Common:* fatigue

*Rare:* gait disturbance, fall (predominantly in elderly patients and when zolpidem tartrate was not taken in accordance with prescribing recommendation) (see section 4.4).

*Not known:* drug tolerance

#### Withdrawal syndrome

Withdrawal syndrome has been reported upon discontinuation of zolpidem tartrate (see section 4.4). Withdrawal symptoms vary and may include rebound insomnia, muscle pain, anxiety, tremor, sweating, agitation, confusion, headache, palpitations, tachycardia, delirium, nightmares, hallucinations, panic attacks, muscle aches/cramps, gastrointestinal disturbances and irritability. In severe cases the following symptoms may occur: derealisation, depersonalisation, hyperacusis, numbness and tingling of the extremities, hypersensitivity to light, noise and physical contact, hallucinations. In very rare cases, seizures may occur.

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via Yellow Card Scheme Website: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store.

## 4.9 Overdose

Patients should be informed of the signs and symptoms of overdose and to ensure that family and friends are also aware of these signs and to seek immediate medical help if they occur.

### Signs and Symptoms

In cases of overdose, involving zolpidem tartrate alone or with other CNS-depressant agents (including alcohol), impairment of consciousness ranging from somnolence to coma, and more severe symptomatology, including fatal outcomes have been reported.

### Management

General symptomatic and supportive measures should be used. If there is no advantage in emptying the stomach, activated charcoal should be given to reduce absorption. Sedating drugs should be withheld even if excitation occurs.

Use of flumazenil may be considered where serious symptoms are observed. Flumazenil is reported to have an elimination half-life of about 40 to 80 minutes. Patients should be kept under close observation because of this short duration of action; further doses of flumazenil may be necessary. However, flumazenil administration may contribute to the appearance of neurological symptoms (convulsions).

Zolpidem tartrate is not dialyzable.

The value of dialysis in the treatment of an overdose has not been determined. Dialysis in patients with renal failure receiving therapeutic doses of zolpidem tartrate has demonstrated no reduction in levels of zolpidem tartrate.

In the management of overdose with any medicinal product, it should be borne in mind that multiple agents may have been taken.

## 5 PHARMACOLOGICAL PROPERTIES

### 5.1 Pharmacodynamic properties

Pharmacotherapeutic group: GABA-A receptor agonist selective for omega-1-type sub-unit hypnotic agent, ATC code: N05CF02.

Zolpidem tartrate is an imidazopyridine which selectively binds the omega-1 receptor subtype (also known as the benzodiazepine-1 subtype) which is the alpha unit of the GABA-A receptor complex. Whereas benzodiazepines non-selectively bind all three omega receptor subtypes, zolpidem tartrate preferentially binds the omega-1 subtype. The modulation of the chloride anion channel via this receptor leads to the

specific sedative effects demonstrated by zolpidem tartrate. These effects are reversed by the benzodiazepine antagonist flumazenil.

The randomized trials only showed convincing evidence of efficacy of 10mg zolpidem tartrate. In a randomized double-blind trial in 462 non-elderly healthy volunteers with transient insomnia, zolpidem tartrate 10mg decreased the mean time to fall asleep by 10 minutes compared to placebo, while for 5mg zolpidem tartrate this was 3 minutes.

In a randomized double-blind trial in 114 non-elderly patients with chronic insomnia, zolpidem tartrate 10mg decreased the mean time to fall asleep by 30 minutes compared to placebo, while for 5mg zolpidem tartrate this was 15 minutes.

In some patients, a lower dose of 5mg could be effective.

*In animals:* The selective binding of zolpidem tartrate to omega-1 receptors may explain the virtual absence at hypnotic doses of myorelaxant and anti-convulsant effects in animals which are normally exhibited by benzodiazepines which are not selective for omega-1 sites.

*In humans:* The preservation of deep sleep (stages 3 and 4-slow-wave sleep) may be explained by the selective omega-1 binding by zolpidem tartrate. All identified effects of zolpidem tartrate are reversed by the benzodiazepine antagonist flumazenil.

#### Paediatric population

Safety and efficacy of zolpidem tartrate has not been established in children aged less than 18 years. A randomized placebo-controlled study in 201 children aged 6-17 years with insomnia associated with Attention Deficit Hyperactivity Disorder (ADHD) failed to demonstrate efficacy of zolpidem tartrate 0.25mg/kg/day (with a maximum of 10mg/day) as compared to placebo.

Psychiatric and nervous system disorders comprised the most frequent treatment emergent adverse events observed with zolpidem tartrate versus placebo and included dizziness (23.5% versus 1.5%), headache (12.5% versus 9.2%), and hallucinations (7.4% versus 0%) (see sections 4.2 and 4.3).

## **5.2 Pharmacokinetic properties**

Zolpidem tartrate has both a rapid absorption and onset of hypnotic action. Bioavailability is 70% following oral administration and demonstrates linear kinetics in the therapeutic dose range. Peak plasma concentration is reached at between 0.5 and 3 hours.

The elimination half-life is short, with a mean of 2.4 hours (0.7-3.5) and a duration of action of up to 6 hours.

Protein binding amounts to 92.5% ± 0.1%. First pass metabolism by the liver amounts to approximately 35%. Repeated administration has been

shown not to modify protein binding indicating a lack of competition between zolpidem tartrate and its metabolites for binding sites.

The distribution volume in adults is  $0.54 \pm 0.02$  L/kg and decreases to  $0.34 \pm 0.05$  L/kg in the very elderly.

All metabolites are pharmacologically inactive and are eliminated in the urine (56%) and in the faeces (37%).

Zolpidem tartrate has been shown in trials to be non-dialysable.

Plasma concentrations in elderly subjects and those with hepatic impairment are increased. In patients with renal insufficiency, whether dialysed or not, there is a moderate reduction in clearance. The other pharmacokinetic parameters are unaffected.

Zolpidem tartrate is metabolised via several hepatic cytochrome P450 enzymes, the main enzyme being CYP3A4 with the contribution of CYP1A2. Since CYP3A4 plays an important role in zolpidem metabolism, possible interactions with drugs that are substrates or inducers of CYP3A4 should be considered.

### **5.3 Preclinical safety data**

Preclinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, repeated dose toxicity, genotoxicity, carcinogenic potential, toxicity to reproduction.

## **6 PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Tablet core:

Lactose monohydrate

Microcrystalline cellulose PH101

Sodium starch glycolate (Type A)

Hypromellose (HPMC E5)

Magnesium stearate

Film-coating:

Hypromellose (HPMC E3)

Macrogol (PEG-400)

Titanium dioxide (E171)

**6.2 Incompatibilities**

Not applicable

**6.3 Shelf life**

3 years.

**6.4 Special precautions for storage**

This medicinal product does not require any special storage conditions.

**6.5 Nature and contents of container**

Aluminium/-PVC blisters containing 28 tablets.

HDPE tablets containers with child-resistant polypropylene cap containing 400 tablets.

Not all packs sizes may be marketed.

**6.6 Special precautions for disposal**

No special requirements.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

**7 MARKETING AUTHORISATION HOLDER**

Key Pharmaceuticals Ltd.

18 Oxleasow Road,

Redditch, B98 0RE,

United Kingdom

**8 MARKETING AUTHORISATION NUMBER(S)**

PL 34424/0010

**9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

29/07/2021

**10 DATE OF REVISION OF THE TEXT**

27/05/2026