

Fosrenol® 750 mg chewable tablets

(lanthanum carbonate)

XXXXXXXXXXXX

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet:

1. What Fosrenol is and what it is used for
2. What you need to know before you take Fosrenol
3. How to take Fosrenol
4. Possible side effects
5. How to store Fosrenol
6. Contents of the pack and other information

1. What Fosrenol is and what it is used for

Fosrenol is used to lower the phosphate level in the blood of adult patients with chronic kidney disease.

Patients who have kidneys that do not work properly are not able to control the level of phosphate in the blood. The amount of phosphate in the blood then rises (your doctor may call this hyperphosphataemia).

Fosrenol is a drug which reduces the body's absorption of phosphate from food by binding with it in your digestive tract. Phosphate which has bonded to Fosrenol cannot be absorbed through the intestinal wall.

2. What you need to know before you take Fosrenol

Do not take Fosrenol

- if you are allergic to lanthanum carbonate hydrate or any of the other ingredients of this medicine (listed in section 6).
- if you have too little phosphate in your blood (hypophosphataemia)

Warnings and precautions

Talk to your doctor or pharmacist before taking Fosrenol if you know that you have, or have had, any of the following:

- stomach or intestinal cancer
- inflammatory bowel disease including ulcerative colitis or Crohn's disease
- abdominal surgery, or infection or inflammation of the abdomen/bowel (peritonitis)
- stomach or intestinal ulcers
- blockage of the intestine or slow motility (movement) in the intestine (e.g., constipation and stomach complications due to diabetes)
- reduced liver or kidney function.

It is very important to chew completely Fosrenol tablets and not to swallow them whole or incompletely chewed. This will reduce the risk of adverse gastrointestinal complications like rupture in the intestine wall, blockage in the intestine, constipation (see section 4).

If you have reduced kidney function, your doctor may decide to check the level of calcium in your blood from time to time. If you have too little calcium, you may then be given extra calcium.

If you need to have an x-ray, please inform your doctor that you are taking Fosrenol as it may affect the results.

Other medicines and Fosrenol

Tell your doctor or pharmacist if you are taking, have recently taken, or might take any other medicines.

Fosrenol can affect how certain drugs are absorbed from your digestive tract. If you are taking chloroquine (for rheumatism and malaria), ketoconazole (for fungal infections), tetracycline or doxycycline antibiotics they should not be taken within 2 hours before or after taking Fosrenol.

It is not recommended that you take oral floxacillin antibiotics (including ciprofloxacin) within 2 hours before or 4 hours after taking Fosrenol.

If you are taking levothyroxine (for an under active thyroid) it should not be taken within 2 hours before or after taking Fosrenol. Your doctor may want to monitor the levels of thyroid-stimulating hormone (TSH) in your blood more closely.

Fosrenol with food and drink

Fosrenol should be taken with, or immediately after food. See section 3 for instructions on how to take Fosrenol.

Pregnancy and breast-feeding

Fosrenol should not be taken during pregnancy. If you are pregnant or breast-feeding, think you may be pregnant, or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

As it is not known whether the drug can be transferred to a child in breast-milk, you should not breast-feed whilst taking Fosrenol. If you are breast-feeding, ask your doctor or pharmacist for advice before taking any medicines.

Driving and using machines

Dizziness and vertigo (a feeling of dizziness or "spinning") are uncommon side effects reported by patients taking Fosrenol. If you experience these side effects, it may affect your ability to drive or operate machinery.

Fosrenol contains Glucose

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

3. How to take Fosrenol

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

You should take Fosrenol with, or immediately after food. Side effects such as nausea and vomiting are more likely if you take Fosrenol before your meal.

The tablets must be chewed completely and not swallowed whole. To aid with chewing, the tablets may be crushed. Additional fluid is not necessary.

If you find chewing the tablets difficult, talk to your doctor as an oral powder form of this medicine is available.

Your doctor will tell you how many tablets you must take with each meal (your daily dose will be divided between meals). The number of tablets that you take will depend on:

- Your diet (the amount of phosphate in the food you eat)
- Your blood phosphate level

To start with, the daily dose of Fosrenol will usually be 1 tablet with each meal (3 tablets per day).

Every 2-3 weeks your doctor will check the level of phosphate in your blood and may increase your dose until the level of phosphate in your blood is acceptable. Fosrenol works by binding phosphate from the food in your gut. It is very important to take Fosrenol at every meal. If you change your diet, contact your doctor as you may need to take extra Fosrenol. Your doctor will tell you what to do in this case.

If you take more Fosrenol than you should

If you take too many tablets contact your doctor to assess the risk and obtain advice. Symptoms of overdose may be nausea and headaches.

If you forget to take Fosrenol

It is important to take Fosrenol with every meal.

If you forget to take your Fosrenol tablets, then take the next dose with your next meal. Do not take a double dose to make up for a forgotten dose.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Some side effects could be serious. If you get any of the following side effects, seek immediate medical attention:

- Rupture in the intestinal wall (signs include: severe stomach pain, chills, fever, nausea, vomiting, or a tender abdomen). This is a rare side effect (may affect up to 1 in 1,000 people).
- Blockage in the intestine (signs include: severe bloating; abdominal pain, swelling or cramps; severe constipation). This is an uncommon side effect (may affect up to 1 in 100 people).
- Contact your doctor if you experience new or severe constipation, it could be an early sign of a blockage in your intestine. Constipation is a common side effect (may affect 1 in 10 people).

Other less serious side effects include the following:

Very Common side effects (may affect more than 1 in 10 people):

- Nausea, vomiting, diarrhoea, stomach pain, headache, itching, rash.

Common side effects (may affect up to 1 in 10 people):

- Heartburn, flatulence.
- Hypocalcaemia (too little calcium in your blood) is also a common side effect; the symptoms of which can include tingling in the hands and feet, muscle and abdominal cramps or spasms of the facial and feet muscles.

Uncommon side effects (may affect up to 1 in 100 people):

- Tiredness; feeling of discomfort; chest pain, weakness; swollen hands and feet; body pain; dizziness; vertigo; belching; inflammation of the stomach and intestines (gastroenteritis); indigestion; irritable bowel syndrome; dry mouth; tooth disorders; inflammation of the gullet or mouth; loose stools; increases in certain liver enzymes, parathyroid hormone; aluminium, calcium and glucose in the blood; increased or reduced phosphate level in the blood; thirst; weight decrease; joint pain; muscle pain; weakness and thinning of the bones (osteoporosis); lack of and increased appetite; inflammation of the larynx; loss of hair; increased sweating; taste disturbance and increased white blood cell count.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the **Google Play** or **Apple App Store**. By reporting side effects, you can help provide more information on the safety of this medicine.

5. How to store Fosrenol

Keep this medicine out of the sight and reach of children.

This medicinal product does not require any special storage conditions.

Do not use this medicine after the expiry date which is stated on the carton and the bottle label. The expiry date refers to the last day of that month.

If this medicine becomes discoloured or shows any other signs of deterioration, please contact your pharmacist who will advise you on what to do.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Fosrenol contains

- The active substance is 750 mg lanthanum (as lanthanum carbonate hydrate).
- The other ingredients are dextrans (hydrated), colloidal anhydrous silica and magnesium stearate.

What Fosrenol looks like and contents of the pack

Fosrenol is a white, round, bevelled-edged flat chewable tablet debossed with 'S405/750' on one side of the tablet. The tablets are supplied in plastic bottles. A multipack containing 90 (6 packs of 15) chewable tablets (750 mg).

Product Licence Holder and Manufacturer

Procured from within the EU by the Product Licence holder Orifarm A/S, Energivej 15, 5260 Odense S, Denmark

Repacked by

Orifarm Supply s.r.o., Palouky 1366, 253 01 Hostivice, Czech Republic

Manufactured by

Hamol Limited, Nottingham site, Thane Road, Nottingham, Nottinghamshire NG90 2DB, UK

RB NL Brands B.V., Schiphol Boulevard 207, 1118BH, Schiphol, The Netherlands

Fosrenol 750 mg chewable tablets
PL 46927/0211

POM

Leaflet revision date: 07/12/2021

**Blind or partially sighted?
Is this leaflet hard to see or read?
Call +45 63 95 27 00
to obtain the leaflet in a format suitable
for you.**