

160 mm

Package Leaflet: Information for the user

Quinine Sulfate 200mg and 300mg Film coated Tablets

PHARMACODE

XXXX

Pharmaco-Reading Direction

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet:

1. What Quinine sulfate tablets are and what they used for
2. What you need to know before you take Quinine sulfate tablets
3. How to take Quinine sulfate tablets
4. Possible side effects
5. How to store Quinine sulfate tablets
6. Contents of the pack and other information

1. What Quinine sulfate tablets are and what they are used for

Quinine Sulfate 200mg and 300mg Film coated Tablets also called as Quinine sulfate tablets in this leaflet belongs to a group of medicines called antiprotozoal agents and they are used:

- To treat malaria
- To treat and prevent night cramps in adults and the elderly when sleep is regularly disrupted

2. What you need to know before you take Quinine sulfate tablets

Do not take Quinine sulfate tablets if you:

- are allergic to quinine (including that in tonic waters or other beverages), quinoline or any of the other ingredients of this medicine (listed in section 6)
- have blood in your urine
- have ringing in your ears
- suffer from muscle weakness (myasthenia gravis)
- have problems with your eyes or difficulty seeing
- Have been told you have a disorder affecting the red blood cells (haemolysis).

Warnings and precautions

Talk to your doctor, pharmacist or nurse before taking Quinine sulfate tablets if you:

- have irregular heartbeats or other heart disease
- have had malaria for a long time
- suffer from severe glucose-6-phosphate dehydrogenase deficiency (G6PD), as this can cause episodes of anaemia after eating certain foods such as fava beans (favism) or take certain drugs including dapsone and drugs to prevent malaria
- Were born with or have any condition that causes an abnormal heart rhythm.

You should not take more than the prescribed dose as a condition called 'cinchonism' may occur, even with normal doses. Please see section 4 'Possible side effects' for symptoms of cinchonism and tell your doctor if you experience any of them.

Other medicines and Quinine sulfate tablets

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. Especially:

- anticoagulants (to stop your blood from clotting)
- cardiac glycosides (for your heart, such as digoxin)
- chloroquine, mefloquine, artemether with lumefantrine, primaquine, halofantrine (also to treat malaria)
- cimetidine (to treat stomach ulcers or acid reflux and indigestion)
- amantadine (to treat Parkinson's Disease or some viral infections)

- ciclosporin (to prevent transplant rejection)
- flecainide, quinidine or amiodarone (to treat irregular heartbeats)
- astemizole or terfenadine (for allergic reactions)
- pimozone or thioridazine (to treat some mental disorders)
- moxifloxacin, rifampicin or antifungals (to treat infections)
- medicines to treat diabetes
- suxamethonium (muscle relaxant)
- HIV medicines
- barbiturates, carbamazepine or phenytoin (used in epilepsy)
- medicines which are known to cause disturbances in heart rhythm
- levacetylmethadol (a pain killer).

Pregnancy, breast-feeding and fertility

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. You should not take quinine sulfate when you are pregnant or breast-feeding, unless the benefits outweigh the risks. This medicine should not be used to treat cramps during pregnancy.

Large doses of quinine can cause foetal abnormalities or induce abortion. You should not breast-feed a baby that might have glucose-6 phosphate dehydrogenase (G6PD) deficiency.

Driving and using machines

Quinine sulfate tablets may affect your vision and cause vertigo (dizziness or spinning sensation). Make sure you are not affected before you drive or operate machinery.

Information on sodium content

This medicine contains less than 1 mmol (23mg) sodium per tablet, that is to say essentially 'sodium-free'.

3. How to take Quinine sulfate tablets

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure. Swallow the tablets with water.

The recommended dose is:

- **Malaria** (you may be given another medicine for malaria with or after this course of quinine):
Adults (including elderly) and children over 12 years - 600mg every eight hours for 5-7 days
Children aged 10-12 years - 10mg per kg of body weight every eight hours for 7 days
Children under 10 years - Not recommended.
If you have kidney or liver disease you may be given a different dose.

• **Night cramps: Adults (including elderly)** - The recommended dose is 200mg at bedtime. The maximum dose is 300mg. It may take up to 4 weeks before you notice any reduction in the frequency of leg cramps.

If you take more Quinine sulfate tablets than you should

If you (or someone else) swallow a lot of tablets at the same time, or you think a child may have swallowed any, contact your nearest hospital casualty department or tell your doctor immediately.

Symptoms include loss of consciousness, difficulty breathing, changes in heartbeat and heart function, kidney failure, fits, 'ringing' in the ears, loss of hearing, headache, being sick, changes in vision, low blood potassium (causing muscle twitching/weakness or abnormal heart rhythm), low blood sugar and sweating. Some of these side effects can be fatal. High doses of quinine may also cause miscarriage in pregnant women.

If you forget to take Quinine sulfate tablets

Do not take a double dose to make up for a forgotten dose. If you forget to take a dose, take it as soon as you remember it and then take the next dose at the right time.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Contact your doctor immediately if the following effects occur:

- **allergic reactions** - itchy skin rash, swelling of the lips, face, throat or tongue, flushing, fever, asthma or sensitivity to light
- sores in the mouth, nose, and eyes, a skin rash, blistering (**Stevens-Johnson syndrome**)
- **cinchonism** - abdominal pain, diarrhoea, disturbed vision (blurred vision, changes in colour perception or field of vision, total blindness), headache, feeling or being sick, ringing in the ears or impaired hearing, rashes, loss of consciousness, fits, shock due to heart problems, irregular heartbeats. If these occur while taking Quinine sulfate tablets for leg cramps, treatment should be stopped and a doctor contacted straight away. This side effect can be fatal.
- Changes to blood cells as seen in blood or urine tests. If you notice that you are bruising or bleeding easily, have frequent nose bleeds, or you have more sore throats and infections than usual, tell your doctor who may want to give you a blood test.
- Muscle weakness, excitement, agitation, 'spinning' sensation, confusion, loss of consciousness. This can lead to coma, and may be fatal.

Tell your doctor if you notice any of the following side effects or notice any other effects not listed:

- diarrhoea, feeling or being sick, abdominal pain, low blood sugar.
- tightness in chest, difficulty breathing or wheezing
- headache, changes in vision, 'ringing' in the ears, loss of hearing.
- swollen, itchy, flaky, red or raised patches of skin, rashes, sensitivity to light.
- aggravation of Myasthenia gravis (disorder characterised by muscle weakness, difficulty chewing or swallowing and slurred speech).
- kidney damage, kidney failure, low production of urine, water retention, slowed heart rate, changes in heart rhythm and the way the heart beats, a fall in blood pressure and weak pulse, eczema, miscarriages (at very high concentrations).

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme Website: www.mhra.gov.uk/yellowcard or search for

MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Quinine sulfate tablets

Keep out of the sight and reach of children. Store in the original package to protect from light and moisture. Do not use take this medicine after the expiry date stated on the carton. The expiry date refers to the last day of that month. Do not throw away medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Quinine sulfate tablets contain

The active substance (the ingredient that makes the tablet work) is 200mg quinine sulfate. The active substance (the ingredient that makes the tablet work) is 300mg quinine sulfate. The other ingredients are microcrystalline cellulose, croscarmellose sodium, povidone, sodium laurilsulfate, purified Talc, hydrogenated vegetable oil, magnesium stearate. The coating material OPADRY TF 266G580006-White contains: hypromellose (E464), hydroxypropyl cellulose (E463), macrogol (E1521), medium chain triglycerides, rice starch.

What Quinine sulfate tablets look like and contents of the pack

Quinine sulfate 200mg Film coated tablet are white to off white film-coated, round shaped, biconvex tablet, plain on both sides, with diameter of 9.70mm.

Quinine sulfate 300mg Film coated tablet are white to off white film-coated, round shaped, biconvex tablet, debossed with 'L' on one side and plain on other side, with diameter of 11.10mm.

These tablets will be packed in Aluminium-PVC White opaque film blister packs of 28s, 30s, 56s, 60s, 84s, 90s, 100s, 112s and 120s. Not all pack sizes may be marketed.

Marketing Authorization Holder

Flamingo Pharma (UK) Ltd.
First Floor, Kirkland House,
11-15 Peterborough Road,
Harrow, Middlesex,
HA1 2AX, United Kingdom.

Manufacturer

Flamingo Pharma (UK) Limited
The Bloc, 38 Springfield Way,
Anlaby, Hull, HU10 6RJ,
United Kingdom

This leaflet was last revised in 01/2026.

To request a copy of this leaflet in Braille, large print or audio please call free of charge: 0800 198 5000 (UK only).

Please be ready to give the following information:

Product Name: Quinine Sulfate 200mg and 300mg Film coated Tablets

Reference Number: PL 43461/0140 and PL 43461/0141

This is a service provided by the Royal National Institute of Blind People.

MPLLQUIXXXXTBCOM
FPLXXX407V01_A

POM

PHARMACODE

XXXX

Pharmaco-Reading Direction

160 mm

300 mm

300 mm