4437

Colour Black

LOCATION : Indrad	COUNTRY : UK	Supersedes A/W No.:				
SIZE : 150 x 375 mm_Front/Back Side		CODE : 8087460-7803	DATE : 19-09-2025			
REMARK : Folding Length 35 mm						
SUBSTRATE :						
Activities	Department	Name	Signature	Date		
Prepared By	Pkg.Dev					
Reviewed By	Pkg.Dev					
Approved By	Quality					

This colour proof is not colour binding. Follow Pantone shade reference for actual colour matching.

Package leaflet: Information for the patient

Alogliptin Torrent 12.5 mg film-coated tablets
alogliptin

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.
- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist .
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

This product is referred to as Alogliptin Torrent tablet in this leaflet.

What is in this leaflet

1. What Alogliptin Torrent is and what it is used for
2. What you need to know before you take Alogliptin Torrent
3. How to take Alogliptin Torrent
4. Possible side effects
5. How to store Alogliptin Torrent
6. Contents of the pack and other information

1. What Alogliptin Torrent is and what it is used for

Alogliptin Torrent contains the active substance alogliptin which belongs to a group of medicines called DPP-4 inhibitors (dipeptidyl peptidase-4 inhibitors) which are “oral anti-diabetics”. It is used to lower blood sugar levels in adults with type 2 diabetes. Type 2 diabetes is also called non-insulin-dependent diabetes mellitus or NIDDM.

Alogliptin Torrent works to increase the levels of insulin in the body after a meal and decrease the amount of sugar in the body. It must be taken together with other anti-diabetic medicines, which your doctor will have prescribed for you, such as sulphonylureas (e.g. glipizide, tolbutamide, glibenclamide), metformin and/or thiazolidinediones (e.g. pioglitazone) and metformin and/or insulin.

Alogliptin Torrent is taken when your blood sugar cannot be adequately controlled by diet, exercise and one or more of these other oral anti-diabetic medicines. It is important that you continue to take your other anti-diabetic medicine, and continue to follow the advice on diet and exercise that your nurse or doctor has given you.

2. What you need to know before you take Alogliptin Torrent

Do not take Alogliptin Torrent
- if you are allergic to alogliptin or any of the other ingredients of this medicine (listed in section 6)
- if you have had a serious allergic reaction to any other similar medicines that you take to control your blood sugar. Symptoms of a serious allergic reaction may include; rash, raised red patches on your skin (hives), swelling of the face, lips, tongue, and throat that may cause difficulty in breathing or swallowing. Additional symptoms may include general itching and feeling of heat especially affecting the scalp, mouth, throat, palms of hands and soles of feet (Stevens-Johnson syndrome).

Warnings and precautions
Talk to your doctor or pharmacist before taking Alogliptin Torrent:
- if you have type 1 diabetes (your body does not produce insulin)
- if you have diabetic ketoacidosis (a complication of diabetes that occurs when the body is unable to breakdown glucose because there is not enough insulin). Symptoms include excessive thirst, frequent urination, loss of appetite, nausea or vomiting and rapid weight loss
- if you are taking an anti-diabetic medicine known as sulphonylurea (e.g. glipizide, tolbutamide, glibenclamide) or insulin. Your doctor may want to reduce your dose of sulphonylurea or insulin when you take any of them together with Alogliptin Torrent in order to avoid too low blood sugar (hypoglycaemia)
- if you have kidney disease, you can still take this medicine but your doctor may reduce the dose
- if you have liver disease
- if you suffer from heart failure
- if you are taking insulin or an anti-diabetic medicine, your doctor may want to reduce your dose of the other anti-diabetic medicine or insulin when you take either of them together with Alogliptin Torrent in order to avoid low blood sugar
- if you have or have had a disease of the pancreas

Contact your doctor if you encounter blistering of the skin, as it may be a sign for a condition called bullous pemphigoid. Your doctor may ask you to stop alogliptin.

Children and adolescents
Alogliptin Torrent is not recommended for children and adolescents under 18 years due to lack of efficacy in these patients.

Other medicines and Alogliptin Torrent
Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Pregnancy and breast-feeding
If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

There is no experience of using Alogliptin Torrent in pregnant women or during breast-feeding. Alogliptin Torrent should not be used during pregnancy or breast-feeding. Your doctor will help you to decide whether to continue breast-feeding or to continue using Alogliptin Torrent.

Driving and using machines
Alogliptin Torrent is not known to affect your ability to drive and use machines. Taking Alogliptin Torrent in combination with other anti-diabetic medicines called sulphonylureas, insulin or combination therapy with thiazolidinedione plus metformin can cause too low blood sugar levels (hypoglycaemia), which may affect your ability to drive and use machines.

Alogliptin Torrent contains sodium
This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially ‘sodium-free’.

3. How to take Alogliptin Torrent

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

Your doctor will prescribe Alogliptin Torrent together with one or more other medicines to control your blood sugar levels. Your doctor will tell you if you need to change the amount of other medicines you take.

The recommended dose of Alogliptin Torrent is 25 mg once a day.

Patients with kidney disease
If you have kidney disease your doctor may prescribe you a reduced dose. This may be 12.5 mg or 6.25 mg once a day, depending on the severity of your kidney disease.

Patients with liver disease
If you have mildly or moderately reduced liver function, the recommended dose of Alogliptin Torrent is 25 mg once a day. This medicine is not recommended for patients with severely reduced liver function due to the lack of data in these patients.

Swallow your tablet(s) whole with water. You can take this medicine with or without food.

If you take more Alogliptin Torrent than you should
If you take more tablets than you should, or if someone else or a child takes your medicine, contact or go to your nearest emergency centre straight away. Take this leaflet or some tablets with you so that your doctor knows exactly what you have taken.

If you forget to take Alogliptin Torrent
If you forget to take a dose, take it as soon as you remember it. Do not take a double dose to make up for a forgotten dose.

If you stop taking Alogliptin Torrent
Do not stop taking Alogliptin Torrent without consulting your doctor first. Your blood sugar levels may increase when you stop taking Alogliptin Torrent.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

STOP taking Alogliptin Torrent and contact a doctor immediately if you notice any of the following serious side effects:

Not known (frequency cannot be estimated from the available data):
- An allergic reaction. The symptoms may include: a rash, hives, swallowing or breathing problems, swelling of your lips, face, throat or tongue and feeling faint.
- A severe allergic reaction: skin lesions or spots on your skin that can progress to a sore surrounded by pale or red rings, blistering and/or peeling of the skin possibly with symptoms such as itching, fever, overall ill feeling, achy joints, vision problems, burning, painful or itchy eyes and mouth sores (Stevens-Johnson syndrome and Erythema multiforme).
- Severe and persistent pain in the abdomen (stomach area) which might reach through to your back, as well as nausea and vomiting, as it could be a sign of an inflamed pancreas (pancreatitis).

You should also discuss with your doctor if you experience the following side effects:

Common (may affect up to 1 in 10 people):
- Symptoms of low blood sugar (hypoglycaemia) may occur when Alogliptin Torrent is taken in combination with insulin or sulphonylureas (e.g. glipizide, tolbutamide, glibenclamide). Symptoms may include: trembling, sweating, anxiety, blurred vision, tingling lips, paleness, mood change or feeling confused. Your blood sugar could fall below the normal level, but can be increased again by taking sugar. It is recommended that you carry some sugar lumps, sweets, biscuits or sugary fruit juice.
- Cold like symptoms such as sore throat, stuffy or blocked nose.
- Rash
- Itchy skin
- Headache
- Stomach ache
- Diarrhoea
- Indigestion, heartburn.

Not known:
- Liver problems such as nausea or vomiting, stomach pain, unusual or unexplained tiredness, loss of appetite, dark urine or yellowing of your skin or the whites of your eyes.
- Inflammation of the connective tissue within the kidneys (interstitial nephritis).
- Blistering of the skin (bullous pemphigoid).

Reporting of side effects
If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via Yellow Card Scheme at Website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the GooglePlay or Apple App store, By reporting side effects, you can help provide more information on the safety of this medicine.

5. How to store Alogliptin Torrent

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and blister after EXP. The expiry date refers to the last day of that month.

HDPE Bottles:
for pack size 30:
Discard 30 days after first opening of the bottle.
for pack size 90:
Discard 90 days after first opening of the bottle.

This medicinal product does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Alogliptin Torrent contains
- The active substance is alogliptin.

Each 12.5 mg tablet contains alogliptin benzoate equivalent to 12.5 mg alogliptin,
- The other ingredients are: Cellulose, microcrystalline (PH 200), Silica, colloidal anhydrous, Starch, pregelatinized, Copovidone (K-28), Croscarmellose sodium, Magnesium Stearate, Hypromellose/HPMC 2910, Titanium Dioxide (E171), Macrogol, Iron oxide yellow (E172).

What Alogliptin Torrent looks like and contents of the pack

Alogliptin Torrent 12.5 mg film-coated tablets are Yellow colored, oval shaped, biconvex, film-coated tablet with debossed “D2” on one side and plain on other side.

Alogliptin Torrent is available in OPA/Al/PVC/Al blister in cardboard boxs containing 28 tablets and in HDPE (High density polyethylene) bottles with child resistant closure in cardboard boxes containing 30 and 90 tablets.

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer

Torrent Pharma (UK) Ltd.
3rd Floor, Nexus Building
4 Gatwick Road
Crawley
West Sussex
RH10 9BG
United Kingdom

This leaflet was last revised in 09/2025.

8087460-7803