

## **SUMMARY OF PRODUCT CHARACTERISTICS**

### **1 NAME OF THE MEDICINAL PRODUCT**

Bimatoprost/Timolol Zentiva 0.3 mg/ml + 5 mg/ml eye drops, solution

### **2 QUALITATIVE AND QUANTITATIVE COMPOSITION**

One ml of solution contains 0.3 mg of bimatoprost and 5 mg of timolol (as 6.8 mg of timolol maleate).

#### Excipients with known effect

Each ml of solution contains 0.05 mg of benzalkonium chloride.

Each ml of solution contains 0.95 mg of phosphates.

For the full list of excipients, see section 6.1.

### **3 PHARMACEUTICAL FORM**

Eye drops, solution.

Practically clear solution, practically free from particles.  
Colourless to slightly yellow solution.

The pH of solution is 6.5 to 7.8, the osmolality is 260 to 320 mOsmol/kg.

### **4 CLINICAL PARTICULARS**

#### **4.1 Therapeutic indications**

Reduction of intraocular pressure (IOP) in adult patients with open-angle glaucoma or ocular hypertension who are insufficiently responsive to topical beta-blockers or prostaglandin analogues.

#### **4.2 Posology and method of administration**

##### Posology

*Recommended dosage in adults (including older people)*

The recommended dose is one drop of Bimatoprost/Timolol in the affected eye(s) once daily, administered either in the morning or in the evening. It should be administered at the same time each day.

Existing literature data for bimatoprost/timolol containing products suggest that evening dosing may be more effective in IOP lowering than morning dosing. However, consideration should be given to the likelihood of compliance when considering either morning or evening dosing (see section 5.1).

If one dose is missed, treatment should continue with the next dose as planned. The dose should not exceed one drop in the affected eye(s) daily.

#### *Renal and hepatic impairment*

Bimatoprost/timolol has not been studied in patients with hepatic or renal impairment. Therefore caution should be used in treating such patients.

#### *Paediatric population*

The safety and efficacy of bimatoprost/timolol in children aged 0 to 18 years has not been established. No data are available.

#### Method of administration

If more than one topical ophthalmic medicinal product is to be used, each one should be instilled at least 5 minutes apart.

When using nasolacrimal occlusion or closing the eyelids for 2 minutes, the systemic absorption is reduced. This may result in a decrease in systemic side effects and an increase in local activity.

### **4.3 Contraindications**

- Hypersensitivity to the active substances or to any of the excipients listed in section 6.1.
- Reactive airway disease including bronchial asthma or a history of bronchial asthma, severe chronic obstructive pulmonary disease.
- Sinus bradycardia, sick sinus syndrome, sino-atrial block, second or third degree atrioventricular block, not controlled with pace-maker. Overt cardiac failure, cardiogenic shock.

### **4.4 Special warnings and precautions for use**

Like other topically applied ophthalmic medicinal products, the active substances (timolol/ bimatoprost) in Bimatoprost/Timolol may be absorbed systemically. No enhancement of the systemic absorption of the individual active substances has been observed. Due to the beta-adrenergic component, timolol, the same types of

cardiovascular, pulmonary and other adverse reactions as seen with systemic beta-blockers may occur. Incidence of systemic ADRs after topical ophthalmic administration is lower than for systemic administration. To reduce the systemic absorption, see section 4.2.

#### Cardiac disorders

Patients with cardiovascular diseases (e.g. coronary heart disease, Prinzmetal's angina and cardiac failure) and hypotension therapy with beta-blockers should be critically assessed and therapy with other active substances should be considered. Patients with cardiovascular diseases should be watched for signs of deterioration of these diseases and of adverse reactions.

Due to its negative effect on conduction time, beta-blockers should only be given with caution to patients with first degree heart block.

#### Vascular disorders

Patients with severe peripheral circulatory disturbance/disorders (i.e. severe forms of Raynaud's disease or Raynaud's syndrome) should be treated with caution.

#### Respiratory disorders

Respiratory reactions, including death due to bronchospasm in patients with asthma have been reported following administration of some ophthalmic beta-blockers.

Bimatoprost/Timolol should be used with caution, in patients with mild/moderate chronic obstructive pulmonary disease (COPD) and only if the potential benefit outweighs the potential risk.

#### Endocrine disorders

Beta-adrenergic blocking medicinal products should be administered with caution in patients subject to spontaneous hypoglycemia or to patients with labile diabetes as beta-blockers may mask the signs and symptoms of acute hypoglycemia.

Beta-blockers may also mask the signs of hyperthyroidism.

#### Corneal diseases

Ophthalmic  $\beta$ -blockers may induce dryness of eyes. Patients with corneal diseases should be treated with caution.

#### Other beta-blocking agents

The effect on intra-ocular pressure or the known effects of systemic beta-blockade may be potentiated when timolol is given to the patients already receiving a systemic beta-blocking agent. The response of these patients should be closely observed. The use of two topical beta-adrenergic blocking agents is not recommended (see section 4.5).

### Anaphylactic reactions

While taking beta-blockers, patients with a history of atopy or a history of severe anaphylactic reaction to a variety of allergens may be more reactive to repeated challenge with such allergens and unresponsive to the usual dose of adrenaline used to treat anaphylactic reactions.

### Choroidal detachment

Choroidal detachment has been reported with administration of aqueous suppressant therapy (e.g. timolol, acetazolamide) after filtration procedures.

### Surgical anaesthesia

$\beta$ -blocking ophthalmological preparations may block systemic  $\beta$ -agonist effects e.g. of adrenaline. The anaesthesiologist should be informed when the patient is receiving timolol.

### Hepatic

In patients with a history of mild liver disease or abnormal alanine aminotransferase (ALT), aspartate aminotransferase (AST) and/or bilirubin at baseline, bimatoprost had no adverse reactions on liver function over 24 months. There are no known adverse reactions of ocular timolol on liver function.

### Ocular

Before treatment is initiated, patients should be informed of the possibility of eyelash growth, darkening of the eyelid or periocular skin and increased brown iris pigmentation since these have been observed during treatment with bimatoprost and bimatoprost/timolol. Increased iris pigmentation is likely to be permanent, and may lead to differences in appearance between the eyes if only one eye is treated. After discontinuation of bimatoprost/timolol, pigmentation of iris may be permanent. After 12 months treatment with bimatoprost/timolol, the incidence of iris pigmentation was 0.2%. After 12 months treatment with bimatoprost eye drops alone, the incidence was 1.5% and did not increase following 3 years treatment. The pigmentation change is due to increased melanin content in the melanocytes rather than to an increase in the number of melanocytes. The long term effects of increased iridial pigmentation are not known. Iris colour changes seen with ophthalmic administration of bimatoprost may not be noticeable for several months to years. Neither nevi nor freckles of the iris appear to be affected by treatment. Periorbital tissue pigmentation has been reported to be reversible in some patients.

Macular oedema, including cystoid macular oedema, has been reported with bimatoprost/timolol. Therefore, Bimatoprost/Timolol should be used with caution in aphakic patients, in pseudophakic patients with a torn posterior lens capsule, or in patients with known risk factors for macular oedema (e.g. intraocular surgery, retinal vein occlusions, ocular inflammatory disease and diabetic retinopathy).

Bimatoprost/Timolol should be used with caution in patients with active intraocular inflammation (e.g. uveitis) because the inflammation may be exacerbated.

### Skin

There is a potential for hair growth to occur in areas where Bimatoprost/Timolol solution comes repeatedly in contact with the skin surface. Thus, it is important to apply Bimatoprost/Timolol as instructed and avoid it running onto the cheek or other skin areas.

### Excipients

#### *Benzalkonium chloride*

This medicinal product contains 0.15 mg benzalkonium chloride in each 3 ml of solution, which is equivalent to 0.05 mg/ml.

Benzalkonium chloride has been reported to cause eye irritation, symptoms of dry eyes and may affect the tear film and corneal surface. Should be used with caution in dry eye patients and in patients where the cornea may be compromised. Patients should be monitored in case of prolonged use.

Benzalkonium chloride may be absorbed by soft contact lenses and may change the colour of the contact lenses. Contact lenses should be removed before using this medicinal product and may be reinserted after 15 minutes.

#### *Phosphates*

This medicinal product contains 2.85 mg of phosphates in each 3 ml of solution, which is equivalent to 0.95 mg/ml.

### Other conditions

Bimatoprost/timolol has not been studied in patients with inflammatory ocular conditions, neovascular, inflammatory, angle-closure glaucoma, congenital glaucoma or narrow-angle glaucoma.

In studies of bimatoprost 0.3 mg/ml in patients with glaucoma or ocular hypertension, it has been shown that more frequent exposure of the eye to more than 1 dose of bimatoprost daily may decrease the IOP-lowering effect. Patients using Bimatoprost/Timolol with other prostaglandin analogues should be monitored for changes to their intraocular pressure.

## **4.5 Interaction with other medicinal products and other forms of interaction**

No specific interaction studies have been performed with the bimatoprost / timolol fixed combination.

There is a potential for additive effects resulting in hypotension, and/or marked bradycardia when ophthalmic beta-blockers solution is administered concomitantly with oral calcium channel blockers, guanethidine, beta-adrenergic blocking agents, parasympathomimetics, anti-arrhythmics (including amiodarone) and digitalis glycosides.

Potentiated systemic beta-blockade (e.g., decreased heart rate, depression) has been reported during combined treatment with CYP2D6 inhibitors (e.g. quinidine, fluoxetine, paroxetine) and timolol.

Mydriasis resulting from concomitant use of ophthalmic beta-blockers and adrenaline (epinephrine) has been reported occasionally.

## **4.6 Pregnancy and lactation**

### Pregnancy

There are no adequate data from the use of the bimatoprost / timolol fixed combination in pregnant women. Bimatoprost/Timolol should not be used during pregnancy unless clearly necessary. To reduce the systemic absorption, see section 4.2.

#### *Bimatoprost*

No adequate clinical data in exposed pregnancies are available. Animal studies have shown reproductive toxicity at high maternotoxic doses (see section 5.3).

#### *Timolol*

Epidemiological studies have not revealed malformative effects but shown a risk for intra uterine growth retardation when beta-blockers are administered by the oral route. In addition, signs and symptoms of beta-blockade (e.g. bradycardia, hypotension, respiratory distress and hypoglycaemia) have been observed in the neonate when beta-blockers have been administered until delivery. If Bimatoprost/Timolol is administered until delivery, the neonate should be carefully monitored during the first days of life. Animal studies with timolol have shown reproductive toxicity at doses significantly higher than would be used in clinical practice (see section 5.3).

### Breast-feeding

#### *Timolol*

Beta-blockers are excreted in breast milk. However, at therapeutic doses of timolol in eye drops it is not likely that sufficient amounts would be present in breast milk to produce clinical symptoms of beta-blockade in the infant. To reduce the systemic absorption, see section 4.2.

#### *Bimatoprost*

It is not known if bimatoprost is excreted in human breast milk but it is excreted in the milk of the lactating rat. Bimatoprost/Timolol should not be used by breast-feeding women.

## Fertility

There are no data on the effects of bimatoprost/timolol on human fertility.

### **4.7 Effects on ability to drive and use machines**

Bimatoprost/Timolol has negligible influence on the ability to drive and use machines. As with any ocular treatment, if transient blurred vision occurs at instillation, the patient should wait until the vision clears before driving or using machines.

### **4.8 Undesirable effects**

#### *Summary of the safety profile*

The adverse reactions reported in clinical studies using bimatoprost/timolol were limited to those earlier reported for either of the single active substances bimatoprost and timolol. No new adverse reactions specific for bimatoprost/timolol have been observed in clinical studies.

The majority of adverse reactions reported in clinical studies using bimatoprost/timolol were ocular, mild in severity and none were serious. Based on 12-month clinical data, the most commonly reported adverse reaction was conjunctival hyperaemia (mostly trace to mild and thought to be of a non-inflammatory nature) in approximately 26% of patients and led to discontinuation in 1.5% of patients.

#### *Tabulated list of adverse reactions*

Table 1 presents the adverse reactions that have been reported during clinical studies with all bimatoprost/timolol formulations (multi-dose and single-dose) (within each frequency grouping, adverse reactions are presented in order of decreasing seriousness) or in the post-marketing period.

The frequency of possible adverse reactions listed below is defined using the following convention:

Very common	$\geq 1/10$
Common	$\geq 1/100$ to $< 1/10$
Uncommon	$\geq 1/1,000$ to $< 1/100$
Rare	$\geq 1/10,000$ to $< 1/1,000$
Very rare	$< 1/10,000$
Not known	Frequency cannot be estimated from available data

**Table 1**

<b>System Organ Class</b>	<b>Frequency</b>	<b>Adverse reaction</b>
<i>Immune system disorders</i>	Not known	hypersensitivity reactions including signs or symptoms of allergic dermatitis, angioedema, eye allergy
<i>Psychiatric disorders</i>	Not known	insomnia <sup>2</sup> , nightmare <sup>2</sup>
<i>Nervous system disorders</i>	Common	headache
	Not known	dysgeusia <sup>2</sup> , dizziness
<i>Eye disorders</i>	Very common	conjunctival hyperaemia.
	Common	punctuate keratitis, corneal erosion <sup>2</sup> , burning sensation <sup>2</sup> , conjunctival irritation <sup>1</sup> , eye pruritus, stinging sensation in the eye <sup>2</sup> , foreign body sensation, dry eye, erythema of eyelid, eye pain, photophobia, eye discharge, visual disturbance <sup>2</sup> , eyelid pruritus, visual acuity worsened <sup>2</sup> , blepharitis <sup>2</sup> , eyelid oedema, eye irritation, lacrimation increased, growth of eyelashes.
	Uncommon	iritis <sup>2</sup> , conjunctival oedema <sup>2</sup> , eyelid pain <sup>2</sup> , abnormal sensation in the eye <sup>1</sup> , asthenopia, trichiasis <sup>2</sup> , iris hyperpigmentation <sup>2</sup> , periorbital and lid changes associated with periorbital fat atrophy and skin tightness resulting in deepening of eyelid sulcus, eyelid ptosis, enophthalmos, lagophthalmos and eyelid retraction <sup>1,2</sup> , eyelash discolouration (darkening) <sup>1</sup> .
	Not known	cystoid macular oedema <sup>2</sup> , eye swelling, vision blurred <sup>2</sup> ,

		ocular discomfort
<i>Cardiac disorders</i>	Not known	bradycardia
<i>Vascular disorders</i>	Not known	hypertension
<i>Respiratory, thoracic and mediastinal disorders</i>	Common	rhinitis <sup>2</sup>
	Uncommon	dyspnoea
	Not known	bronchospasm (predominantly in patients with pre-existing bronchospastic disease) <sup>2</sup> , asthma
<i>Skin and subcutaneous tissue disorders</i>	Common	blepharal pigmentation <sup>2</sup> , hirsutism <sup>2</sup> , skin hyperpigmentation (periocular).
	Not known	alopecia, skin discolouration (periocular)
<i>General disorders and administration site conditions</i>	Not known	fatigue

<sup>1</sup>adverse reactions only observed with bimatoprost/timolol single-dose formulation

<sup>2</sup>adverse reactions only observed with bimatoprost/timolol multi-dose formulation

Like other topically applied ophthalmic drugs, bimatoprost/timolol is absorbed into the systemic circulation. Absorption of timolol may cause similar undesirable effects as seen with systemic beta-blocking agents. The incidence of systemic ADRs after topical ophthalmic administration is lower than for systemic administration. To reduce the systemic absorption, see section 4.2.

Additional adverse reactions that have been seen with either of the active substances (bimatoprost or timolol), and may potentially occur also with this medicinal product are listed below in Table 2:

**Table 2**

<b>System Organ Class</b>	<b>Adverse reaction</b>
<i>Immune system disorders</i>	systemic allergic reactions including anaphylaxis <sup>1</sup>
<i>Metabolism and nutrition disorders</i>	hypoglycaemia <sup>1</sup>
<i>Psychiatric disorders</i>	depression <sup>1</sup> , memory loss <sup>1</sup> , hallucination <sup>1</sup>
<i>Nervous system disorders</i>	syncope <sup>1</sup> , cerebrovascular accident <sup>1</sup> , increase in signs and symptoms of myasthenia

	gravis <sup>1</sup> , paraesthesia <sup>1</sup> , cerebral ischaemia <sup>1</sup>
<i>Eye disorders</i>	decreased corneal sensitivity <sup>1</sup> , diplopia <sup>1</sup> , ptosis <sup>1</sup> , choroidal detachment following filtration surgery (see section 4.4) <sup>1</sup> , keratitis <sup>1</sup> , blepharospasm <sup>2</sup> , retinal haemorrhage <sup>2</sup> , uveitis <sup>2</sup> ,
<i>Cardiac disorder</i>	atrioventricular block <sup>1</sup> , cardiac arrest <sup>1</sup> , arrhythmia <sup>1</sup> , cardiac failure <sup>1</sup> , congestive heart failure <sup>1</sup> , chest pain <sup>1</sup> , palpitations <sup>1</sup> , oedema <sup>1</sup>
<i>Vascular disorders</i>	hypotension <sup>1</sup> , Raynaud's phenomenon <sup>1</sup> , cold hands and feet <sup>1</sup>
<i>Respiratory, thoracic and mediastinal disorders</i>	asthma exacerbation <sup>2</sup> , COPD exacerbation <sup>2</sup> , cough <sup>1</sup>
<i>Gastrointestinal disorders</i>	nausea <sup>1,2</sup> , diarrhoea <sup>1</sup> , dyspepsia <sup>1</sup> , dry mouth <sup>1</sup> , abdominal pain <sup>1</sup> , vomiting <sup>1</sup>
<i>Skin and subcutaneous tissue disorders</i>	psoriasiform rash <sup>1</sup> or exacerbation of psoriasis <sup>1</sup> , skin rash <sup>1</sup>
<i>Musculoskeletal and connective tissue disorders</i>	myalgia <sup>1</sup>
<i>Reproductive system and breast disorders</i>	sexual dysfunction <sup>1</sup> , decreased libido <sup>1</sup>
<i>General disorders and administration site conditions</i>	asthenia <sup>1,2</sup>
<i>Investigations</i>	liver function tests (LFT) abnormal <sup>2</sup>

<sup>1</sup> adverse reactions observed with timolol

<sup>2</sup> adverse reactions observed with bimatoprost

#### Adverse reactions reported in phosphate containing eye drops

Cases of corneal calcification have been reported very rarely in association with the use of phosphate containing eye drops in some patients with significantly damaged corneas.

### *Reporting of suspected adverse reactions*

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store.

## **4.9 Overdose**

A topical overdose with Bimatoprost/Timolol is not likely to occur or to be associated with toxicity.

### Bimatoprost

If Bimatoprost/Timolol is accidentally ingested, the following information may be useful: in two-week oral rat and mouse studies, doses of bimatoprost up to 100 mg/kg/day did not produce any toxicity. This dose expressed as mg/m<sup>2</sup> is at least 70-times higher than the accidental dose of one bottle of bimatoprost/timolol in a 10 kg child.

### Timolol

Symptoms of systemic timolol overdose include: bradycardia, hypotension, bronchospasm, headache, dizziness, shortness of breath, and cardiac arrest. A study of patients with renal failure showed that timolol did not dialyse readily.

If overdose occurs treatment should be symptomatic and supportive.

## **5 PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: ophthalmological,– beta-blocking agents – ATC code: S01ED51

### Mechanism of action

Bimatoprost/Timolol consists of two active substances: bimatoprost and timolol. These two components decrease elevated intraocular pressure (IOP) by complementary mechanisms of action and the combined effect results in additional

IOP reduction compared to either compound administered alone. Bimatoprost/timolol has a rapid onset of action.

Bimatoprost is a potent ocular hypotensive active substance . It is a synthetic prostamide, structurally related to prostaglandin  $F_{2\alpha}$  ( $PGF_{2\alpha}$ ) that does not act through any known prostaglandin receptors. Bimatoprost selectively mimics the effects of newly discovered biosynthesised substances called prostamides. The prostamide receptor, however, has not yet been structurally identified. The mechanism of action by which bimatoprost reduces intraocular pressure in man is by increasing aqueous humour outflow through the trabecular meshwork and enhancing uveoscleral outflow.

Timolol is a  $\beta_1$  and  $\beta_2$  non-selective adrenergic receptor blocking agent that does not have significant intrinsic sympathomimetic, direct myocardial depressant, or local anaesthetic (membrane-stabilising) activity. Timolol lowers IOP by reducing aqueous humour formation. The precise mechanism of action is not clearly established, but inhibition of the increased cyclic AMP synthesis caused by endogenous beta-adrenergic stimulation is probable.

#### Clinical effects

The IOP-lowering effect of bimatoprost/timolol is non-inferior to that achieved by adjunctive therapy of bimatoprost (once daily) and timolol (twice daily).

Existing literature data for bimatoprost/timolol suggest that evening dosing may be more effective in IOP lowering than morning dosing. However, consideration should be given to the likelihood of compliance when considering either morning or evening dosing.

#### Paediatric population

The safety and efficacy of bimatoprost/timolol in children aged 0 to 18 years has not been established.

## **5.2 Pharmacokinetic properties**

#### Bimatoprost/timolol medicinal product

Plasma bimatoprost and timolol concentrations were determined in a crossover study comparing the monotherapy treatments to bimatoprost/timolol treatment in healthy subjects. Systemic absorption of the individual components was minimal and not affected by co-administration in a single formulation.

In two 12-month studies where systemic absorption was measured, no accumulation was observed with either of the individual components.

### Bimatoprost

Bimatoprost penetrates the human cornea and sclera well *in vitro*. After ocular administration, the systemic exposure of bimatoprost is very low with no accumulation over time. After once daily ocular administration of one drop of 0.03% bimatoprost to both eyes for two weeks, blood concentrations peaked within 10 minutes after dosing and declined to below the lower limit of detection (0.025 ng/ml) within 1.5 hours after dosing. Mean  $C_{max}$  and  $AUC_{0-24hrs}$  values were similar on days 7 and 14 at approximately 0.08 ng/ml and 0.09 ng·hr/ml respectively, indicating that a steady drug concentration was reached during the first week of ocular dosing.

Bimatoprost is moderately distributed into body tissues and the systemic volume of distribution in humans at steady-state was 0.67 l/kg. In human blood, bimatoprost resides mainly in the plasma. The plasma protein binding of bimatoprost is approximately 88%.

Bimatoprost is the major circulating species in the blood once it reaches the systemic circulation following ocular dosing. Bimatoprost then undergoes oxidation, N-deethylation and glucuronidation to form a diverse variety of metabolites.

Bimatoprost is eliminated primarily by renal excretion, up to 67% of an intravenous dose administered to healthy volunteers was excreted in the urine, 25% of the dose was excreted via the faeces. The elimination half-life, determined after intravenous administration, was approximately 45 minutes; the total blood clearance was 1.5 l/hr/kg.

### Characteristics in older people

After twice daily dosing, the mean  $AUC_{0-24hrs}$  value of 0.0634 ng·hr/ml bimatoprost in the elderly (subjects 65 years or older) were significantly higher than 0.0218 ng·hr/ml in young healthy adults. However, this finding is not clinically relevant as systemic exposure for both elderly and young subjects remained very low from ocular dosing. There was no accumulation of bimatoprost in the blood over time and the safety profile was similar in elderly and young patients.

### Timolol

After ocular administration of a 0.5% eye drops solution in humans undergoing cataract surgery, peak timolol concentration was 898 ng/ml in the aqueous humour at one hour post-dose. Part of the dose is absorbed systemically where it is extensively metabolised in the liver. The half-life of timolol in plasma is about 4 to 6 hours. Timolol is partially metabolised by the liver with timolol and its metabolites excreted by the kidney. Timolol is not extensively bound to plasma.

## **5.3 Preclinical safety data**

### Bimatoprost/timolol medicinal product

Repeated dose ocular toxicity studies on bimatoprost/timolol showed no special hazard for humans. The ocular and systemic safety profile of the individual components is well established.

### Bimatoprost

Non-clinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, genotoxicity, carcinogenic potential. Studies in rodents produced species-specific abortion at systemic exposure levels 33- to 97-times that achieved in humans after ocular administration.

Monkeys administered ocular bimatoprost concentrations of  $\square$ 0.03% daily for 1 year had an increase in iris pigmentation and reversible dose-related periocular effects characterised by a prominent upper and/or lower sulcus and widening of the palpebral fissure. The increased iris pigmentation appears to be caused by increased stimulation of melanin production in melanocytes and not by an increase in melanocyte number. No functional or microscopic changes related to the periocular effects have been observed, and the mechanism of action for the periocular changes is unknown.

### Timolol

Non-clinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, repeated dose toxicity, genotoxicity, carcinogenic potential, toxicity to reproduction.

## **6 PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Benzalkonium chloride

Sodium chloride

Disodium phosphate heptahydrate

Citric acid monohydrate

Hydrochloric acid (for pH adjustment)

Sodium hydroxide (for pH adjustment)

Purified water

### **6.2 Incompatibilities**

Not applicable.

### **6.3 Shelf life**

3 years

4 weeks after first opening.

**6.4 Special precautions for storage**

This medicinal product does not require any special storage conditions.

**6.5 Nature and contents of container**

White LDPE bottles with dark blue HDPE screw cap and white LDPE dropper insert. Each bottle has a fill volume of 3 ml.

The following pack sizes are available: cartons containing 1 bottle or 3 bottles.

Not all pack sizes may be marketed.

**6.6 Special precautions for disposal**

No special requirements.

**7 MARKETING AUTHORISATION HOLDER**

Zentiva Pharma UK Limited  
12 New Fetter Lane,  
London,  
EC4A 1JP  
United Kingdom

**8 MARKETING AUTHORISATION NUMBER(S)**

PL 17780/0888

**9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

21/02/2025

**10 DATE OF REVISION OF THE TEXT**

21/02/2025