

# Provigil® 100mg Tablets

## Modafinil 100mg Tablets

(Modafinil)

1829  
15.09.16[11]

### PACKAGE LEAFLET: INFORMATION FOR THE USER

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

This medication is available using any of the above names but will be referred to as Provigil throughout the following leaflet. Provigil Tablets are also available in strength of 200mg.

#### What is in this leaflet:

1. What Provigil is and what it is used for
2. What you need to know before you take Provigil
3. How to take Provigil
4. Possible side effects
5. How to store Provigil
6. Contents of the pack and other information

#### 1. WHAT PROVIGIL IS AND WHAT IT IS USED FOR

The active ingredient in the tablets is modafinil. Modafinil can be taken by adults who suffer from narcolepsy to help them to stay awake. Narcolepsy is a condition that causes excessive daytime sleepiness and a tendency to fall asleep suddenly in inappropriate situations (sleep attacks). Modafinil may improve your narcolepsy and reduce the likelihood that you will have sleep attacks but there may still be other ways that you can improve your condition and your doctor will advise you.

#### 2. WHAT YOU NEED TO KNOW BEFORE YOU TAKE PROVIGIL

##### Do not take Provigil if you:

- Are **allergic** to modafinil, or any of the other ingredients of this medicine (listed in section 6 'What Provigil contains').
- Have an **irregular heartbeat**.
- Have **uncontrolled, moderate to severe high blood pressure** (hypertension).

##### Warnings and precautions

Talk to your doctor or pharmacist before taking Provigil if you:

- Have any **heart problems** or **high blood pressure**. Your doctor will need to check these regularly while you are taking Provigil.
- Have ever had **depression, low mood, anxiety, psychosis** (loss of contact with reality) or **mania** (over-excitement or feeling of extreme happiness) or **bipolar disorder** because Provigil may make your condition worse.
- Have **kidney** or **liver problems** (because you will need to take a lower dose).
- Have had **alcohol** or **drug problems** in the past.

##### Other things to talk to your doctor or pharmacist about

- Some people have reported having **suicidal** or **aggressive thoughts** or **behaviour** while taking this medicine.  
**Tell your doctor straight away** if you notice that you are becoming **depressed, feel aggressive or hostile** towards other people or have **suicidal thoughts** or other changes in your behaviour (see section 4). You may want to consider asking a family member or close friend to help you look out for signs of depression or other changes in your behaviour.
- This medicine has the potential for you to become reliant (dependent) on it after long-term use. If you need to take it for a long time your doctor will check regularly that it is still the best medicine for you.

##### Children and adolescents

Children aged less than 18 years should not take this medicine.

##### Other medicines and Provigil

Tell your doctor or pharmacist if you are taking have recently taken or might take any other medicines, including medicines obtained without prescription. Provigil and certain other medicines can affect each other and your doctor may need to adjust the doses that you are taking. It is especially important if you are taking any of the following medicines as well as Provigil:

- Hormonal **contraceptives** (including the contraceptive pill, implants, intrauterine devices (IUDs) and patches). You will need to consider other birth control methods while taking Provigil, and for two months after stopping treatment, because Provigil reduces their effectiveness.
- **Omeprazole** (for acid reflux, indigestion or ulcers).
- Antiviral medicines to treat HIV infection (protease inhibitors e.g. indinavir or ritonavir).

- **Ciclosporin** (used to prevent organ transplant rejection, or for arthritis or psoriasis).
- Medicines for **epilepsy** (e.g. carbamazepine, phenobarbital or phenytoin).
- Medicines for depression (e.g. amitriptyline, citalopram or fluoxetine) or anxiety (e.g. diazepam).
- Medicines for thinning the blood (e.g. warfarin). Your doctor will monitor your blood clotting times during treatment.
- Calcium channel blockers or betablockers for **high blood pressure** or heart problems (e.g. amlodipine, verapamil or propranolol).
- Statin medicines for lowering **cholesterol** (e.g. atorvastatin or simvastatin).

##### Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

It is not known if your medicine may harm your unborn baby.

##### Driving and using machines

Provigil can cause blurred vision or dizziness in up to 1 in 10 people. If you are affected or you find that while using this medication you still feel very sleepy, do not attempt to drive or operate machinery.

##### Provigil contains Lactose

Provigil contain lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

#### 3. HOW TO TAKE PROVIGIL

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

Tablets should be swallowed whole with water.

##### Adults

The usual dose is 200 mg a day. This can be taken once daily (in the morning) or divided into two doses a day (100 mg in the morning and 100 mg at midday).

Your doctor in some cases may decide to increase your daily dose up to 400 mg.

##### Elderly patients (over 65 years of age)

The usual dose is 100 mg a day.

Your doctor will only increase your dose (up to the maximum 400 mg a day) provided that you do not have any liver or kidney problems.

##### Adults with severe kidney and liver problems

The usual dose is 100 mg a day.

Your doctor will review your treatment regularly to check that it is right for you.

##### If you take more Provigil than you should

If you take too many tablets you may feel sick, restless, disorientated, confused, agitated, anxious or excited.

You may also have difficulty sleeping, diarrhoea, hallucinations (sensing things that are not real), chest pain, a change in the speed of your heart beat or an increase in blood pressure. Contact your nearest hospital casualty department or tell your doctor or pharmacist immediately. Take this leaflet and any remaining tablets with you.

##### If you forget to take Provigil

If you forget to take your medicine take the next dose at the usual time, do not take a double dose to make up for the forgotten one.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

#### 4. POSSIBLE SIDE EFFECTS

Like all medicines, this medicine can cause side effects, although not everybody gets them.

##### Stop taking this medicine and tell your doctor straight away if:

- You have sudden difficulty breathing or wheeziness or your face, mouth or throat begins to swell.
- You notice a skin rash or itching (especially if it affects your whole body). Severe rashes may cause blistering or peeling of the skin, ulcers in your mouth, eyes, nose or genitals. You may also have a high temperature (fever) and abnormal blood test results.
- You feel any change in your mental health and wellbeing. The signs may include:
  - mood swings or abnormal thinking,
  - aggression or hostility,
  - forgetfulness or confusion,
  - feeling of extreme happiness,
  - over-excitement or hyperactivity,
  - anxiety or nervousness,
  - depression, suicidal thoughts or behaviour,
  - agitation or psychosis (a loss of contact with reality which may include delusions or sensing things that are not real), feeling detached or numb, or personality disorder.

Other side effects include the following:

**Very common** side effects (may affect more than 1 in 10 people):

- Headache.

**Common** side effects (may affect up to 1 in 10 people):

- Dizziness.
- Sleepiness, extreme tiredness or difficulty sleeping (insomnia).
- Awareness of your heart beat, which may be faster than normal.
- Chest pain.
- Flushing.
- Dry mouth.
- Loss of appetite, feeling sick, stomach pain, indigestion, diarrhoea or constipation.
- Weakness. Numbness or tingling of the hands or feet ('pins and needles').
- Blurred vision.
- Abnormal blood test results showing how your liver is working (increased liver enzymes).
- Irritability

**Uncommon** side effects (may affect up to 1 in 100 people):

- Back pain, neck pain, muscle pain, muscle weakness, leg cramps, joint pain, twitching or tremor.
- Vertigo (spinning sensation).
- Difficulty moving muscles smoothly or other movement problems, muscle tension, coordination problems.
- Hayfever symptoms including itchy/runny nose or watery eyes.
- Increased cough, asthma or shortness of breath.
- Skin rash, acne or itchy skin.
- Sweating.
- Changes in blood pressure (high or low), abnormal heart trace (ECG), and irregular or unusually slow heart beat.
- Difficulty swallowing, swollen tongue or mouth ulcers.
- Excess wind, reflux (bringing back fluid from the stomach), increased appetite, weight changes, thirst or taste alteration.
- Being sick (vomiting).
- Migraine.
- Speech problems.
- Diabetes with increased blood sugar.
- High blood cholesterol.
- Swollen hands and feet.
- Disrupted sleep or abnormal dreams.
- Loss of sex drive.
- Nose bleed, sore throat or inflamed nasal passages (sinusitis).
- Abnormal vision or dry eyes.
- Abnormal urine or more frequent urination.
- Abnormal periods.
- Abnormal blood test results showing that the numbers of your white blood cells have changed.
- Restlessness with increased body movement.

#### Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at:

[www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard).

By reporting side effects, you can help provide more information on the safety of this medicine.

#### 5. HOW TO STORE PROVIGIL

Keep out of the sight and reach of children.

Do not take this medicine after the expiry date shown on the pack.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

If your tablets become discoloured or show any other signs of deterioration, consult your doctor or pharmacist who will tell you what to do.

#### 6. CONTENTS OF THE PACK AND OTHER INFORMATION

##### What Provigil contains

Each tablet contains 100 mg of modafinil as the active ingredient. In addition the tablets also contain: Lactose monohydrate, pregelatinised starch (maize), microcrystalline cellulose, croscarmellose sodium, povidone and magnesium stearate

##### What Provigil looks like and the contents of the pack

Provigil is available as white to off-white, capsule-shaped tablets debossed with '100' on one side. They are packed in blister strips and supplied in boxes of 30 and 90 tablets.

#### Manufacturer and Product Licence holder

Manufactured by:

Laboratoires Macors, Rue des Caillottes, ZI Plaine des Isles, 89000 Auxerre, France.

Cephalon France, 20 rue Charles Martigny, 94700 Maisons-Alfort, France.

Procured from within the EU by Product Licence holder Star Pharmaceuticals Ltd, 5 Sandridge Close, Harrow, Middlesex, HA1 1XD. Repackaged by Servipharm Ltd.

POM

PL No: 20636/1829

Leaflet revision and issue date (Ref) 15.09.16[11]

Provigil is a registered trademark of Cephalon INC.