

Package leaflet: Information for the user

SEREVENT® EVOHALER®
(salmeterol xinafoate)

Your medicine is marketed using above name but will be referred to as Serevent Evohaler or Serevent throughout this leaflet.

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

In this leaflet:

1. What Serevent Evohaler is and what it is used for
2. What you need to know before you use Serevent Evohaler
3. How to use Serevent Evohaler
4. Possible side effects
5. How to store Serevent Evohaler
6. Contents of the pack and other information

1. WHAT SEREVENT EVOHALER IS AND WHAT IT IS USED FOR

- Serevent Evohaler contains the medicine salmeterol. It is a 'long-acting bronchodilator'. It helps the airways in the lungs to stay open. This makes it easier for air to get in and out. The effects are usually felt within 10 to 20 minutes and last for 12 hours or more.
- The doctor has prescribed it to help prevent breathing problems. These could be caused by asthma. Taking Serevent Evohaler regularly will help prevent asthma attacks. This also includes asthma brought on by exercise or at night.
- Taking Serevent Evohaler regularly will also help prevent breathing problems caused by other chest illnesses such as Chronic Obstructive Pulmonary Disease (COPD).
- Serevent Evohaler helps to stop breathlessness and wheezing coming on. It does not work once you are breathless or wheezy. If that happens, you need to use a fast-acting 'reliever' medicine, such as salbutamol.
- Serevent Evohaler is supplied to you in an inhaler. You breathe the medicine directly into your lungs.
- Serevent Evohaler contains norflurane. This is less harmful to the environment than older inhalers. Older inhalers may taste differently to Serevent Evohaler. This will make no difference to how your medicine works.

If you are being treated for asthma, you should always be given both a Serevent and a steroid inhaler to use together.

2. WHAT YOU NEED TO KNOW BEFORE YOU USE SEREVENT EVOHALER

Do not take Serevent Evohaler if:

you are allergic (hypersensitive) to salmeterol or to the other ingredient norflurane (HFA 134a).

Take special care with Serevent Evohaler

- **If your asthma or breathing gets worse tell your doctor straight away.** You may find that you feel more wheezy, your chest feels tight more often or you may need to use more of your fast-acting 'reliever' medicine. If any of these happen, do not increase your number of puffs of Serevent. Your chest condition may be getting worse and you could become seriously ill. See your doctor as you may need a change in asthma treatment.
- Once your asthma is well controlled your doctor may consider it appropriate to gradually reduce the dose of Serevent.
- If you have been prescribed Serevent for your asthma, continue to use any other asthma medication you are already taking. These could include a steroid inhaler or steroid tablets. Continue taking the same doses as before, unless your doctor tells you otherwise. Do this even if you feel much better. **Do not stop taking your steroid inhaler (or any steroid tablets) when you start using Serevent.**
- Your doctor may want to check your health regularly if you have an overactive thyroid gland, diabetes mellitus (Serevent may increase your blood sugar) or heart disease, including an irregular or fast heartbeat.

Other medicines and Serevent

- Tell your doctor if you are taking or have recently taken any other medicines. This includes those for asthma or any other medicines obtained without a prescription. This is because Serevent may not be suitable to be taken with other medicines.
- Inform your doctor before using Serevent if you are currently being treated for any fungal infections with medicines containing ketoconazole or itraconazole, or if you are being treated for HIV with ritonavir. These medicines may increase the risk of you experiencing side effects with Serevent, including irregular heart beats, or may make side-effects worse.

- Beta-blockers should be avoided when taking Serevent, unless your doctor tells you to take them. Beta-blockers, including atenolol, propranolol and sotalol, are mostly used for high blood pressure or other heart conditions. Tell your doctor if you are taking beta-blockers or have recently been prescribed beta-blockers as they may reduce or abolish the effects of salmeterol.
- Serevent can reduce the amount of potassium in your blood. If this happens you may notice an uneven heartbeat, muscle weakness or cramp. This is more likely to happen if you take Serevent with some medicines used to treat high blood pressure (diuretics) and other medicines used to treat breathing problems such as theophylline or steroids. Your doctor may ask for you to have blood tests to check the amount of potassium in your blood. If you have any concerns discuss them with your doctor.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

Driving and using machines

The possible side effects associated with Serevent are unlikely to affect your ability to drive or use machines.

3. HOW TO USE SEREVENT EVOHALER

Always use this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

- **If you are being treated for asthma, you should always be given both a Serevent and a steroid inhaler to use together.**
- Use Serevent every day, until your doctor advises you to stop.
- You will start to feel your medicine working within the first day of use.

Serevent is for inhalation by mouth only.

Adults and adolescents aged 12 years and older with Asthma

- The usual starting dose is 2 puffs twice a day.
- For people with more severe asthma, your doctor may increase your dose to 4 puffs twice a day.

Children with Asthma

- In children aged 4 to 12 the usual dose is 2 puffs twice a day.
- Serevent is not recommended for use in children below 4 years of age.

Adults with Chronic Obstructive Pulmonary Disease (COPD) including bronchitis and emphysema

- The usual starting dose is 2 puffs twice a day.
- Not applicable for children and adolescents.


Instructions for use

Your doctor, nurse or pharmacist should show you how to use your inhaler. They should check how you use it from time to time. Not using the inhaler properly or as prescribed, may mean that the medicine will not help your asthma or COPD as it should.

The medicine is contained in a pressurised canister in a plastic casing with a mouthpiece.

Testing your inhaler

- 1 When using your inhaler for the first time, test that it is working. Remove the mouthpiece cover by gently squeezing the sides with your thumb and forefinger and pull apart.
- 2 To make sure that it works, shake it well, point the mouthpiece away from you and press the canister to release two puffs into the air. If you have not used the inhaler for a week or more, also release two puffs of medicine into the air.



Using your inhaler

It is important to start to breathe in as slowly as possible just before using your inhaler.

- 1 Stand or sit upright when using your inhaler.
- 2 Remove the mouthpiece cover (as shown in the first picture). Check inside and outside to make sure that the mouthpiece is clean and free of objects.



- 3 Shake the inhaler 4 or 5 times to ensure that any loose objects are removed and that the contents of the inhaler are evenly mixed.

- 4 Hold the inhaler upright with your thumb on the base, below the mouthpiece. Breathe out as far as is comfortable.



5 Place the mouthpiece in your mouth between your teeth. Close your lips around it. Do not bite.

6 Breathe in through your mouth. Just after starting to breathe in, press down on the top of the canister to release a puff of medicine. Do this while still breathing in steadily and deeply.



7 Hold your breath, take the inhaler from your mouth and your finger from the top of the inhaler. Continue holding your breath for a few seconds, or as long as is comfortable.

8 Wait about half a minute between taking each puff of medicine and then repeat steps 3 to 7.

9 After use always replace the mouthpiece cover immediately to keep out dust. Replace the cover by firmly pushing and snapping the cap into position.

Practice in front of a mirror for the first few times. If you see a 'mist' coming from the top of your inhaler or the sides of your mouth you should start again

If you or your child find it difficult to use Serevent Evohaler, it may be used with other devices to make its use easier e.g. a Haleraid® or Volumatic® spacer device. Serevent is also available in an alternative device. Talk to your doctor, nurse or pharmacist for further advice.

Cleaning your inhaler

To stop your inhaler blocking up, it is important to clean it at least once a week.

To clean your inhaler:

- Remove the mouthpiece cover.
- Do not remove the metal canister from the plastic casing at any time.
- Wipe the inside and outside of the mouthpiece and the plastic casing with a dry cloth or tissue.
- Replace the mouthpiece cover.

Do not put the metal canister in water.

If you use more Serevent Evohaler than you should

It is important to use the inhaler as instructed. If you accidentally take a larger dose than recommended, talk to your doctor or pharmacist. You may notice your heart beating faster than usual and that you feel shaky and/or dizzy. You may also have a headache, muscle weakness and aching joints.

If you forget to use Serevent Evohaler

Do not take a double dose to make up for a missed dose. Just take your next dose at the usual time.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. POSSIBLE SIDE EFFECTS

Like all medicines, this medicine can cause side effects, although not everybody gets them. To reduce the chances of side effects, your doctor will prescribe the lowest dose of Serevent to control your asthma or COPD. These are the side effects reported by people taking Serevent.

Allergic reactions: you may notice your breathing suddenly gets worse after using Serevent. You may be very wheezy and cough. You may also notice itching and swelling (usually of the face, lips, tongue or throat). If you get these effects or they happen suddenly after using Serevent, **tell your doctor straight away.** Allergic reactions to Serevent are very rare (they affect less than 1 person in 10,000).

Other side effects are listed below:

Common (affects less than 1 person in 10):

- Muscle cramps
- Feeling shaky; fast or uneven heartbeat (palpitations), headache, shaking hands (tremor). Tremors are more likely if you are taking more than two puffs twice daily. These side effects do not last long and happen less as treatment with Serevent continues

Uncommon (affects less than 1 person in 100):

- Rash
- Very fast heart rate (tachycardia). This is more likely to happen if you are taking more than two puffs twice daily
- Feeling nervous.

Rare (affects less than 1 person in 1,000):

- Feeling dizzy
- Being unable to sleep or finding sleep difficult
- A reduction in the amount of potassium in your blood (you may get an uneven heartbeat, muscle weakness, cramp).

Very rare (affects less than 1 person in 10,000):

- **Breathing difficulties or wheezing that gets worse straight after taking Serevent.** If this happens **stop using your Serevent Evohaler.** Use your fast-acting 'reliever' inhaler to help your breathing and **tell your doctor straight away.**
- Uneven heartbeat or your heart gives an extra beat (arrhythmias). If this happens do not stop using Serevent but tell your doctor
- Increases in the amount of sugar (glucose) in your blood (hyperglycaemia). If you have diabetes, more frequent blood sugar monitoring and possibly adjustment of your usual diabetic treatment may be required
- Sore mouth or throat
- Feeling sick (nausea)
- Aching, swollen joints or chest pain

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the **Google Play** or **Apple App Store**. By reporting side effects, you can help provide more information on the safety of this medicine.

5. HOW TO STORE SEREVENT EVOHALER

- **KEEP OUT OF THE SIGHT AND REACH OF CHILDREN.**
- Straight after use, replace the mouthpiece cover firmly and click it into position. Do not use excessive force.
- Do not store above 30°C.
- Protect from frost and heat. As with most inhaled medications in aerosol canisters, the therapeutic effect of this medication may decrease when the canister is cold.
- This is a pressurised container. Do not pierce, break or burn it even when apparently empty.
- Do not use Serevent after the expiry date stated on the label and carton. The expiry date refers to the last day of the month.
- If the inhaler shows any other signs of deterioration, you should seek the advice of your pharmacist who will tell you what to do.
- Medicines should not be disposed of via household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

6. CONTENT OF THE PACK AND OTHER INFORMATION

What Serevent Evohaler contains

- Each actuation (puff) provides 25 micrograms of the active ingredient salmeterol (as xinafoate).
- It also includes a CFC-free propellant Norflurane.
- There are 120 metered actuations (puffs) in each canister.

What Serevent Evohaler looks like and contents of the pack

Pressurised aluminium canister closed with a metering valve containing a pressurised inhalation suspension inserted into a green plastic actuator and mouthpiece. The mouthpiece has a light green plastic dust cap.

Product Licence holder

Procured from within the EU and repackaged by the Product Licence holder: S&M Medical Ltd, Chemilines House, Alperton Lane, Wembley, HA0 1DX.

Manufacturer

This product is manufactured by Glaxo Wellcome Production, Rue Lavoisier 23, 27000 Evreux France.

POM PL: 19488/1188

Leaflet revision date: 29 July 2019

Blind or partially sighted? Is this leaflet hard to see or read? Call 02087997607 to obtain the leaflet in large print, tape, CD or Braille.

Serevent® and Evohaler® are registered trademarks of Glaxo Group Limited, UK.

S1188 LEAFLET Serevent 20190729

Package leaflet: Information for the user

SALMETEROL 25 micrograms EVOHALER

(salmeterol xinafoate)

Your medicine is marketed using above name but will be referred to as Salmeterol Evohaler or Salmeterol throughout this leaflet.

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

In this leaflet:

1. What Salmeterol Evohaler is and what it is used for
2. What you need to know before you use Salmeterol Evohaler
3. How to use Salmeterol Evohaler
4. Possible side effects
5. How to store Salmeterol Evohaler
6. Contents of the pack and other information

1. WHAT SALMETEROL EVOHALER IS AND WHAT IT IS USED FOR

- Salmeterol Evohaler contains the medicine salmeterol. It is a 'long-acting bronchodilator'. It helps the airways in the lungs to stay open. This makes it easier for air to get in and out. The effects are usually felt within 10 to 20 minutes and last for 12 hours or more.
- The doctor has prescribed it to help prevent breathing problems. These could be caused by asthma. Taking Salmeterol Evohaler regularly will help prevent asthma attacks. This also includes asthma brought on by exercise or at night.
- Taking Salmeterol Evohaler regularly will also help prevent breathing problems caused by other chest illnesses such as Chronic Obstructive Pulmonary Disease (COPD).
- Salmeterol Evohaler helps to stop breathlessness and wheezing coming on. It does not work once you are breathless or wheezy. If that happens, you need to use a fast-acting 'reliever' medicine, such as salbutamol.
- Salmeterol Evohaler is supplied to you in an inhaler. You breathe the medicine directly into your lungs.
- Salmeterol Evohaler contains norflurane. This is less harmful to the environment than older inhalers. Older inhalers may taste differently to Salmeterol Evohaler. This will make no difference to how your medicine works.

If you are being treated for asthma, you should always be given both a Salmeterol and a steroid inhaler to use together.

2. WHAT YOU NEED TO KNOW BEFORE YOU USE SALMETEROL EVOHALER

Do not take Salmeterol Evohaler if:

you are allergic (hypersensitive) to salmeterol or to the other ingredient norflurane (HFA 134a).

Take special care with Salmeterol Evohaler

- **If your asthma or breathing gets worse tell your doctor straight away.** You may find that you feel more wheezy, your chest feels tight more often or you may need to use more of your fast-acting 'reliever' medicine. If any of these happen, do not increase your number of puffs of Salmeterol. Your chest condition may be getting worse and you could become seriously ill. See your doctor as you may need a change in asthma treatment.
- Once your asthma is well controlled your doctor may consider it appropriate to gradually reduce the dose of Salmeterol.
- If you have been prescribed Salmeterol for your asthma, continue to use any other asthma medication you are already taking. These could include a steroid inhaler or steroid tablets. Continue taking the same doses as before, unless your doctor tells you otherwise. Do this even if you feel much better. **Do not stop taking your steroid inhaler (or any steroid tablets) when you start using Salmeterol.**
- Your doctor may want to check your health regularly if you have an overactive thyroid gland, diabetes mellitus (Salmeterol may increase your blood sugar) or heart disease, including an irregular or fast heartbeat.

Other medicines and Salmeterol

- Tell your doctor if you are taking or have recently taken any other medicines. This includes those for asthma or any other medicines obtained without a prescription. This is because Salmeterol may not be suitable to be taken with other medicines.
- Inform your doctor before using Salmeterol if you are currently being treated for any fungal infections with medicines containing ketoconazole or itraconazole, or if you are being treated for HIV with zidovudine. These medicines may increase the risk of you experiencing side effects with Salmeterol, including irregular heart beats, or may make side-effects worse.

- Beta-blockers should be avoided when taking Salmeterol, unless your doctor tells you to take them. Beta-blockers, including atenolol, propranolol and sotalol, are mostly used for high blood pressure or other heart conditions. Tell your doctor if you are taking beta-blockers or have recently been prescribed beta-blockers as they may reduce or abolish the effects of salmeterol.
- Salmeterol can reduce the amount of potassium in your blood. If this happens you may notice an uneven heartbeat, muscle weakness or cramp. This is more likely to happen if you take Salmeterol with some medicines used to treat high blood pressure (diuretics) and other medicines used to treat breathing problems such as theophylline or steroids. Your doctor may ask for you to have blood tests to check the amount of potassium in your blood. If you have any concerns discuss them with your doctor.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

Driving and using machines

The possible side effects associated with Salmeterol are unlikely to affect your ability to drive or use machines.

3. HOW TO USE SALMETEROL EVOHALER

Always use this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

- **If you are being treated for asthma, you should always be given both a Salmeterol and a steroid inhaler to use together.**
- Use Salmeterol every day, until your doctor advises you to stop.
- You will start to feel your medicine working within the first day of use.

Salmeterol is for inhalation by mouth only.

Adults and adolescents aged 12 years and older with Asthma

- The usual starting dose is 2 puffs twice a day.
- For people with more severe asthma, your doctor may increase your dose to 4 puffs twice a day.

Children with Asthma

- In children aged 4 to 12 the usual dose is 2 puffs twice a day.
- Salmeterol is not recommended for use in children below 4 years of age.

Adults with Chronic Obstructive Pulmonary Disease (COPD) including bronchitis and emphysema

- The usual starting dose is 2 puffs twice a day.
- Not applicable for children and adolescents.


Instructions for use

Your doctor, nurse or pharmacist should show you how to use your inhaler. They should check how you use it from time to time. Not using the inhaler properly or as prescribed, may mean that the medicine will not help your asthma or COPD as it should.

The medicine is contained in a pressurised canister in a plastic casing with a mouthpiece.

Testing your inhaler

- 1 When using your inhaler for the first time, test that it is working. Remove the mouthpiece cover by gently squeezing the sides with your thumb and forefinger and pull apart.
- 2 To make sure that it works, shake it well, point the mouthpiece away from you and press the canister to release two puffs into the air. If you have not used the inhaler for a week or more, also release two puffs of medicine into the air.



Using your inhaler

It is important to start to breathe in as slowly as possible just before using your inhaler.

- 1 Stand or sit upright when using your inhaler.
- 2 Remove the mouthpiece cover (as shown in the first picture). Check inside and outside to make sure that the mouthpiece is clean and free of objects.



- 3 Shake the inhaler 4 or 5 times to ensure that any loose objects are removed and that the contents of the inhaler are evenly mixed.

- 4 Hold the inhaler upright with your thumb on the base, below the mouthpiece. Breathe out as far as is comfortable.



5 Place the mouthpiece in your mouth between your teeth. Close your lips around it. Do not bite.

6 Breathe in through your mouth. Just after starting to breathe in, press down on the top of the canister to release a puff of medicine. Do this while still breathing in steadily and deeply.



7 Hold your breath, take the inhaler from your mouth and your finger from the top of the inhaler. Continue holding your breath for a few seconds, or as long as is comfortable.

8 Wait about half a minute between taking each puff of medicine and then repeat steps 3 to 7.

9 After use always replace the mouthpiece cover immediately to keep out dust. Replace the cover by firmly pushing and snapping the cap into position.

Practice in front of a mirror for the first few times. If you see a 'mist' coming from the top of your inhaler or the sides of your mouth you should start again

If you or your child find it difficult to use Salmeterol Evohaler, it may be used with other devices to make its use easier e.g. a Haleraid® or Volumatic® spacer device. Salmeterol is also available in an alternative device. Talk to your doctor, nurse or pharmacist for further advice.

Cleaning your inhaler

To stop your inhaler blocking up, it is important to clean it at least once a week.

To clean your inhaler:

- Remove the mouthpiece cover.
- Do not remove the metal canister from the plastic casing at any time.
- Wipe the inside and outside of the mouthpiece and the plastic casing with a dry cloth or tissue.
- Replace the mouthpiece cover.

Do not put the metal canister in water.

If you use more Salmeterol Evohaler than you should

It is important to use the inhaler as instructed. If you accidentally take a larger dose than recommended, talk to your doctor or pharmacist. You may notice your heart beating faster than usual and that you feel shaky and/or dizzy. You may also have a headache, muscle weakness and aching joints.

If you forget to use Salmeterol Evohaler

Do not take a double dose to make up for a missed dose. Just take your next dose at the usual time.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. POSSIBLE SIDE EFFECTS

Like all medicines, this medicine can cause side effects, although not everybody gets them. To reduce the chances of side effects, your doctor will prescribe the lowest dose of Salmeterol to control your asthma or COPD. These are the side effects reported by people taking Salmeterol.

Allergic reactions: you may notice your breathing suddenly gets worse after using Salmeterol. You may be very wheezy and cough. You may also notice itching and swelling (usually of the face, lips, tongue or throat). If you get these effects or they happen suddenly after using Salmeterol, **tell your doctor straight away.** Allergic reactions to Salmeterol are very rare (they affect less than 1 person in 10,000).

Other side effects are listed below:

Common (affects less than 1 person in 10):

- Muscle cramps
- Feeling shaky; fast or uneven heartbeat (palpitations), headache, shaking hands (tremor). Tremors are more likely if you are taking more than two puffs twice daily. These side effects do not last long and happen less as treatment with Salmeterol continues

Uncommon (affects less than 1 person in 100):

- Rash
- Very fast heart rate (tachycardia). This is more likely to happen if you are taking more than two puffs twice daily
- Feeling nervous

Rare (affects less than 1 person in 1,000):

- Feeling dizzy
- Being unable to sleep or finding sleep difficult
- A reduction in the amount of potassium in your blood (you may get an uneven heartbeat, muscle weakness, cramp)

Very rare (affects less than 1 person in 10,000):

- **Breathing difficulties or wheezing that gets worse straight after taking Salmeterol.** If this happens **stop using your Salmeterol Evohaler.** Use your fast-acting 'reliever' inhaler to help your breathing and **tell your doctor straight away.**
- Uneven heartbeat or your heart gives an extra beat (arrhythmias). If this happens do not stop using Salmeterol but tell your doctor
- Increases in the amount of sugar (glucose) in your blood (hyperglycaemia). If you have diabetes, more frequent sugar monitoring and possibly adjustment of your usual diabetic treatment may be required
- Sore mouth or throat
- Feeling sick (nausea)
- Aching, swollen joints or chest pain

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the **Google Play** or **Apple App Store**. By reporting side effects, you can help provide more information on the safety of this medicine.

5. HOW TO STORE SALMETEROL EVOHALER

- **KEEP OUT OF THE SIGHT AND REACH OF CHILDREN.**
- Straight after use, replace the mouthpiece cover firmly and click it into position. Do not use excessive force.
- Do not store above 30°C.
- Protect from frost and heat. As with most inhaled medications in aerosol canisters, the therapeutic effect of this medication may decrease when the canister is cold.
- This is a pressurised container. Do not pierce, break or burn it even when apparently empty.
- Do not use Salmeterol after the expiry date stated on the label and carton. The expiry date refers to the last day of the month.
- If the inhaler shows any other signs of deterioration, you should seek the advice of your pharmacist who will tell you what to do.
- Medicines should not be disposed of via household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

6. CONTENT OF THE PACK AND OTHER INFORMATION

What Salmeterol Evohaler contains

- Each actuation (puff) provides 25 micrograms of the active ingredient salmeterol (as xinafoate).
- It also includes a CFC-free propellant Norflurane.
- There are 120 metered actuations (puffs) in each canister.

What Salmeterol Evohaler looks like and contents of the pack

Pressurised aluminium canister closed with a metering valve containing a pressurised inhalation suspension inserted into a green plastic actuator and mouthpiece. The mouthpiece has a light green plastic dust cap.

Product Licence holder

Procured from within the EU and repackaged by the Product Licence holder: S&M Medical Ltd, Chemilines House, Alperton Lane, Wembley, HA0 1DX.

Manufacturer

This product is manufactured by Glaxo Wellcome Production, Rue Lavoisier 23, 27000 Evreux France.

POM PL: 19488/1188

Leaflet revision date: 29 July 2019

Blind or partially sighted? Is this leaflet hard to see or read? Call 02087997607 to obtain the leaflet in large print, tape, CD or Braille.

S1188 LEAFLET Salmeterol 20190729