

Rare side effects (may affect up to 1 in 1 000 people)

Low platelet count (thrombocytopenia), which increases risk of bleeding or bruising (small purple-red marks in skin or other tissue caused by bleeding), high blood calcium level, high blood sugar level, headache, abdominal discomfort, yellowing of the skin or eyes (jaundice), excess of biliary substances in the blood (cholestasis), photosensitivity reaction, uncontrolled blood levels of glucose in patients with a diagnosis of diabetes mellitus, sugars in the urine (glucosuria).

Very rare side effects (may affect up to 1 in 10 000 people)

Abnormal breakdown of red blood cells (haemolytic anaemia), inability of the bone marrow to work properly, reduction of white blood cells (leukopenia, agranulocytosis), serious allergic reactions (e.g. hypersensitivity), increased pH due to low blood chloride level (disturbed acid-base balance, alkalosis hypochloraemic), acute respiratory distress (signs include severe shortness of breath, fever, weakness, and confusion), inflammation of the pancreas, lupus-like syndrome (a condition mimicking a disease called systemic lupus erythematosus where the body's immune system attacks the body), inflammation of blood vessels (vasculitis necrotising).

Not known (frequency cannot be estimated from the available data)

Skin and lip cancer (non-melanoma skin cancer), blood cell deficiency (aplastic anaemia), decrease in vision and eye pain (possible signs of fluid accumulation in the vascular layer of the eye (choroidal effusion) or acute-angle closure glaucoma), skin disorders such as inflamed blood vessels in the skin, increased sensitivity to sunlight, rash, redness of the skin, blistering of the lips, eyes or mouth, skin peeling, fever (possible signs of erythema multiforme), weakness, kidney impairment.

Low levels of sodium accompanied by symptoms relating to the brain or nerves (feeling sick, progressive disorientation, lack of interest or energy) occurs in isolated cases.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly (see details below). By reporting side effects you can help provide more information on the safety of this medicine.

Yellow Card Scheme

Website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store

5. How to store MicardisPlus

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton after "EXP". The expiry date refers to the last day of that month.

This medicine does not require any special temperature storage conditions. Store in the original package in order to protect from moisture. Remove your MicardisPlus tablet from the sealed blister only directly prior to intake.

Occasionally, the outer layer of the blister pack separates from the inner layer between the blister pockets. You do not need to take any action if this happens.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

If the medicine becomes discolored or shows any other signs of deterioration, you should seek the advice of your pharmacist who will tell you what to do.

6. Contents of the pack and other information**What MicardisPlus contains**

- The active substances are telmisartan and hydrochlorothiazide. Each tablet contains 80 mg telmisartan and 12.5 mg hydrochlorothiazide.
- The other ingredients are lactose monohydrate, magnesium stearate, maize starch, meglumine, microcrystalline cellulose, povidone K25, red iron oxide (E172), sodium hydroxide, sodium starch glycollate (type A), sorbitol (E420).

What MicardisPlus looks like and contents of the pack

MicardisPlus 80 mg/12.5 mg tablets are red and white, oblong-shaped, two-layer tablets engraved with the company logo and the code 'H8'.

MicardisPlus is available in blister packs containing 28 tablets.

Not all pack sizes may be available in your country.

Manufacturer

Boehringer Ingelheim Hellas Single Member S.A.
5th km Paiania – Markopoulo
Koropi Attiki, 19441
Greece

Or

Rottendorf Pharma GmbH
Ostenfelder Strasse 51 - 61
59320 Ennigerloh
Germany

Or

Boehringer Ingelheim France
100-104 Avenue de France
75013 Paris
France

Procured from within the EU and repackaged for the PL Holder: Abacus Medicine Ltd., Abbey House, 282 Farnborough Road, Farnborough, GU14 7NA, United Kingdom

PL 45763/1677

POM

MicardisPlus® is a registered trademark of Boehringer Ingelheim Pharma GmbH & Co. KG

For any information about this medicine, please contact the local representative of the Product Licence Holder:

Abacus Medicine Ltd.

tel: +44(0)2036301244

email: MedInfo@abacusmedicine.co.uk

or

drugsafetyuk@abacusmedicine.co.uk

This leaflet was last revised on 18.07.2025.

**Blind or partially sighted?
Is this leaflet hard to see or read?
Call 02036301310 or write to
info@abacusmedicine.com to
obtain the leaflet in a format
suitable for you.**