



CanesOasis®



Read all of this leaflet carefully before you take this medicine because it contains important information you need to know.

This medicine contains:

These sachets are used for the relief of the symptoms of cystitis in women. Cystitis is an inflammation of the bladder, which causes painful irritation and an unpleasant burning sensation when passing water.

Do not take this medicine if you:

- Talk to your doctor if you have:**

Talk to your doctor or pharmacist if you are taking:

- lithium for manic depression — — —
- methenamine for urinary infections
- other medicines as these may be affected by taking this medicine.

- have an intolerance to some sugars.

Pregnancy and breastfeeding:

This medicine is not recommended if you are pregnant or breastfeeding. Please see your doctor or pharmacist before taking this medicine.

Contains sucrose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.



3. How to take this medicine

This is a 2-day course of treatment.

This medicine is not to be used by men or children.

— If symptoms persist or worsen after the 2-day course is completed, consult your doctor or pharmacist. Do not repeat the treatment without medical advice.

If you take too many sachets by mistake, contact your doctor or pharmacist straight away.

Most people do not have any side effects while taking this medicine.

- However, if you experience any side effects, or anything unusual happens, stop taking the medicine immediately, and see your doctor or pharmacist.

- Passing water more frequently.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet.

By reporting side effects you can help provide
more information on the safety of this medicine.

Keep out of the sight and reach of children.

Do not use this medicine after the expiry date printed on the pack.
Store below 25°C, in a dry place.



Each 5.6g sachet of granules contains the active ingredient: sodium citrate dihydrate 4.0g.

The other ingredients are: sucrose (sugar) 1.5g, colloidal anhydrous silica, saccharin sodium and cranberry flavouring.

This product is available in a pack size of 6 sachets.

The Marketing Authorisation holder and manufacturer is Wrafton Laboratories Limited, Braunton, Devon, EX33 2DL, UK.

Need Advice? ☎ Call 0845 758 5030
(Calls charged at local rate)

Distributed by: Bayer plc, 400 South Oak Way,
Reading, RG2 6AD, UK.

Text revised: December 2020

F4T91LVJ2

Bayer