

## **Package leaflet: Information for the user**

### **Azelair 0.15% Nasal Spray**

Azelastine hydrochloride

**Read all of this leaflet carefully before you start using this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.
- In this leaflet Azelair 0.15% Nasal Spray is called Azelair.

#### **What is in this leaflet:**

1. What Azelair is and what it is used for
2. What you need to know before you use Azelair
3. How to use Azelair
4. Possible side effects
5. How to store Azelair
6. Contents of the pack and other information

#### **1. What Azelair is and what it is used for**

Azelair contains azelastine hydrochloride which belongs to a group of medicines called antihistamines. Antihistamines work by preventing the effects of histamine that the body produces as part of an allergic reaction.

Azelair is used to treat allergic rhinitis in adults, adolescents and children 6 years and older. This is an allergic reaction to substances such as, for example, pollen, house dust mites or animal hair.

Usually, it affects you by causing a runny nose, sneezing, itching or blocked nose. Azelair should help control these symptoms.

#### **2. What you need to know before you use Azelair**

##### **Do not use Azelair**

- if you are allergic (hypersensitive) to azelastine hydrochloride or any of the other ingredients of this medicine (listed in section 6).

##### **Warnings and precautions**

Talk to your doctor or pharmacist before using Azelair.

##### **Children**

Azelair is not recommended for use in children below 6 years of age.

##### **Other medicines and Azelair**

Even if Azelair is not known to be affected by other medicines, tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

### **Pregnancy and breast-feeding**

There is only limited information on the effects of Azelair on the unborn or breast fed child.

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before using this medicine.

### **Driving and using machines**

Azelair has minor influence on the ability to drive and use machines.

Rarely, you may experience fatigue or dizziness due to the disease itself or when using Azelair. In these cases, do not drive or operate machinery. Please be aware that drinking alcohol may enhance these effects.

## **3. How to use Azelair**

Always use this medicine exactly as your doctor or pharmacist has told you. Azelair is for nasal use.

### **Important:**

**If you are not sure of the correct dose or how to use this medicine, ask your doctor or pharmacist.**

### **Adults and adolescents 12 years and older:**

- The recommended dose is two sprays in each nostril once daily. In some cases, two sprays in each nostril twice daily may be required.
- The maximum daily dose is two sprays in each nostril twice daily.

### **Children 6 to 11 years:**

- The recommended dose is one spray in each nostril twice daily.

Clinical experience of up to 4 weeks duration showed good efficacy and safety in children. Longer experiences in children have not been available; however, clinical trials of up to one year duration using a double higher daily dose showed good safety in adults and adolescents.

Azelair is not recommended for the use in children under 6 years of age due to a lack of data on safety and/or efficacy.

If possible, you should use Azelair regularly until your symptoms have disappeared. If you interrupt the use of Azelair your symptoms are likely to return.

Azelair is suitable for long-term use.

Use longer than 4 weeks is not recommended in children 6-11 years due to lack of clinical data.

### **How to use the spray**

1. Blow your nose first.
2. Remove the protective cap (Diagram 1).
3. Before the first use, press the pump six times until an even spray emerges (Diagram 2). When Azelair has not been used for three or more days, the pump must be pressed a sufficient number of times until a fine mist emerges.

4. Spray once into each nostril keeping head upright. Do not tilt head backwards (Diagram 3).
5. Wipe and replace the protective cap.



Diagram 1




Diagram 2




Diagram 3

### **If you use more Azelair than you should**

If you spray too much Azelair into your nose you are unlikely to have any problems. If you are worried, contact your doctor.

If anyone, especially a child, accidentally drinks Azelair, contact your doctor or nearest hospital casualty department immediately.

### **If you forget to use Azelair**

Use the spray as soon as you remember, then take the next dose at the usual time. Do not take a double dose to make up for a missed dose.

If you have any further questions on the use of this product, ask your doctor or pharmacist.

## **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

These effects include:

Common (may affect up to 1 in 10 people):

- An unpleasant taste in your mouth (especially if you tilt your head backwards when you are using the nasal spray).

Uncommon (may affect up to 1 in 100 people):

- Slight irritation of the inside of the nose (stinging, itching), sneezing and nose bleed.

Rare (may affect up to 1 in 1,000 people):

- The unpleasant taste might cause you to feel sick. Fatigue (weariness, exhaustion), dizziness, weakness or a feeling of drowsiness can occur that may also be caused by the disease itself.

Very rare (may affect up to 1 in 10,000 people):

- An allergic reaction, rash, itching or nettle rash.

### **Reporting of side effects**

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

## **5. How to store Azelair**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date given on the bottle label and the outer carton. The expiry date refers to the last day of the month.

Dispose of any unused medicine 6 months after you first open the nasal spray.  
Do not store the spray in the fridge or freezer.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## **6. Contents of the pack and other information**

### **What Azelair contains**

The active substance is azelastine hydrochloride.

1 ml of nasal spray contains 1.5 mg azelastine hydrochloride. Each squirt (0.14 ml) contains 0.21 mg azelastine hydrochloride equivalent to 0.19 mg azelastine.

The other ingredients are hypromellose, sucralose, liquid sorbitol (crystallising), disodium edetate, sodium citrate, and purified water.

### **What Azelair looks like and contents of the pack**

Azelair Nasal Spray is a clear, colourless solution.

It comes in a brown glass bottle fitted with a spray pump. The 10 ml bottle contains 5 or 10 ml of the nasal spray. The 20 ml bottle contains 17 ml, 20ml or 22ml of the nasal spray.

Not all pack sizes may be marketed.

### **Marketing Authorisation Holder**

Cooper Consumer Health B.V.,  
Verrijn Stuartweg 60, 1112 AX Diemen,  
The Netherlands

### **Manufacturer**

Madaus GmbH  
Lütticher Straße 5  
53842 Troisdorf  
Germany

This leaflet was last revised in **July 2025**