

# **SUMMARY OF PRODUCT CHARACTERISTICS**

## **1 NAME OF THE MEDICINAL PRODUCT**

Cefaclor 125mg/5ml Suspension

## **2 QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each 5 ml of reconstituted suspension contains cefaclor monohydrate equivalent to 125mg of cefaclor

Excipient with known effect: Sucrose 2990.9mg per 5 ml of Oral Suspension

For the full list of excipients, see section 6.1

## **3 PHARMACEUTICAL FORM**

Powder for oral suspension.

Homogenous, white to slightly yellow coloured granule powder mixture; when dispersed in water: it is white to slightly yellow in colour with a taste and odour of fresh fruit.

## **4 CLINICAL PARTICULARS**

### **4.1 Therapeutic indications**

Cefaclor is indicated for the treatment of infections caused by sensitive micro-organisms:

- Respiratory tract infections, including pneumonia, bronchitis (including deterioration in chronic bronchitis), tonsillitis, pharyngitis and the management of sinusitis.
- Otitis media.
- Acute and chronic urinary tract infections including cystitis and pyelonephritis.

- Skin and soft tissue infections.
- Cefaclor may also be used in the eradication of streptococci from the nasopharynx, but data establishing efficacy in the subsequent prevention of either rheumatic fever or bacterial endocarditis are not available.

## 4.2 Posology and method of administration

### Posology

#### Adult:

250mg every eight hours, may be doubled to 500mg every eight hours for more severe infections or those caused by less susceptible organisms.

Total daily dosage should not exceed 4g.

#### Paediatric population:

The usual recommended dose is 20mg/kg/day in divided doses administered every eight hours. For bronchitis and pneumonia, the dosage is 20mg/kg/d in divided doses administered 3 times daily. The total daily dose may be administered in divided doses every twelve hours in otitis media and pharyngitis. This dose may be increased to 40mg/kg/day in divided doses for more severe infections, caused by less susceptible organisms.

The total daily dosage should not exceed 1g.

In the treatment of beta-haemolytic streptococcal infections, therapy should be continued for at least 10 days.

Doses may generally be administered every eight hours as follows:

<u>Age</u>	<u>Dose</u>
1 month to 1year	62.5mg
1-5 years	125mg
Over 5 years	250mg

The safe use of Cefaclor in babies aged below one month has not been proven.

#### *Cefaclor Suspension*

	<b>125mg/5ml</b>	<b>250mg/5ml</b>
--	------------------	------------------

<1 year (9kg)	2.5ml tid	
1- 5 years (9-18kg)	5.0ml tid	
Over 5 years		5.0ml tid

**Elderly:**

The normal adult dose is appropriate.

**Patients with impaired renal function:**

The normal adult dose is appropriate. (see special warnings and precautions for use)

**Patients Undergoing Haemodialysis:**

Due to a 25-30% decrease in plasma half-life, a loading dose of 250mg - 1g before dialysis is recommended with a maintenance dose between dialysis sessions of 250mg - 500mg every six to eight hours.

**Method of administration**

Oral administration.

**4.3 Contraindications**

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1

Hypersensitivity to cephalosporins

**4.4 Special warnings and precautions for use**

*Warnings*

Prior to treatment with cefaclor, every effort should be made to determine whether the patient has had previous hypersensitivity reactions to cefaclor, cephalosporins, penicillins or other drugs. Cefaclor should be given cautiously to penicillin-sensitive patients, as cross-hypersensitivity, including anaphylaxis, among beta-lactam antibiotics has been clearly documented.

If an allergic reaction to cefaclor occurs, the drug should be discontinued and the patient treated with the appropriate agents.

Pseudomembranous colitis has been reported with virtually all broad-spectrum antibiotics, including macrolides, semi-synthetic penicillins and cephalosporins. It is important, therefore, to consider its diagnosis in patients who develop diarrhoea in association with the use of antibiotics. Such colitis may range in severity from mild to life-threatening. Mild cases usually respond to drug discontinuance alone. In moderate to severe cases, appropriate measures should be taken.

#### *Precautions*

Cefaclor should be administered with caution in the presence of markedly impaired renal function. Since the half-life of cefaclor in anuric patients is 2.3 to 2.8 hours (compared to 0.6-0.9 hours in normal subjects), dosage adjustments for patients with moderate or severe renal impairment are not usually required. Clinical experience with cefaclor under such conditions is limited; therefore, careful clinical observation and laboratory studies should be made.

Broad-spectrum antibiotics should be prescribed with caution in individuals with a history of gastro-intestinal disease, particularly colitis.

Prolonged use of cefaclor may result in the overgrowth of non-susceptible organisms. If superinfection occurs during therapy, appropriate measures should be taken.

Positive direct Coombs' tests have been reported during treatment with the cephalosporin antibiotics. In haematological studies or in transfusion cross-matching procedures, when anti-globulin tests are performed on the minor side, or in Coombs' testing of newborns whose mothers have received cephalosporin antibiotics before parturition, it should be recognised that a positive Coombs' test may be due to the drug.

A false-positive reaction for glucose in the urine may occur with Benedict's or Fehling's solutions or with copper sulphate test tablets.

## **4.5 Interaction with other medicinal products and other forms of interaction**

### *Warfarin*

There have been rare reports of increased prothrombin time, with or without clinical bleeding, in patients receiving cefaclor and warfarin concomitantly. The anticoagulant effect of warfarin may be enhanced by simultaneous treatment with Cefaclor therefore prothrombin times should be monitored and warfarin dosage adjusted if necessary.

### *Probenecid*

The excretion of all cephalosporins (cefaclor) may be reduced by concomitant administration of probenecid leading to increased plasma-cephalosporin concentrations.

## **4.6 Fertility, Pregnancy and lactation**

### *Pregnancy*

Animal studies have shown no teratogenic effects and no evidence of impaired fertility. However, since there are no adequate or well controlled studies in pregnant women, care should be taken while prescribing Cefaclor for pregnant women.

### *Breast-feeding*

Studies during lactation have detected trace amounts in breast milk following administration of single 500mg doses. Average levels of about 0.2 micrograms/ml or less were detected up to 5 hours later. Trace amounts were detected at one hour. As the effect on nursing infants is not known, caution should be exercised when cefaclor is administered to a nursing woman.

## **4.7 Effects on ability to drive and use machines**

Cefaclor does not affect the ability to drive or operate machinery.

## **4.8 Undesirable effects**

### *Gastro-intestinal:*

The most frequent side-effect has been diarrhoea, though it is rarely severe enough to warrant cessation of therapy. Colitis, including rare instances of pseudomembranous colitis, has been reported. Nausea and vomiting have also occurred.

### *Hypersensitivity:*

Allergic reactions such as morbilliform eruptions, pruritus and urticaria have been observed. These reactions usually subside upon discontinuation of therapy. Serum sickness-like reactions (erythema multiforme minor, rashes or other skin manifestations accompanied by arthritis/arthralgia, with or without fever) have been reported.

Lymphadenopathy and proteinuria are infrequent; there are no circulating immune complexes and no evidence of sequelae. Occasionally, solitary symptoms may occur, but do not represent a serum sickness-like reaction. Such reactions are apparently due to hypersensitivity and have usually occurred during or following a second (or subsequent) course of therapy with cefaclor, and have been reported more frequently in children than in adults. Signs and symptoms usually occur a few days after initiation of therapy and usually subside within a few days of cessation of therapy. Antihistamines and corticosteroids appear to enhance resolution of the syndrome. No serious sequelae have been reported.

There are rare reports of erythema multiforme major (Stevens - Johnson syndrome), toxic epidermal necrolysis, and anaphylaxis. Anaphylaxis may be more common in

patients with a history of penicillin allergy. Anaphylactoid events may present as solitary symptoms, including angioedema, asthenia, oedema (including face and limbs), dyspnoea, paraesthesias, syncope, or vasodilatation.

Rarely, hypersensitivity symptoms may persist for several months.

**Haematological:**

Eosinophilia, positive Coombs' tests and, rarely, thrombocytopenia. Transient lymphocytosis, leucopenia and, rarely, haemolytic anaemia, aplastic anaemia, agranulocytosis and reversible neutropenia of possible clinical significance. See 'Interactions with other medicaments and other forms of interaction'.

**Hepatic:**

Transient hepatitis and cholestatic jaundice have been reported rarely, including slight elevations in AST, ALT or alkaline phosphatase values.

**Renal:**

Reversible interstitial nephritis has occurred rarely, also slight elevations in blood urea or serum creatinine or abnormal urinalysis.

**Central nervous system:**

Reversible hyperactivity, agitation, nervousness, insomnia, confusion, hypertonia, dizziness, hallucinations and somnolence have been reported rarely.

**Miscellaneous:**

Genital pruritus, vaginitis and vaginal moniliasis.

**Reporting of suspected adverse reactions**

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard).

## **4.9 Overdose**

The symptoms of Cefaclor overdose are non-specific and are generally nausea, vomiting, diarrhoea and gastric upsets.

Treatment for Cefaclor overdose is mainly supportive. If a large amount has been ingested (over 4g) then gastric lavage will be necessary.

General management may consist of supportive therapy.

## 5 PHARMACOLOGICAL PROPERTIES

### 5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Second generation cephalosporins antibiotics

ATC classification: J01DC04

Mode of action: The reported mode of action is predominantly by the inhibition of cell wall synthesis in susceptible bacteria. This is mainly achieved by inhibiting the trans-peptidation reaction, the final stage of the cell wall synthesis process, thus preventing the complete formation of peptidoglycan cross-links. Other earlier stages in this synthesis process may also be inhibited and there may be some induction of bacterial lysis

Mechanism of resistance: (1) Permeability – failure of cephalosporins to reach receptor sites, (2) Destroyed by beta-lactamase, (3) Alteration of PBPs

Breakpoints:

Species-related breakpoints

Species	Species-related breakpoints (S≤/R>)
<i>Staphylococcus</i>	See notes F and G
<i>Streptococcus A, B, C, G</i>	See note H
<i>H. influenzae</i>	0.5/0.5
<i>M. catarrhalis</i>	0.5/0.5

F. Susceptibility of staphylococci to cephalosporins is inferred from the methicillin susceptibility

G. For cefaclor, high dose therapy is required for the treatment of staphylococcal infections.

H. The susceptibility of streptococcus groups A, B, C and G can be inferred from their susceptibility to benzylpenicillin

Susceptibility Table

The prevalence of acquired resistance may vary geographically and with time for selected species and local information on resistance is desirable, particularly when treating severe infections. As necessary, expert advice should be sought when

the local prevalence of resistance is such that the utility of the agent in at least some types of infections is questionable.

#### Commonly susceptible species

##### Gram-positive aerobes

*Staphylococci* (coagulase positive, coagulase negative and penicillin producing strains)

Alpha and beta-haemolytic *streptococci*

*Streptococcus pneumoniae*

*Streptococcus pyogenes* (group A beta-haemolytic streptococci)

##### Gram negative aerobes

*Escherichia coli*

*Haemophilus influenzae*

*Klebsiella* species

*Moraxella catarrhalis*

*Proteus mirabilis*

#### Species for which acquired resistance may be a problem

##### Inherently resistant organisms

*Enterococci*

Methicillin resistant *staphylococci*

## **5.2 Pharmacokinetic properties**

### *Absorption*

Cefaclor is well absorbed after oral administration to fasting subjects. Total absorption is the same whether the drug is given with or without food; however, when it is taken with food, the peak concentration achieved is 50-75% of that observed when the drug is administered to fasting subjects and generally appears from  $\frac{3}{4}$  to one hour later.

### *Linearity*

Following administration of 250mg, 500mg and 1g doses to fasting subjects, average peak serum levels of approximately 7, 13 and 23 mg/l, respectively, were obtained within 30–60 minutes.

### *Biotransformation and Elimination*

Approximately 60–85% of the drug is excreted unchanged in the urine within eight hours, the greater portion being excreted within the first two hours. During the eight

hour period, peak urine concentrations following the 250mg, 500mg and 1g doses were approximately 600, 900 and 1,900 mg/l, respectively.

The serum half-life in normal subjects is 0.6–0.9 hours. In patients with reduced renal function, the serum half-life of cefaclor is slightly prolonged. In those with complete absence of renal function, the plasma half-life of the intact molecule is 2.3–2.8 hours. Excretion pathways in patients with markedly impaired renal function have not been determined. Haemodialysis shortens the half-life by 25-30%.

### **5.3 Preclinical safety data**

No further preclinical safety data.

## **6 PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Antischaumemulsion SE2

Xanthan Gum

Sodium Starch Glycollate

Strawberry Flavour

Raspberry Flavour

Sodium Lauryl Sulfate

Methylhydroxyethylcellulose

Sucrose

### **6.2 Incompatibilities**

Cefaclor has no major incompatibilities.

### **6.3 Shelf life**

Dry Powder: 2 years.

Reconstituted Suspension: 14 days.

#### **6.4 Special precautions for storage**

Dry Powder: Store below 25°C in a dry place, protected from light.

Reconstituted Suspension: Store at 2-8°C.

#### **6.5 Nature and contents of container**

Type III amber glass bottle with a polypropylene screw cap containing powder for the preparation of 100ml of suspension.

#### **6.6 Special precautions for disposal**

The suspensions should be prepared by the addition of 60ml of water.

No special requirements for disposal.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

### **7 MARKETING AUTHORISATION HOLDER**

Clydesdale Pharma Ltd  
Unit 3-4 Campbell Court  
Campbell Road  
Tadley  
RG26 5EG  
UK

### **8 MARKETING AUTHORISATION NUMBER(S)**

PL 51718/0005

**9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

15<sup>th</sup> October 1998

**10 DATE OF REVISION OF THE TEXT**

24/04/2020