

SUMMARY OF PRODUCT CHARACTERISTICS

1 NAME OF THE MEDICINAL PRODUCT

Ibuprofen 200mg Soft Capsules

Bells Healthcare Migraine Relief 200mg Soft Capsules

Bells Healthcare Period Pain Relief 200mg Soft Capsules

Bells Healthcare Fever and Headache Relief 200 mg Soft Capsules

Bells Healthcare Back and Muscle Pain Relief 200 mg Soft Capsules

Well Pharmaceuticals Ibuprofen 200mg Soft Capsules

Numark Back & Muscle Pain Relief 200 mg Soft Capsules

Almus Fever and Headache Relief 200mg Soft Capsules

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each capsule contains Ibuprofen 200 mg.

Excipient(s) with known effect:

sorbitol (E420), ponceau 4R red (E124)

For the full list of excipients, see section 6.1.

3 PHARMACEUTICAL FORM

Capsule, soft.

Wine red colour oval shaped soft gelatin capsules containing pink to dark pink coloured viscous liquid, printed '142' in white colour on capsule shell.

4 CLINICAL PARTICULARS

4.1 Therapeutic indications

Ibuprofen 200mg Soft Capsules are indicated adults and children over 12 years for the symptomatic relief of:

- rheumatic or muscular pain
- backache
- neuralgia
- migraine
- headache
- dental pain

- dysmenorrhoea
- feverishness
- colds and influenza symptoms

4.2 Posology and method of administration

For oral administration and short-term use only.

Adults, elderly and children over 12 years:

The lowest effective dose should be used for the shortest duration necessary to relieve symptoms (see section 4.4).

The patient should consult a doctor if the symptoms persist or worsen, or the product is required for more than 10 days.

Adolescents (age range: ≥ 12 years to < 18 years): If this medicinal product is required for more than 3 days, or if symptoms persist or worsen, consult your doctor.

Take one or two capsules with water to start with, repeat up to three times a day as required.

Leave at least 4 hours between doses and do not take more than six capsules in any 24-hour period.

Not suitable for children under 12 years.

4.3 Contraindications

Hypersensitivity to ibuprofen or to any of the excipients.

Patients who have previously shown hypersensitivity reactions (e.g. asthma, rhinitis, angioedema or urticaria), in response to ibuprofen, aspirin or other non-steroidal anti-inflammatory drugs.

Active or history of recurrent peptic ulcer/haemorrhage (two or more distinct episodes of proven ulceration or bleeding).

History of gastrointestinal bleeding or perforation, related to previous NSAIDs therapy.

Severe heart failure (NYHA Class IV), renal failure or hepatic failure (see section 4.4).

Third trimester of pregnancy (see section 4.6 Fertility, pregnancy and lactation).

Children under 12 years.

4.4 Special warnings and precautions for use

Undesirable effects may be minimised by using the lowest effective dose for the shortest duration necessary to control symptoms (see GI and cardiovascular risks below).

The elderly have an increased frequency of adverse reactions to NSAIDs especially gastrointestinal bleeding and perforation which may be fatal.

Respiratory:

Bronchospasm may be precipitated in patients suffering from, or with a history of, bronchial asthma or allergic disease.

Other NSAIDs:

The use of ibuprofen with concomitant NSAIDs including cyclooxygenase-2 selective inhibitors should be avoided (see section 4.5)

SLE and mixed connective tissue disease:

Systemic lupus erythematosus and mixed connective tissue disease – increased risk of aseptic meningitis (see section 4.8).

Renal:

Renal impairment as renal function may further deteriorate (see sections 4.3 and 4.8)

There is a risk of renal impairment in dehydrated children and adolescents.

Renal tubular acidosis and hypokalaemia may occur following acute overdose and in patients taking ibuprofen products over long periods at high doses (typically greater than

4 weeks), including doses exceeding the recommended daily dose.

Hepatic:

Hepatic dysfunction (see Sections 4.3 and 4.8)

Cardiovascular and cerebrovascular effects:

Caution (discussion with doctor or pharmacist) is required prior to starting treatment in patients with a history of hypertension and/or heart failure as fluid retention, hypertension and oedema have been reported in association with NSAID therapy.

Clinical studies suggest that the use of ibuprofen, particularly at a high dose (2400 mg daily) may be associated with a small increased risk of arterial thrombotic events (for example myocardial infarction or stroke). Overall, epidemiological studies do not suggest that low dose ibuprofen (e.g. ≤1200mg daily) is associated with an increased risk of arterial thrombotic events.

Patients with uncontrolled hypertension, congestive heart failure (NYHA II-III), established ischaemic heart disease, peripheral arterial disease, and/or cerebrovascular disease should only be treated with ibuprofen after careful consideration and high doses (2400 mg/day) should be avoided.

Careful consideration should also be exercised before initiating long-term treatment of patients with risk factors for cardiovascular events (e.g. hypertension, hyperlipidaemia, diabetes mellitus, smoking), particularly if high doses of ibuprofen (2400 mg/day) are required.

Cases of Kounis syndrome have been reported in patients treated with ibuprofen. Kounis syndrome has been defined as cardiovascular symptoms secondary to an allergic or hypersensitive reaction-associated with constriction of coronary arteries and potentially leading to myocardial infarction.

Impaired female fertility:

There is some evidence that drugs which inhibit cyclo-oxygenase/ prostaglandin synthesis may cause impairment of female fertility by an effect on ovulation. This is reversible on withdrawal of treatment.

Gastrointestinal:

NSAIDs should be given with care to patients with a history of gastrointestinal disease (ulcerative colitis, Crohn's disease) as these conditions may be exacerbated (see section 4.8).

GI bleeding, ulceration or perforation, which can be fatal, has been reported with all NSAIDs at anytime during treatment, with or without warning symptoms or a previous history of GI events.

The risk of GI bleeding, ulceration or perforation is higher with increasing NSAID doses, in patients with a history of ulcer, particularly if complicated with haemorrhage or perforation (see section 4.3), and in the elderly. These patients should commence treatment on the lowest dose available.

Patients with a history of GI toxicity, particularly the elderly, should report any unusual abdominal symptoms (especially GI bleeding) particularly in the initial stages of treatment.

Caution should be advised in patients receiving concomitant medications which could increase the risk of ulceration or bleeding, such as oral corticosteroids, anticoagulants such as warfarin, selective serotonin-reuptake inhibitors or anti-platelet agents such as aspirin (see section 4.5).

When GI bleeding or ulceration occurs in patients receiving ibuprofen, the

treatment should be withdrawn.

Dermatological:

Severe cutaneous adverse reactions (SCARs)

Severe cutaneous adverse reactions (SCARs) , including exfoliative dermatitis, erythema multiforme, Stevens-Johnson syndrome (SJS), and toxic epidermal necrolysis, (TEN), Drug Reactions with Eosinophilia and Systemic Symptoms (DRESS syndrome), and acute generalised exanthematous pustulosis (AGEP), which can be life-threatening or fatal, have been reported in association with the use of NSAIDs ibuprofen (see section 4.8).

Most of these reactions occurred within the first month.

If signs and symptoms suggestive of these reactions appear ibuprofen should be withdrawn immediately and an alternative treatment considered (as appropriate).

Masking of symptoms of underlying infections

Ibuprofen can mask symptoms of infection, which may lead to delayed initiation of appropriate treatment and thereby worsening the outcome of the infection. This has been observed in bacterial community acquired pneumonia and bacterial complications to varicella. When ibuprofen is administered for fever or pain relief in relation to infection, monitoring of infection is advised. In non-hospital settings, the patient should consult a doctor if symptoms persist or worsen.

Lactose intolerance:

Patients with rare hereditary problems of galactose intolerance, total lactase deficiency or glucose-galactase malabsorption should not take this medicine.

Intolerance to some sugars:

Patients with rare hereditary problems of galactose intolerance, fructose intolerance, the Lapp lactase deficiency, glucose-galactose malabsorption or sucrase-isomaltase insufficiency should not take this medicine.

The label will include:

Read the enclosed leaflet before taking this product

Do not take if you:

- have (or have had two or more episodes of) a stomach ulcer, perforation or bleeding
- are allergic to ibuprofen, to any of the ingredients, or to aspirin or other painkillers
- are taking other NSAID pain killers or aspirin with a daily dose above 75mg

Speak to a pharmacist or your doctor before taking if you:

- have or have had asthma, diabetes, high cholesterol, high blood pressure, a stroke, heart, liver, kidney or bowel problems
- are a smoker
- are pregnant

This medicine contains 17.4 mg of potassium per dose.

To be taken into consideration by patients with reduced kidney function or patient on a controlled potassium diet.

Contains 58.6 mg of sorbitol per dose, a source of 14.6 mg of fructose per dose.

Patients with rare hereditary problems of fructose intolerance should not take this medicine.

If symptoms persist or worsen, or if new symptoms occur, consult your doctor or pharmacist.

This medicine contains Ponceau 4R. May cause allergic reactions.

4.5 Interaction with other medicinal products and other forms of interaction

Ibuprofen (like other NSAIDs) should be avoided in combination with:

- *Aspirin (acetylsalicylic acid):* Concomitant administration of ibuprofen and acetylsalicylic acid is not generally recommended because of the potential of increased adverse effects, unless low-dose aspirin (not above 75mg daily) has been advised by a doctor (see Section 4.4).

Experimental data suggest that ibuprofen may competitively inhibit the effect of low dose aspirin on platelet aggregation when they are dosed concomitantly. Although there are uncertainties regarding extrapolation of these data to the clinical situation, the possibility that regular, long-term use of ibuprofen may reduce the cardioprotective effect of low-dose acetylsalicylic acid cannot be excluded. No clinically relevant effect is considered to be likely for occasional ibuprofen use (see section 5.1).

- *Other NSAIDs including cyclooxygenase-2 selective inhibitors:*
Avoid concomitant use of two or more NSAIDs as this may increase the risk of adverse effects (see section 4.4)

Ibuprofen should be used with caution in combination with:

- *Anticoagulants:* NSAIDs may enhance the effects of anti-coagulants, such as warfarin (See section 4.4).

- *Antihypertensives (ACE inhibitors and Angiotensin II Antagonists) and diuretics:* since NSAIDs may diminish the effect of these drugs. In some patients with compromised renal function (e.g. dehydrated patients or elderly patients with compromised renal function) the co-administration of an ACE inhibitor or Angiotensin II antagonist and agents that inhibit cyclo-oxygenase may result in further deterioration of renal function, including possible acute renal failure, which is usually reversible. These interactions should be considered in patients taking a coxib concomitantly with ACE inhibitors or angiotensin II antagonists. Therefore, the combination should be administered with caution, especially in the elderly. Patients should be adequately hydrated and consideration should be given to monitoring of renal function after initiation of concomitant therapy, and periodically thereafter.

- *Diuretics* can increase the risk of nephrotoxicity of NSAIDs.

- *Corticosteroids:* as these may increase the risk of gastrointestinal ulceration or bleeding (see Section 4.4)

- *Anti-platelet agents and selective serotonin reuptake inhibitors (SSRIs):* increased risk of gastrointestinal bleeding (see section 4.4).

- *Cardiac glycosides:* NSAIDs may exacerbate cardiac failure, reduce GFR and increase plasma glycoside levels.

- *Lithium:* There is evidence for potential increase in plasma levels of lithium.

- *Methotrexate:* There is evidence for the potential increase in plasma levels of methotrexate.

- *Ciclosporin:* Increased risk of nephrotoxicity.

- *Mifepristone:* NSAIDs should not be used for 8-12 days after mifepristone administration as NSAIDs can reduce the effect of mifepristone.

- *Tacrolimus*: Possible increased risk of nephrotoxicity when NSAIDs are given with tacrolimus.
- *Zidovudine*: Increased risk of haematological toxicity when NSAIDs are given with zidovudine. There is evidence of an increased risk haemarthroses and haematoma in HIV (+) haemophiliacs receiving concurrent treatment with zidovudine and ibuprofen.
- *Quinolone antibiotics*: Animal data indicate that NSAIDs can increase the risk of convulsions associated with quinolone antibiotics. Patients taking NSAIDs and quinolones may have an increased risk of developing convulsions.

4.6 Fertility, pregnancy and lactation

Inhibition of prostaglandin synthesis may adversely affect the pregnancy and/or the embryo/foetal development. Data from epidemiological studies suggest an increased risk of miscarriage and of cardiac malformation and gastroschisis after use of a prostaglandin synthesis inhibitor in early pregnancy. The absolute risk for cardiovascular malformation was increased from less than 1%, up to approximately 1.5 %. The risk is believed to increase with dose and duration of therapy. In animals, administration of a prostaglandin synthesis inhibitor has been shown to result in increased pre- and post-implantation loss and embryo-foetal lethality. In addition, increased incidences of various malformations, including cardiovascular, have been reported in animals given a prostaglandin synthesis inhibitor during the organogenetic period.

From the 20th week of pregnancy onward, ibuprofen use may cause oligohydramnios resulting from foetal renal dysfunction. This may occur shortly after treatment initiation and is usually reversible upon discontinuation. In addition, there have been reports of ductus arteriosus constriction following treatment in the second trimester, most of which resolved after treatment cessation. Therefore, during the first and second trimester of pregnancy, ibuprofen should not be given unless clearly necessary. If ibuprofen is used by a woman attempting to conceive, or during the first and second trimester of pregnancy, the dose should be kept as low and duration of treatment as short as possible. Antenatal monitoring for oligohydramnios and ductus arteriosus constriction should be considered after exposure to ibuprofen for several days from gestational week 20 onward. Ibuprofen should be discontinued if oligohydramnios or ductus arteriosus constriction are found.

During the third trimester of pregnancy, all prostaglandin synthesis inhibitors

may expose the foetus to:

- cardiopulmonary toxicity (with premature constriction/closure of the ductus arteriosus and pulmonary hypertension);
 - renal dysfunction (see above) which may progress to renal failure with oligohydroamniosis;
 - the mother and the neonate, at the end of the pregnancy, to
 - possible prolongation of bleeding time, an anti-aggregating effect which may occur even at very low doses.
 - inhibition of uterine contractions resulting in delayed or prolonged labour.
- Consequently, ibuprofen is contraindicated during the third trimester of pregnancy (see section 4.3).

Breast-feeding

In limited studies, ibuprofen appears in the breast milk in very low concentration and is unlikely to affect the breast-fed infant adversely.

Fertility

See section 4.4 regarding female fertility.

4.7 Effects on ability to drive and use machines

None expected at recommended dose and duration of therapy.

4.8 Undesirable effects

The following list of adverse effects relates to those experienced with ibuprofen at OTC doses (maximum 1200 mg ibuprofen per day), in short-term use. In the treatment of chronic conditions, under long-term treatment, additional adverse events may occur.

Adverse events which have been associated with Ibuprofen are given below, listed by system organ class and frequency. Frequencies are defined as: very common ($\geq 1/10$), common ($\geq 1/100$ to $< 1/10$), uncommon ($\geq 1/1000$ to $< 1/100$), rare ($\geq 1/10,000$ to $< 1/1000$), very rare ($< 1/10,000$) and not known (cannot be estimated from the available data). Within each frequency grouping, adverse events are presented in order of decreasing seriousness.

The adverse events observed most often are gastrointestinal in nature. Adverse events are mostly dose-dependent, in particular the risk of occurrence of gastrointestinal bleeding is dependent on the dosage range and duration of treatment.

Clinical studies suggest that the use of ibuprofen, particularly at a high dose (2400mg/day) may be associated with a small increased risk of arterial thrombotic events (for example myocardial infarction or stroke) (see section 4.4).

System Organ Class	Frequency	Adverse Event
Blood and Lymphatic System Disorders	Very rare	Haematopoietic disorders (anaemia, leucopenia, thrombocytopenia, pancytopenia, agranulocytosis). First signs are: fever, sore throat, superficial mouth ulcers, flu-like symptoms, severe exhaustion, unexplained bleeding and bruising.
Immune System Disorders	Uncommon	Hypersensitivity reactions consisting of ¹ : Urticaria and pruritus
	Very rare	Severe hypersensitivity reactions. Symptoms could be facial, tongue and laryngeal swelling, dyspnoea, tachycardia, hypotension (anaphylaxis, angioedema or severe shock).
	Not Known	Respiratory tract reactivity comprising asthma, aggravated asthma, bronchospasm or dyspnoea.
Nervous System Disorders	Uncommon	Headache
	Very rare	Aseptic meningitis ²
Cardiac Disorders	Not Known	Cardiac failure and oedema. Kounis syndrome.

Vascular Disorders	Not Known	Hypertension
Gastrointestinal Disorders	Uncommon	Abdominal pain, nausea, dyspepsia
	Rare	Diarrhoea, flatulence, constipation and vomiting
	Very rare	Peptic ulcer, perforation or gastrointestinal haemorrhage, melaena, haematemesis, sometimes fatal, particularly in the elderly. Ulcerative stomatitis, gastritis
	Not Known	Exacerbation of colitis and Crohn's disease (section 4.4).
Hepatobiliary Disorders	Very rare	Liver disorders
Skin and Subcutaneous Tissue Disorders	Uncommon	Various skin rashes
	Very rare	Severe cutaneous adverse reactions (SCARs) (including erythema multiforme, exfoliative dermatitis, Stevens-Johnson syndrome (SJS) and toxic epidermal necrolysis (TEN)).
	Not Known	Drug reaction with eosinophilia and systemic symptoms (DRESS syndrome) Acute generalised exanthematous pustulosis (AGEP) Photosensitivity reactions
Renal and Urinary Disorders	Very rare	Acute renal failure, papillary necrosis,

		especially in long-term use, associated with increased serum urea and oedema.
	Not Known	Renal insufficiency, Renal tubular acidosis ³ , ureteric colic, dysuria.
Investigations	Very rare	Decreased haemoglobin levels
Metabolism and nutrition disorders	Not Known	Hypokalaemia ³ , decreased appetite
Infections and infestations	Very rare	Exacerbation of infection related inflammations (e.g. development of necrotising fasciitis) coinciding with the use of non-steroidal anti-inflammatory drugs has been described. ⁴

Description of Selected Adverse Reactions

¹ Hypersensitivity reactions have been reported following treatment with ibuprofen. These may consist of (a) non-specific allergic reactions and anaphylaxis, (b) respiratory tract activity comprising asthma, aggravated asthma, bronchospasm, dyspnoea or (c) assorted skin disorders, including rashes of various types pruritus, urticaria, purpura, angioedema and more rarely exfoliative and bullous dermatoses (including epidermal necrolysis and erythema multiforme).

²The pathogenic mechanism of drug-Induced aseptic meningitis is not fully understood. However, the available data on NSAID-related aseptic meningitis points to a hypersensitivity reaction (due to a temporal relationship with drug intake, and disappearance of symptoms after drug discontinuation). Of note, single cases of symptoms of aseptic meningitis (such as stiff neck, headache, nausea, vomiting, fever or disorientation) have been observed during treatment with ibuprofen, in patients with existing auto-immune disorders (such as systemic lupus erythematosus, mixed connective tissue disease).

³*Renal tubular acidosis and hypokalaemia have been reported in the post-marketing setting typically following prolonged use of the ibuprofen component at higher than recommended doses.

⁴This is possibly associated with the mechanism of action of the non-steroidal anti-inflammatory drugs. If signs of an infection occur or get worse during use of Ibuprofen the patient is therefore recommended to go to a doctor without delay. It is to be investigated whether there is an indication for anti-infective/antibiotic therapy.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store.

4.9 Overdose

In children ingestion of more than 400 mg/kg may cause symptoms.

In adults the dose response effect is less clear cut. The half-life in overdose is 1.5-3 hours.

Symptoms

Most patients who have ingested clinically important amounts of NSAIDs will develop no more than nausea, vomiting, epigastric pain, or more rarely diarrhoea. Tinnitus, headache and gastrointestinal bleeding are also possible. In more serious poisoning, toxicity is seen in the central nervous system, manifesting as drowsiness, occasionally excitation and disorientation or coma. Occasionally patients develop convulsions. In serious poisoning metabolic acidosis may occur and the prothrombin time/ INR may be prolonged, probably due to interference with the actions of circulating clotting factors. Acute renal failure and liver damage may occur. Exacerbation of asthma is possible in asthmatics.

In serious poisoning metabolic acidosis may occur and the prothrombin time/INR may be prolonged, probably due to interference with the actions of circulating clotting factors. Acute renal failure and liver damage may occur.

Prolonged use at higher than recommended doses may result in severe hypokalaemia and renal tubular acidosis. Symptoms may include reduced level of consciousness and generalised weakness (see section 4.4 and section 4.8).

Management

Management should be symptomatic and supportive and include the maintenance of a clear airway and monitoring of cardiac and vital signs until stable. Consider oral administration of activated charcoal if the patient presents within 1 hour of ingestion of a potentially toxic amount. If frequent or prolonged, convulsions should be treated with intravenous diazepam or lorazepam. Give bronchodilators for asthma.

5 PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

ATC Code: M01A E01 Propionic acid derivative

Ibuprofen is a propionic acid derivative NSAID that has demonstrated its efficacy by inhibition of prostaglandin synthesis. In humans, ibuprofen reduces inflammatory pain, swellings and fever. Furthermore, ibuprofen reversibly inhibits platelet aggregation.

Clinical evidence demonstrates that when 400mg of ibuprofen is taken the pain relieving effects can last for up to 8 hours.

Experimental data suggest that ibuprofen may competitively inhibit the effect of low dose acetylsalicylic acid on platelet aggregation when they are dosed concomitantly. Some pharmacodynamic studies show that , when a single dose of ibuprofen 400mg were taken within 8 h before or within 30 min after immediate release aspirin dosing (81mg), a decreased effect of ASA on the formation of thromboxane or platelet aggregation occurred.

Although there are uncertainties regarding extrapolation of these data to the clinical situation, the possibility that regular, long-term use of ibuprofen may reduce the cardioprotective effect of low dose acetylsalicylic acid cannot be excluded.

No clinically relevant effect is considered to be likely for occasional ibuprofen use (see section 4.5).

5.2 Pharmacokinetic properties

Ibuprofen is well absorbed from the gastrointestinal tract. Ibuprofen is

extensively bound to plasma proteins.

Ibuprofen 200mg Soft Capsules consist of ibuprofen 200 mg dissolved in a hydrophilic solvent inside a gelatin shell. On ingestion, the gelatin shell disintegrates in the gastric juice releasing the solubilised ibuprofen immediately for absorption. The median peak plasma concentration is achieved approximately 30 minutes after administration.

The median peak plasma concentration for Ibuprofen tablets is achieved approximately 1-2 hours after administration.

Ibuprofen is metabolised in the liver to two major metabolites with primary excretion via the kidneys, either as such or as major conjugates, together with a negligible amount of unchanged ibuprofen. Excretion by the kidney is both rapid and complete.

Elimination half-life is approximately 2 hours.

In limited studies, ibuprofen appears in the breast milk in very low concentrations.

No significant differences in pharmacokinetic profile are observed in the elderly.

5.3 Preclinical safety data

No relevant information, additional to that contained elsewhere in the SPC.

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Capsule contents:

Macrogol 400
Sorbitan mono-oleate
Sorbitol
Potassium hydroxide (E525)
Water, purified.

Capsule shell:

Gelatin (E441)
Macrogol 400
Sorbitol (E420)
Ponceau 4 R red (E124)
Water, purified.
Triglycerides – medium chain

Capsule printing:

Shellac, glaze
Titanium dioxide (E171)
Propylene glycol

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

3 years

6.4 Special precautions for storage

Store below 25°C

Store in the original pack.

6.5 Nature and contents of container

Blisters PVC / Aluminium and PVdC / Aluminium packed into cartons.

Each carton may contain 10, 12 or 16 capsules in blister strips

Not all pack sizes may be marketed.

6.6 Special precautions for disposal

No special requirements.

7 MARKETING AUTHORISATION HOLDER

Bell, Sons & Co (Druggists) Ltd.,
Gifford House,
Slaidburn Crescent,
Southport,
Merseyside,
PR9 9AL,
United Kingdom.

8 MARKETING AUTHORISATION NUMBER(S)

PL 01305/0105

**9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE
AUTHORISATION**

18/08/2025

10 DATE OF REVISION OF THE TEXT

22/05/2026