

SUMMARY OF PRODUCT CHARACTERISTICS

1 NAME OF THE MEDICINAL PRODUCT

Gentamicin Paediatric 20mg/2ml Solution for Injection,

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each vial (2 ml) contains Gentamicin Sulphate Ph. Eur. equivalent to 20 mg Gentamicin base.

For the full list of excipients, see section 6.1.

3 PHARMACEUTICAL FORM

Solution for Injection.

Clear, colourless solution

4 CLINICAL PARTICULARS

4.1 Therapeutic indications

Gentamicin is an aminoglycoside antibiotic with broad-spectrum bactericidal activity. It is indicated to treat severe infections caused by bacteria susceptible to gentamicin such as, but not limited to:

- Urinary tract infections
- Respiratory tract infections
- Intra-abdominal infections
- CNS infections
- Severe neonatal infections

It is usually active against most strains of the following organisms: *Escherichia coli*, *Klebsiella spp.*, *Proteus spp.* (indole positive and indole negative), *Pseudomonas aeruginosa*, *Staphylococci*, *Enterobacter spp.*, *Citrobacter spp.* and *Providencia spp.*

Consideration should be given to official local guidance on the appropriate use of antibacterial agents.

4.2 Posology and method of administration

Posology

Adults

The recommended dose in adults with normal renal function is 3-5 mg/kg/day, depending on the severity of infection, administered as one single dose (preferred) or in two divided doses. The dose should be adjusted according to clinical response and serum concentration levels (see below). Dose calculations should be based on ideal body weight. A dosing frequency of more than twice daily may be adopted for some specific pathogens or some sites of infection as recommended in national and local guidance.

Once daily dosing is not recommended in cases of endocarditis, depending on the responsible pathogens. National and local guidance on treatment with gentamicin and serum level monitoring in endocarditis should be followed.

In patients with normal renal function, 160 mg once daily may be used for the treatment of urinary tract infections.

Paediatric population

The daily dose recommended in children aged 1 year and above and adolescents with normal renal function, is 3 – 6 mg/kg/day as one single dose (preferred) or two divided doses.

The daily dose in infants after the first month of life is 4.5 – 7.5 mg/kg/day as one single dose (preferred) or two divided doses.

The daily dose in neonates and pre-term infants (aged 0 – 4 weeks old) is 4 – 7 mg/kg/day. Due to the longer half-life, newborns are given the required daily dose in one single dose.

Elderly

There is some evidence that elderly patients may be more susceptible to aminoglycoside toxicity whether secondary to previous auditory/vestibular impairment or borderline renal dysfunction. Accordingly, therapy should be closely monitored by frequent determination of gentamicin serum levels, assessment of renal function and signs of ototoxicity.

Renal impairment

In impaired renal function, the recommended daily dose has to be decreased and adjusted to the renal function. This can be achieved by reducing the dose and/or increasing the dose interval.

In all patients with renal impairment, serum gentamicin peak and trough concentration and renal function must be monitored frequently (see below).

Nomograms are available for the calculation of dose, which depends on the patient's age, weight and renal function. Local guidance should be followed where available.

No clear recommendation can be made for once daily dosing; dosing should be guided by plasma concentration levels. In patients with moderate renal impairment, in whom once daily dosing would be considered appropriate if their renal function were normal, the dose interval should be at least 24 hours and extended according to the degree of renal impairment and the results of serum gentamicin monitoring. Limited data are available in patients with severe renal impairment (creatinine clearance < 30 ml/min) after once daily dose administration.

The following table may be useful for adults on multiple daily dose regimens:

Blood Urea		Creatinine clearance (GFR) (ml/min)	Dose & frequency of administration
(mg/100 ml)	(mmol/l)		
< 40	6 – 7	> 70	80 mg* 8 hourly
40 – 100	6 – 17	30 – 70	80 mg* 12 hourly
100 – 200	17 – 34	10 – 30	80 mg* daily
> 200	> 34	5 – 10	80 mg* every 48 hours
Twice weekly intermittent haemodialysis		< 5	80 mg* after dialysis

*60 mg if body weight < 60kg.

Monitoring advice

Regular serum concentration monitoring of gentamicin is recommended for all patients, and especially in the elderly, newborns, obesity and in patients with impaired renal function, as well as patients with cystic fibrosis. Gentamicin should not be prescribed if serum concentrations cannot be monitored.

There are no universally accepted guidelines for therapeutic drug monitoring of gentamicin. Local monitoring and dose adjustment guidelines should be followed where available.

Pre-dose (“trough level”) monitoring is recommended to ensure that the interval between doses is correct. Trough levels are measured at the end of a dosing interval and should not exceed 1mg/L for once daily dosing or 2 mg/L for multiple daily dosing. Levels in excess of these indicate the need to extend the interval between doses, not reduction of the dose.

Post-dose (“peak level”) monitoring is recommended to check the adequacy of a dose or to ensure that it is not excessive and likely to cause toxicity. Peak levels should be measured one hour after an intravenous bolus or intramuscular bolus dose, or 30 minutes after the end of an infusion. A plasma concentration < 4 mg/L indicates that the dose is likely to be inadequate and a dose increase should be considered; plasma concentrations > 10 mg/L indicate an increased risk for toxicity, particularly ototoxicity, and a dose reduction should be considered.

Any change in dose should be re-assessed with pre- and post-dose levels to confirm the adequacy of the new dose and the appropriateness of the dose interval.

Method of administration

The recommended dose and precautions for intramuscular and intravenous administration are identical. Gentamicin when given intravenously should be injected directly into a vein or into the drip set tubing over no less than three minutes. If administered by infusion, this should be over 20 - 30 minutes and in no greater volume of fluid than 100 ml. Longer infusion times of up to 60 minutes may be used, in particular for a once daily dosing regimen. Once daily dosing should only be administered through the intravenous route.

4.3 Contraindications

- Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

- Myasthenia gravis.

4.4 Special warnings and precautions for use

Ototoxicity and nephrotoxicity

Ototoxicity has been reported following the use of aminoglycosides, including gentamicin. Symptoms include loss of balance and hearing loss, which may be irreversible (see section 4.8). Important risk factors include renal impairment, high doses, prolonged duration of treatment and age (neonates/infants and possibly the elderly). Due to the potential for ototoxicity and nephrotoxicity, monitoring of vestibule, cochlea and renal function is recommended before, during and shortly after treatment (see section 4.8). Serum levels are determined so as to avoid peak concentrations above 10 mg/l and troughs above 1 mg/l when administering gentamicin once daily and 2 mg/l when administering gentamicin twice daily.

As there is some evidence that risk of both ototoxicity and nephrotoxicity is related to the level of total exposure, duration of therapy should be the shortest possible compatible with clinical recovery. In some patients with impaired renal function there has been a transient rise in blood-urea-nitrogen which has usually reverted to normal during or following cessation of therapy. It is important to adjust the frequency of dosage according to the degree of renal function.

There is an increased risk of ototoxicity in patients with mitochondrial DNA mutations, (particularly the nucleotide 1555 A to G substitution in the 12S rRNA gene), even if aminoglycoside serum levels are within the recommended range during treatment. Alternative treatment options should be considered in such patients. Mitochondrial mutations are rare, and the penetrance of this observed effect is unknown.

To avoid adverse events, continuous monitoring (before, during and after treatment) of hepatic and laboratory parameters is also recommended

In patients with a maternal history of relevant mutations or aminoglycoside induced deafness, alternative treatments or genetic testing prior to administration should be considered.

Drug-resistant microorganisms

Treatment with gentamicin may produce an excessive growth of drug-resistant microorganisms. If this happens, an appropriate treatment should be initiated.

Clostridium difficile infection

Diarrhoea and pseudomembranous colitis have been observed when gentamicin is combined with other antibiotics. These diagnoses should be considered in every patient that develops diarrhoea during or immediately after treatment. Gentamicin should be discontinued if the patient suffers severe diarrhoea and/or bloody diarrhoea during treatment and an appropriate treatment should be initiated. Drugs that inhibit peristalsis should not be administered (see section 4.8).

Pregnancy

Gentamicin should only be used in pregnancy if considered essential by the physician (see section 4.6).

Conditions characterised by muscular weakness.

Gentamicin should be used with care in conditions characterised by muscular weakness.

Obesity

In cases of significant obesity gentamicin serum concentrations should be closely monitored and a reduction in dose should be considered.

Gentamicin Paediatric solution contains sodium:

This medicine contains 6.81 mg of sodium per 2 ml vial.

4.5 Interaction with other medicinal products and other forms of interaction

Concurrent administration of gentamicin and other potentially ototoxic or nephrotoxic drugs should be avoided. Potent diuretics such as etacrynic acid and furosemide are believed to enhance the risk of ototoxicity whilst amphotericin B, cisplatin and ciclosporin are potential enhancers of nephrotoxicity.

Any potential nephrotoxicity of cephalosporins, and in particular cephaloridine, may also be increased in the presence of gentamicin. Consequently, if this combination is used monitoring of kidney function is advised.

Neuromuscular blockade and respiratory paralysis have been reported from administration of aminoglycosides to patients who have received curare-type muscle relaxants during anaesthesia.

Indometacin possibly increases plasma concentrations of gentamicin in neonates.

Concurrent use with oral anticoagulants may increase the hypothermibrinanaemic effect.

Concurrent use of bisphosphonates may increase the risk of hypocalcaemia.

Concurrent use of the Botulinum Toxin and gentamicin may increase the risk of toxicity due to enhanced neuromuscular block.

Antagonism of effect may occur with concomitant administration of gentamicin with either neostigmine or pyridostigmine.

4.6 Fertility, pregnancy and lactation

Pregnancy

There are no proven cases of intrauterine damage caused by gentamicin. However, in common with most drugs known to cross the placenta, usage in pregnancy should only be considered in life-threatening situations where the expected benefits outweigh possible risks.

Breast-feeding

In the absence of gastro-intestinal inflammation, the amount of gentamicin ingested from the milk is unlikely to result in significant blood levels in breast-fed infants.

4.7 Effects on ability to drive and use machines

Not known.

4.8 Undesirable effects

The following CIOMS frequency rating is used, when applicable: Very common ($\geq 1/10$); common ($\geq 1/100$ to $\leq 1/10$); uncommon ($\geq 1/1,000$ to $\leq 1/100$); rare ($\geq 1/10,000$ to $\leq 1/1,000$); very rare ($\leq 1/10,000$); not known (cannot be estimated from the available data).

Infections and infestations:

Not known: antibiotic-associated colitis, pseudomembranous colitis, Superinfection (caused by gentamicin-resistant bacteria),

Blood and lymphatic system disorders:

Not known: anaemia, blood dyscrasia

Immune system disorders:

Not known: hypersensitivity, anaphylaxis/anaphylactic reaction (including anaphylactic shock)

Metabolism and nutrition disorders:

Not known: hypomagnesaemia on prolonged therapy

Psychiatric disorders:

Not known: depression, hallucinations, confusion

Nervous system disorders:

Not known: central neuropathy (including convulsions, lethargy, encephalopathy), peripheral neuropathy

Ear and labyrinth disorders:

Not known: transitory hearing loss, irreversible hearing loss, deafness, particularly after exposure to ototoxic drugs or in the presence of renal dysfunction (see section 4.4).

Gastrointestinal disorders:

Not known: stomatitis, nausea, vomiting

Hepatobiliary disorders:

Not known: abnormal liver function, transaminases increased

Skin and subcutaneous tissue disorders:

Not known: rash, purpura, urticaria, pruritus, Steven Johnson syndrome, toxic epidermal necrosis

Renal and urinary disorders:

Very rare: acute renal failure, Fanconi-like syndrome in patients treated with a prolonged course of high-dose

Not known: nephrotoxicity (usually reversible) has been reported.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store.

4.9 Overdose

Haemodialysis and peritoneal dialysis will aid removal from the blood but the former is probably more efficient. Calcium salts given intravenously have been used to counter the neuromuscular blockade caused by gentamicin.

5 PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Antibacterials for systemic use, ATC Code: J01GB03

Gentamicin is a mixture of antibiotic substances produced by the growth of *Micromonospora purpurea*. It is bactericidal with greater antibacterial activity than streptomycin, neomycin or kanamycin.

Gentamicin exerts a number of effects on cells of susceptible bacteria. It affects the integrity of the plasma membrane and the metabolism of RNA, but its most important effect is inhibition of protein synthesis at the level of the 30s ribosomal subunit.

5.2 Pharmacokinetic properties

Gentamicin is not readily absorbed from the gastro-intestinal tract. Gentamicin is 70 – 85% bound to plasma albumin following administration and is excreted 90% unchanged in urine. The half-life for its elimination in normal patients is 2 – 3 hours.

- Effective plasma concentration is 4 – 8 µg/ml.
- The volume of distribution (VD) is 0.31 kg.
- The elimination rate constant is:
 1. 0.02 hr⁻¹ for anuric patients*
 2. 0.30 hr⁻¹ normal

*Therefore in those with anuria care must be exercised following the initial dose, any subsequent administration being reduced in-line with plasma concentrations of gentamicin.

Paediatric population (premature infants and neonates)

Distribution

The distribution volume of gentamicin is about equivalent to the volume of extracellular water. In the newborn water makes up 70 – 75% of bodyweight, compared with 50 – 55% in adults. The extracellular water compartment is larger (40% of body weight compared with 25% of body weight in adults). Therefore, the volume of distribution of gentamicin per kg bodyweight is affected and decreases with increasing age from 0.5 – 0.7 l/kg for a premature newborn to 0.25 l/kg for an adolescent. The larger volume of distribution per kg bodyweight means that for adequate peak blood concentration a higher dose per kg bodyweight needs to be administered.

Elimination

Gentamicin is not metabolized in the body but is excreted unchanged in microbiologically active form predominantly via the kidneys. In patients with normal renal function the elimination half-life is about 2 – 3 hours. In neonates elimination rate is reduced due to immature renal function.

Elimination half-life averages approximately 8 hours in neonates at a gestational age of 26 – 34 weeks compared with about 6.7 hours in neonates at a gestational age of 35 – 37 weeks.

Correspondingly, clearance values increase from about 0.05 l/h in neonates at a gestational age of 27 – 0.2 l/h in neonates at a gestational age of 40 weeks.

5.3 Preclinical safety data

Not applicable.

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Sodium Chloride
2M Sodium Hydroxide
1M Sulphuric Acid
Water for Injections

6.2 Incompatibilities

In general, gentamicin injection should not be mixed. In particular the following are incompatible in mixed solution with gentamicin injection: penicillins, cephalosporins, erythromycin, heparins, sodium bicarbonate. * Dilution in the body will obviate the danger of physical and chemical incompatibility and enable gentamicin to be given concurrently with the drugs listed above either as a bolus injection into the drip tubing, with adequate flushing, or at separate sites. In the case of carbenicillin, administration should only be at a separate site.

*Carbon Dioxide may be liberated on addition of the two solutions. Normally this will dissolve in the solution but under some circumstances small bubbles may form.

6.3 Shelf life

2 years.

6.4 Special precautions for storage

Do not store above 25°C. Do not refrigerate or freeze

6.5 Nature and contents of container

Gentamicin Paediatric Injection is supplied in vials.

6.6 Special precautions for disposal

Not applicable.

7 MARKETING AUTHORISATION HOLDER

Zentiva Pharma UK Limited
12 New Fetter Lane
London
EC4A 1JP
United Kingdom

8 MARKETING AUTHORISATION NUMBER(S)

PL 17780/0507

9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

10/03/2010

10 DATE OF REVISION OF THE TEXT

06/11/2020