

Barcode

some of your Elozar), contact your doctor right away. If you cannot reach your doctor, go to the nearest hospital and take the pack with you.

Patients who have taken too much aripiprazole have experienced the following symptoms:

- rapid heartbeat, agitation/aggressiveness, problems with speech.
- unusual movements (especially of the face or tongue) and reduced level of consciousness.

Other symptoms may include:

- acute confusion, seizures (epilepsy), coma, a combination of fever, faster breathing, sweating,
- muscle stiffness, and drowsiness or sleepiness, slower breathing, choking, high or low blood pressure, abnormal rhythms of the heart.

Contact your doctor or hospital immediately if you experience any of the above.

If you forget to take Elozar
If you miss a dose, take the missed dose as soon as you remember but do not take two doses in one day.

If you stop taking Elozar
Do not stop your treatment just because you feel better. It is important that you carry on taking Elozar for as long as your doctor has told you to.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Common side effects (may affect up to 1 in 10 people):

- diabetes mellitus,
- difficulty sleeping,
- feeling anxious,
- feeling restless and unable to keep still, difficulty sitting still,
- akathisia (an uncomfortable feeling of inner restlessness and a compelling need to move constantly),
- uncontrollable twitching, jerking or writhing movements,
- trembling,
- headache,
- tiredness,
- sleepiness,
- light-headedness,
- shaking and blurred vision,
- decreased number of or difficulty making bowel movements,
- indigestion,
- feeling sick,
- more saliva in mouth than normal,
- vomiting,
- feeling tired.

Uncommon side effects (may affect up to 1 in 100 people):

- increased or decreased blood levels of the hormone prolactin,
- too much sugar in the blood,
- depression,
- altered or increased sexual interest,
- uncontrollable movements of mouth, tongue and limbs (tardive dyskinesia),
- muscle disorder causing twisting movements (dystonia),
- restless legs,
- double vision,
- eye sensitivity to light,
- fast heartbeat,
- a fall in blood pressure on standing up which causes dizziness, light-headedness or fainting,
- hiccups.

The following side effects have been reported since the marketing of oral aripiprazole but the frequency for them to occur is not known:

- low levels of white blood cells,
- low levels of blood platelets,
- allergic reaction (e.g. swelling in the mouth, tongue, face and throat, itching, hives),
- onset or worsening of diabetes, ketoacidosis (ketones in the blood and urine) or coma,
- high blood sugar,
- not enough sodium in the blood,
- loss of appetite (anorexia),
- weight loss,
- weight gain,
- thoughts of suicide, suicide attempt and suicide,
- feeling aggressive,
- agitation,
- nervousness,
- combination of fever, muscle stiffness, faster breathing, sweating, reduced consciousness and sudden changes in blood pressure and heart rate, fainting (neuroleptic malignant syndrome),
- seizure,
- serotonin syndrome (a reaction which may cause feelings of great happiness, drowsiness, clumsiness, restlessness, feeling of being drunk, fever, sweating or rigid muscles),
- speech disorder,
- fixation of the eyeballs in one position,
- sudden unexplained death,
- life-threatening irregular heartbeat,
- heart attack,
- slower heartbeat,
- blood clots in the veins especially in the legs (symptoms include swelling, pain and redness in the leg), which may travel through blood vessels to the lungs causing chest pain and difficulty in breathing (if you notice any of these symptoms, seek medical advice immediately),
- high blood pressure,
- fainting,
- accidental inhalation of food with risk of pneumonia (lung infection),
- spasm of the muscles around the voice box,
- inflammation of the pancreas,
- difficulty swallowing,
- diarrhoea,
- abdominal discomfort,
- stomach discomfort,
- liver failure,
- inflammation of the liver,
- yellowing of the skin and white part of eyes,
- reports of abnormal liver tests values,
- skin rash,
- skin sensitivity to light,

- baldness,
- excessive sweating,
- serious allergic reactions such as Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS). DRESS appears initially as flu-like symptoms with a rash on the face and then with an extended rash, high temperature, enlarged lymph nodes, increased levels of liver enzymes seen in blood tests and an increase in a type of white blood cell (eosinophilia),
- abnormal muscle breakdown which can lead to kidney problems,
- muscle pain,
- stiffness,
- involuntary loss of urine (incontinence),
- difficulty in passing urine,
- withdrawal symptoms in newborn babies in case of exposure during pregnancy,
- prolonged and/or painful erection,
- difficulty controlling core body temperature or overheating,
- chest pain,
- swelling of hands, ankles or feet,
- in blood tests: increased or fluctuating blood sugar, increased glycosylated haemoglobin.
- Inability to resist the impulse, drive or temptation to perform an action that could be harmful to you or others, which may include:
 - strong impulse to gamble excessively despite serious personal or family consequences
 - altered or increased sexual interest and behaviour of significant concern to you or to others, for example, an increased sexual drive
 - uncontrollable excessive shopping
 - binge eating (eating large amounts of food in a short time period) or compulsive eating (eating more food than normal and more than is needed to satisfy your hunger)
 - a tendency to wander away.Tell your doctor if you experience any of these behaviours; he/she will discuss ways of managing or reducing the symptoms.

In elderly patients with dementia, more fatal cases have been reported while taking aripiprazole. In addition, cases of stroke or “mini” stroke have been reported.

Additional side effects in children and adolescents
Adolescents aged 13 years and older experienced side effects that were similar in frequency and type to those in adults except that sleepiness, uncontrollable twitching or jerking movements, restlessness, and tiredness were very common (greater than 1 in 10 patients) and upper abdominal pain, dry mouth, increased heart rate, weight gain, increased appetite, muscle twitching, uncontrolled movements of the limbs, and feeling dizzy, especially when getting up from a lying or sitting position, were common (greater than 1 in 100 patients).

Reporting of side effects
If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects, you can help provide more information on the safety of this medicine.

5. How to store Elozar

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the blister and on the carton after EXP. The expiry date refers to the last day of that month.

This medicinal product does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Elozar contains

- The active substance is aripiprazole.
 - Each orodispersible tablet contains 10 mg of aripiprazole.
 - Each orodispersible tablet contains 15 mg of aripiprazole.
- The other ingredients are lactose monohydrate, cellulose microcrystalline (E460), croscarmellose sodium (E468), silica colloidal anhydrous (E551), magnesium stearate (E470b), purified water, vanilla flavour and aspartame (E951).
Elozar 10 mg orodispersible tablets: iron oxide red (E 172)
Elozar 15 mg orodispersible tablets: iron oxide yellow (E 172)

What Elozar looks like and contents of the pack
Elozar 10 mg orodispersible tablets are round, flat, pink tablets, engraved with ‘10’ on one side and plain on the other.

Elozar 15 mg orodispersible tablets are round, flat, yellow tablets, engraved with ‘15’ on one side and plain on the other.

Elozar orodispersible tablets are supplied in Aluminium-Aluminium paper peel-off blisters packed in cartons containing 14, 28, or 49 orodispersible tablets.

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer
Marketing Authorisation Holder
Novumgen Limited
20-22 Wenlock Road, London,
N1 7GU, United Kingdom

Manufacturer
Rontis Hellas Medical and Pharmaceutical Products S.A.
Larissa Industrial Area, P.O. Box 3012,
Larissa, 41 500,
Greece

This leaflet was last revised in May 2026.



Barcode

Barcode / Artwork Number may be added

	
Product : Elozar 10 mg / 15 mg orodispersible tablets	
Version : V5	Date : May 2026
Sequence Number : 0006	Colors (number & pantone codes) :  Pantone 100% black