

**ARTWORK PANEL**



Master Ref: SI564148  
 Trident Reference No: **TR2840092**  
 SKU Description: Buccastem Buccal Tablets 3mg x 50 UK NZ LFT  
 Brand: Alliance  
 Range: Buccastem  
 Product: Tablets  
 Pack Type: Leaflet  
 Pack Size: 50Tablets  
 Action: **A**  
 Date: 18/04/23  
 Country: UK

CAD Ref: PIR\_HEALTH\_PIL\_140 x 352mm  
 Manufacturing Site:  
 Printer:  
 Substrate: 50gsm ±5gsm Bond C Opaque printing paper

Barcode Type: N/A  
 Magnification: N/A  
 Pharmacode No/NE: 001010010(593)

Font Size (Main Body Copy): 7.00pt

Technical & Non Printing Items  
 Cutter Guides

Colours  
 Process Black  
 1  
 LT  
 H



Please note that any low resolution paper Canon colour copies associated with this job should be referred to for content, layout and colour separation only.

**UNDER NO CIRCUMSTANCES SHOULD THIS ARTWORK BE ALTERED WITHOUT PRIOR PERMISSION FROM TRIDENT.**

**STUDIO USE ONLY** v6.0  
 Sm\*Art check results: G=0; O=231; R=0; - JW - 18/04/23 17:20:59

Package leaflet: Information for the user ALLIANCE

## Buccastem® 3mg Buccal Tablets

(PROCHLORPERAZINE MALEATE)

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

**What is in this leaflet**

1. What Buccastem Tablets are and what they are used for
2. What you need to know before you take Buccastem Tablets
3. How to take Buccastem Tablets
4. Possible side effects
5. How to store Buccastem Tablets
6. Contents of the pack and other information

**1. What Buccastem Tablets are and what they are used for**

Buccastem Tablets are effective in treating nausea (feeling sick) and vomiting (being sick) from whatever cause. They are also used to treat migraine and dizziness due to ear problems and other causes. Buccastem Tablets contain prochlorperazine maleate. Prochlorperazine belongs to a large group of drugs known as phenothiazines, which have a variety of effects.

**2. What you need to know before you take Buccastem Tablets**

You should only take Buccastem Tablets when your doctor prescribes them for you.

**Do not take Buccastem Tablets:**

- if you are allergic to prochlorperazine maleate or any of the other ingredients of this medicine (listed in section 6)
- if you have problems with your liver
- if you have blood problems
- if you suffer from epilepsy, Parkinson's Disease or glaucoma
- if you have problems with your prostate gland.

**Warnings and precautions**

**Talk to your doctor before taking Buccastem Tablets:**

- if you are elderly
- if you were born with or have a family history of prolonged QT interval, or if you have acquired QT prolongation (seen on ECG, electrical recording of the heart)
- if you have heart disorders or have a history of heart attack (myocardial infarction)
- if you have salt imbalance in the blood (especially low level of potassium or magnesium)
- if you are pregnant, thinking of becoming pregnant or breast feeding
- if you have risk factors for a blood clot or stroke such as high blood pressure, high cholesterol levels, diabetes or smoking
- if you or someone else in your family has a history of blood clots, as medicines like these have been associated with formation of blood clots
- if you have a condition called myasthenia gravis, which causes muscle weakness
- if you have AIDS.

Buccastem Tablets may cause you to be more sensitive to sunlight. When taking Buccastem Tablets it is particularly important that you should avoid exposure to direct sunlight and use sunscreen.

**Other medicines and Buccastem Tablets**

Tell your doctor if you are taking, have recently taken or might take any other medicines. In particular, tell your doctor or pharmacist if you are taking:

- medicines known to affect the rhythm of your heart. This includes medicines used for abnormal heart rhythm (antiarrhythmics), for depression (tricyclic antidepressants) and for psychiatric disorders (antipsychotics).
- azithromycin, clarithromycin, or erythromycin (antibiotics used for treating infections). Taking these medicines at the same time as Buccastem Tablets may increase the chance of you getting side effects that affect your heart.
- sedatives or tranquilisers, used to help you sleep (such as diazepam and temazepam)
- antihypertensive medicines, used to treat high blood pressure (such as atenolol, propranolol or clonidine)
- anticholinergic medicines, used for conditions such as depression (such as amitriptyline and imipramine)
- anticonvulsant medicines, used to prevent or treat fits (such as carbamazepine and lamotrigine)
- lithium (used to treat mania and depression)
- desferrioxamine (used to treat high levels of iron in your body) .

**Buccastem Tablets with food and drink**

Buccastem Tablets are best taken **after food**. Do not drink alcohol when taking the tablets as it may interact with medicines like Buccastem.

**Pregnancy and breast feeding**

If you are pregnant or thinking of becoming pregnant, you should only take Buccastem Tablets on your doctor's instructions.

The following symptoms may occur in newborn babies of mothers that have used Buccastem Tablets in the last trimester (last three months of their pregnancy): shaking, muscle stiffness and/or weakness, sleepiness, agitation, breathing problems, and difficulty in feeding. If your baby develops any of these symptoms you may need to contact your doctor.

Experience with Buccastem Tablets is limited. You should therefore not take the tablets if you are breast-feeding.

Ask your doctor or pharmacist for advice before taking any medicine.

**Driving and using machines**

Buccastem Tablets can make you feel drowsy. Therefore you should avoid driving or using dangerous machines until you know how the tablets affect you.

**Buccastem Tablets contain sucrose**



If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

**3. How to take Buccastem Tablets**

Always take this medicine exactly as your doctor has told you. Check with your doctor if you are not sure. The **recommended dose** is one or two tablets twice a day for adults and children over 12 years of age. Buccastem Tablets are **not recommended** for children **under 12 years of age**.

**INSTRUCTIONS FOR USE**

Please read carefully before taking the tablet(s).

Pharmacode Reads This Way  
 First Bar  
 Last Bar

Pharmacode Reads This Way  
 First Bar  
 Last Bar

352mm

140mm

## ARTWORK PANEL

# ALLIANCE

Master Ref: S1564148  
 Trident Reference No: **TR2840092**  
 SKU Description: Buccastem Buccal Tablets 3mg x 50 UK NZ LFT  
 Brand: Alliance  
 Range: Buccastem  
 Product: Tablets  
 Pack Type: Leaflet  
 Pack Size: 50 Tablets  
 Action: **A**  
 Date: 18/04/23  
 Country: UK

CAD Ref: PIR\_HEALTH\_PIL\_140 x 352mm  
 Manufacturing Site:  
 Printer:  
 Substrate: 50gsm ±5gsm Bond C Opaque printing paper

Barcode Type: N/A  
 Magnification: N/A  
 Pharmacode No/NE: 001010010(593)

Font Size (Main Body Copy): 7.00pt

Technical & Non Printing Items  
 Cutter Guides

Colours  
 Process Black  
 1  
 LT  
 H



Connaught House, Connaught Road,  
 Kingswood Business Park, Hull, HU7 3AP, England.  
 T: +44 (0) 1482 828100

Please note that any low resolution paper Canon colour copies associated with this job should be referred to for content, layout and colour separation only.

UNDER NO CIRCUMSTANCES SHOULD THIS ARTWORK BE ALTERED WITHOUT PRIOR PERMISSION FROM TRIDENT.

STUDIO USE ONLY v6.0  
 Sm\*Art check results: G=0; O=231; R=0; - JW - 18/04/23 17:20:59

- Place the tablet high up along your top gum, under the upper lip either side of your mouth as indicated above. The tablet **must not** be swallowed whole or chewed.
- The tablet will soften and adhere to the gum. Allow it to dissolve slowly and completely - this may take between 1 and 2 hours. Most people find that after a few minutes they no longer notice the tablet.
- The tablet **should not** be moved about the mouth with the tongue as this will cause it to dissolve too quickly.
- If you wear dentures, the tablet may be placed in any comfortable position between your lip and gum. The tablet(s) is best taken **after** meals.

#### If you take more Buccastem Tablets than you should

If you accidentally take too many tablets you **must** seek medical attention immediately. Show any left-over medicines or the empty packet to the doctor.

#### If you forget to take Buccastem Tablets

If you forget to take a dose do not double the dose next time. Just carry on taking the medicine as the doctor has told you.

### 4. Possible side effects

Like all medicines, Buccastem Tablets can cause side effects, although not everybody gets them.

#### Serious side effects

If you experience any of the following side effects, you should **stop taking Buccastem Tablets and contact your doctor immediately**:

- Symptoms of an allergic reaction such as rash and swelling of the face, tongue or throat.
- A high temperature, pale complexion, muscle stiffness and changes in levels of alertness, you may have developed a serious condition called neuroleptic malignant syndrome.
- Convulsions (fits).
- Signs of infection such as shivering, headache, sweating, high temperature, flushing, a sore throat or mouth and swollen glands.
- Very fast, uneven or forceful heartbeats (palpitations) and breathing problems such as wheezing, shortness of breath, tightness in the chest and chest pain.
- Blood clots in the veins especially in the legs (symptoms include swelling, pain and redness in the leg), which may travel through blood vessels to the lungs causing chest pain and difficulty in breathing.
- Symptoms of decreased sodium concentration in the blood, such as nausea and vomiting, headache, confusion, loss of energy, restlessness, muscle weakness, spasms or cramps, seizures and coma. In elderly people with dementia, a small increase in the number of deaths has been reported for patients taking medicines of the same type as Buccastem Tablets compared with those not taking these medicines.

#### Other side effects include:

##### Rare: may affect up to 1 in 1,000 people

- jaundice
- blood problems.

##### Very rare: may affect up to 1 in 10,000 people

- breast swelling (in men as well as in women)
- milk secretion from the breasts which is not due to breast-feeding
- absence of menstrual periods

##### Unknown: frequency cannot be estimated from the available data

- drowsiness
- dizziness
- dry mouth
- inability to sleep (insomnia)
- agitation
- a feeling of restlessness or anxiety that can range from mild to severe
- skin reactions
- increased sensitivity to sunlight
- low blood pressure (this makes you feel dizzy or faint, particularly when you stand up), particularly in elderly or volume depleted patients (those who have lost both water and salts from the body)
- high blood sugar level, which can cause increased thirst, a dry mouth, needing to urinate frequently, tiredness and recurrent infections such as thrush
- a condition known as cholestasis where bile cannot flow properly in the body, and can cause itchiness, jaundice (yellowing of the skin and/or the whites of the eyes), pale stools and dark urine
- abnormal movements, tremors and muscle rigidity, and unusual movements of the face and tongue
- impotence, ejaculation problems and persistent/ painful erections
- local irritation to the gum and mouth
- numbness of mouth
- tingling, pricking or burning of mouth
- altered or decreased ability to taste.

#### Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. In the UK, you can also report side effects directly via the Yellow Card Scheme at [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

### 5. How to store Buccastem Tablets

Keep this medicine out of the sight and reach of children.  
 Keep the blisters in the outer carton in order to protect from light.  
 This medicinal product does not require any special temperature storage condition.  
**Do not use** after the expiry date (EXP month/year) shown on the outer carton and blister. The expiry date refers to the last day of that month.

### 6. Contents of the pack and other information

#### What Buccastem Tablets contain

Each tablet contains 3 mg of the active ingredient prochlorperazine maleate.  
 The other ingredients are: compressible sugar (sucrose), povidone K30, xanthan gum, locust bean gum, talc, magnesium stearate and riboflavin sodium phosphate.

#### What Buccastem Tablets look like and contents of the pack

The buccal tablet is pale yellow and has J1 on one side and is plain on the other.  
 The product is only available on prescription in packs of 50 tablets.

#### Marketing Authorisation Holder and Manufacturer

Marketing Authorisation Holders: Alliance Pharmaceuticals Limited, Avonbridge House, Bath Road, Chippenham, Wiltshire, SN15 2BB, UK.  
 Manufacturer: Piramal Healthcare UK Limited, Whalton Road, Morpeth, NE61 3YA, United Kingdom.  
 This leaflet was last revised in March 2023.  
 Buccastem is a registered trademark of Alliance Pharmaceuticals Limited.  
 Alliance and associated devices are registered trademarks of Alliance Pharmaceuticals Limited.  
 © Alliance Pharmaceuticals Limited 2023

TR2840092  
 20769353

140mm

352mm