

Package leaflet: Information for the user

Risedronate sodium 30 mg film-coated tablets

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Risedronate sodium 30 mg is and what it is used for
2. What you need to know before you take Risedronate sodium 30 mg
3. How to take Risedronate sodium 30 mg
4. Possible side effects
5. How to store Risedronate sodium 30 mg
6. Contents of the pack and other information

1. What Risedronate sodium 30 mg is and what it is used for

Risedronate sodium 30 mg belongs to a group of non-hormonal medicines called bisphosphonates which are used to treat bone diseases. It works directly on your bones to make them stronger and therefore less likely to break.

Risedronate sodium 30 mg is used to treat Paget's disease of the bone (osteitis deformans).

Bone is a living tissue. Old bone is constantly removed from your skeleton and replaced with new bone. Paget's disease occurs when this process, called remodelling, happens too quickly and in a disordered way. The new bone that is produced is weaker than normal and the affected bones may become enlarged, painful and may fracture.

Risedronate sodium 30 mg changes the bone remodelling process back to normal, returning the strength to the bone structure.

2. What you need to know before you take Risedronate sodium 30 mg

Do not take Risedronate sodium 30 mg:

- if you are allergic to risedronate sodium or any of the other ingredients of this medicine (listed in section 6);
- if your doctor has told you that you have a condition called hypocalcaemia (a low blood calcium level);
- if you may be pregnant, are pregnant or are planning to become pregnant;
- if you are breast-feeding;
- if you have severe kidney problems.

Warnings and precautions

Talk to your doctor or pharmacist before taking Risedronate sodium 30 mg

- if you are unable to stay in an upright position (sitting or standing) for at least 30 minutes;
- if you have abnormal bone and mineral metabolism (for example lack of vitamin D, parathyroid hormone abnormalities, both leading to a low blood calcium level);
- if you have or have had problems in the past with your oesophagus (the tube that connects your mouth with your stomach). For instance you may have or have had pain or difficulty in swallowing food, or you have previously been told that you have Barrett's oesophagus (a condition associated with changes in the cells that line the lower oesophagus);
- if you have been told by your doctor that you have an intolerance to some sugars (such as lactose);
- if you have or ever had pain, swelling or numbness of the jaw or a "heavy jaw feeling" or loosening of a tooth;
- if you are under dental treatment or will undergo dental surgery, tell your dentist that you are being treated with Risedronate sodium 30 mg.

Your doctor will advise you on what to do when taking Risedronate sodium 30 mg if you have any of the above.

Children and adolescents

Risedronate sodium 30 mg tablets should not be taken by children under 18 years of age.

Other medicines and Risedronate sodium 30 mg

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Medicines containing any of the following lessen the effect of Risedronate sodium 30 mg if taken at the same time:

- calcium;
- magnesium;
- aluminium (for example some indigestion mixtures);
- iron.

Take these medicines at least 30 minutes after your Risedronate sodium 30 mg tablet.

Risedronate sodium 30 mg with food and drink

It is very important that you **do not** take your Risedronate sodium 30 mg tablet with food or drinks (other than plain water) so that it can work properly. Products containing any of the following: calcium (including dairy products such as milk), magnesium, aluminium (for example, some indigestion mixtures) or iron lessen the effect of this medicine. (see section 2, "Other medicines and Risedronate sodium 30 mg").

Take food and drinks (other than plain water) at least 30 minutes after your Risedronate sodium 30 mg tablet.

Pregnancy and breast-feeding

Do not take Risedronate sodium 30 mg if you are pregnant, think you may be pregnant or are planning to have a baby (See section 2, “Do not take Risedronate sodium 30 mg”). Ask your doctor or pharmacist for advice before taking this medicine.

The potential risk associated with the use of risedronate sodium in pregnant women is unknown.

Do not take Risedronate sodium 30 mg if you are breast-feeding (see section 2, “Do not take Risedronate sodium 30 mg”).

Ask your doctor for advice before taking any medicine.

Driving and using machines

Risedronate sodium 30 mg is not known to affect your ability to drive and use machines.

Risedronate sodium 30 mg contains lactose

Risedronate sodium 30 mg contains a small amount of lactose.

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking Risedronate sodium 30 mg.

3. How to take Risedronate sodium 30 mg

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

Recommended dose:

Take one Risedronate sodium 30 mg tablet once a day.

Usually 2 months will be enough to treat your Paget’s disease. However, another course of treatment may be required.

It is very important that you **do not** take Risedronate sodium 30 mg with food or drinks (other than plain water) so that it can work properly.

When to take the Risedronate sodium 30 mg

It is best to take your tablet at least 30 minutes before the first food, drink (other than plain water) or other medicine of the day.

If you are unable to take your Risedronate sodium 30 mg tablet at this time, you may take it on an empty stomach, at the same time every day, in one of the following ways:

- Between meals: at least 2 hours after your last food, drink (other than plain water) or medicine. Do not eat or drink (other than plain water) for 2 hours after taking the tablet.
- In the evening: at least 2 hours after the last food, drink (other than plain water) or medicine of the day. Risedronate sodium 5 mg should be taken at least 30 minutes before going to bed.

How to take the Risedronate sodium 30 mg

- Take the tablet whilst you are in an upright position (you may sit or stand) to avoid heartburn.
- Swallow it with at least one glass (120 ml) of plain water.

- Swallow it whole. Do not suck or chew it.
- Do not lie down for 30 minutes after taking your tablet.

Your doctor will tell you if you need calcium and vitamin supplements, if you are not taking enough from your diet.

If you take more Risedronate sodium 30 mg than you should

If you or somebody else has accidentally taken more Risedronate sodium 30 mg tablets than prescribed, drink one full glass of milk and seek medical attention.

If you forget to take Risedronate sodium 30 mg

If you have forgotten to take your tablet, you can take it before breakfast, between meals, or in the evening according to the instructions above.

Do not take two tablets in one day to make up for the tablet you missed.

If you stop taking Risedronate sodium 30 mg

Do not stop the treatment prematurely because Paget's disease is a long-term disease. Please talk to your doctor before you consider stopping treatment.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Stop taking Risedronate sodium 30 mg and contact a doctor immediately if you experience any of the following:

- Symptoms characteristic of severe tissue swelling (angioedema reaction):
 - Swelling of face, tongue or throat
 - Difficulties in swallowing
 - Hives and difficulties in breathing.
- Severe skin reactions that can include blistering of the skin.

Tell your doctor promptly if you experience the following side effects:

- Eye inflammation, usually with pain, redness and light sensitivity.
- Bone necrosis of the jaw (osteonecrosis) associated with delayed healing and infection, often following tooth extraction (see section 2, "Warnings and precautions").
- Symptoms from the oesophagus such as pain when you swallow, difficulties in swallowing, chest pain or new or worsened heartburn.

Unusual fracture of the thigh bone particularly in patients on long-term treatment for osteoporosis may occur rarely. Contact your doctor if you experience pain, weakness or discomfort in your thigh, hip or groin as this may be an early indication of a possible fracture of the thigh bone.

However, in clinical studies, the other side effects that were observed were usually mild and did not cause the patient to stop taking their tablets.

Common side effects (may affect up to 1 in 10 people)

- Indigestion, feeling sick, stomach ache, stomach cramps or discomfort, constipation, feelings of fullness, bloating, diarrhoea;
- Pain in your bones, muscles or joints;
- Headache.

Uncommon side effects (may affect up to 1 in 100 people)

- Inflammation or ulcer of the oesophagus (the tube that connects your mouth with your stomach) causing difficulty and pain in swallowing (see also section 2, “Warning and precautions”), inflammation of the stomach and duodenum (bowel draining the stomach);
- Inflammation of the coloured part of the eye (iris) (red, painful eyes with a possible change in vision).

Rare side effects (may affect up to 1 in 1,000 people)

- Inflammation of the tongue (red, swollen, possibly painful), narrowing of the oesophagus (the tube that connects your mouth with your stomach);
- Abnormal liver tests have been reported. These can only be diagnosed from a blood test.

Very rare side effects (may affect up to 1 in 10,000 people)

Talk to your doctor if you have ear pain, discharge from the ear, and/or an ear infection. These could be signs of bone damage in the ear.

During post-marketing experience, the following have been reported (unknown frequency):

- Hair loss;
- Liver disorders, some cases were severe.

Rarely, at the beginning of treatment, a patient’s blood calcium and phosphate levels may fall. These changes are usually small and cause no symptoms.

The additional following adverse events has also been observed in a clinical study in patients with Paget’s disease: vision difficulties, breathing difficulties, coughing, inflammation of the large intestine, surface of the eye damage, cramps, dizziness, dryness of the eye, flu-like symptoms, muscle weakness, abnormal growth of cells, a frequent need to pass water at night, unusual lumps or swellings, chest pain, runny nose, ringing in the ears and weight loss.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store.

By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Risedronate sodium 30 mg

Keep this medicine out of the sight and reach of children.
Do not use this medicine after the expiry date which is stated on the carton after EXP.
The expiry date refers to the last day of that month.

No special storage conditions required.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Risedronate sodium 30 mg contains

The active substance is risedronate sodium. Each tablet contains 30 mg risedronate sodium, equivalent to 27.8 mg risedronic acid.

The other ingredients are:

Tablet core: lactose monohydrate, crospovidone, magnesium stearate and microcrystalline cellulose.

Film coating: hypromellose, macrogol, hydroxypropyl cellulose, colloidal anhydrous silica and titanium dioxide (E171).

What Risedronate sodium 30 mg looks like and contents of the pack

Risedronate sodium 30 mg film-coated tablets are white to off-white coloured tablets debossed with the letter "J" on one side and "30" on the other.

The tablets are supplied in blister packs of 28 or 30 tablets.

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer

Marketing Authorisation Holder:

Waymade Plc trading as Sovereign Medical
Sovereign House,
Miles Gray Road,
Basildon, Essex SS14 3FR,
United Kingdom.

Manufacturer:

Waymade Plc
Sovereign House,
Miles Gray Road,
Basildon, Essex SS14 3FR,
United Kingdom.

This leaflet was last revised in October 2017

To request a copy of this leaflet in Braille, large print or audio format, contact the licence holder at the above address or telephone: 01268 535200 (select option Medical Information) / e-mail: info@waymade.co.uk