

# SUMMARY OF PRODUCT CHARACTERISTICS

## 1 NAME OF THE MEDICINAL PRODUCT

Metoclopramide 10mg Tablets

## 2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each tablet contains Metoclopramide Hydrochloride Monohydrate, equivalent to 10mg metoclopramide hydrochloride.

Excipient with known effects: Lactose monohydrate.

Each tablet contains 85mg Lactose monohydrate.

For the full list of excipients, see section 6.1.

## 3 PHARMACEUTICAL FORM

Tablet.

White, circular, biconvex uncoated tablets impressed “C” on one face and the identifying letters “M” and “P” on either side of a central division line on the reverse.

The score line is only to facilitate breaking for ease of swallowing and not to divide into equal doses.

## 4 CLINICAL PARTICULARS

### 4.1 Therapeutic indications

#### Adult population

Metoclopramide is indicated in adults for:

- Prevention of delayed chemotherapy induced nausea and vomiting (CINV).
- Prevention of radiotherapy induced nausea and vomiting (RINV).
- Symptomatic treatment of nausea and vomiting, including acute migraine induced nausea and vomiting. Metoclopramide can be used in combination with oral analgesics to improve the absorption of analgesics in acute migraine.

#### Paediatric population

Metoclopramide 10mg tablets are indicated in children (aged 15-18 years) for:  
- Prevention of delayed chemotherapy induced nausea and vomiting (CINV) as a second line option.

## 4.2 Posology and method of administration

### Posology

#### *All indications (adult patients)*

The recommended single dose is 10 mg, repeated up to three times daily.  
The maximum recommended daily dose is 30 mg or 0.5mg/kg body weight.  
The maximum recommended treatment duration is 5 days.

#### *Paediatric population*

Metoclopramide is contraindicated in children aged less than 1 year (see section 4.3).

#### *Paediatric patients aged 15-18 years*

Prevention of delayed chemotherapy induced nausea and vomiting (CINV)  
The recommended dose is 0.1 to 0.15 mg/kg body weight, repeated up to three times daily by oral route. The maximum dose in 24 hours is 0.5 mg/kg body weight.

#### Dosing table

Age	Body Weight	Dose	Frequency
15-18 years	Over 60kg	10mg	Up to 3 times daily

The maximum treatment duration is 5 days for prevention of delayed chemotherapy induced nausea and vomiting (CINV).

Tablets are not suitable for use in children weighing less than 61kg. Other pharmaceutical forms/strengths may be more appropriate for administration to this population.

A minimal interval of 6 hours between two administrations is to be respected, even in case of vomiting or rejection of the dose (see section 4.4).

#### Special population

##### *Elderly*

In elderly patients a dose reduction should be considered, based on renal and hepatic function and overall frailty.

##### *Renal impairment*

In patients with end stage renal disease (Creatinine clearance  $\leq$  15 ml/min), the daily dose should be reduced by 75%. In patients with moderate to severe renal impairment (Creatinine clearance 15-60 ml/min), the dose should be reduced by 50% (see section 5.2).

##### *Hepatic impairment*

In patients with severe hepatic impairment, the dose should be reduced by 50% (see section 5.2).

Other pharmaceutical forms/strengths may be more appropriate for administration to these populations.

#### Method of administration

For oral use only.

### **4.3 Contraindications**

- Hypersensitivity to the active substance or to any of the excipients listed in section 6.1
- Gastrointestinal haemorrhage, mechanical obstruction or gastro-intestinal perforation for which the stimulation of gastrointestinal motility constitutes a risk
- Confirmed or suspected pheochromocytoma, due to the risk of severe hypertension episodes
- History of neuroleptic or metoclopramide-induced tardive dyskinesia
- Epilepsy (increased crises frequency and intensity)
- Parkinson's disease
- Combination with levodopa or dopaminergic agonists (see section 4.5)
- Known history of methaemoglobinaemia with metoclopramide or of NADH cytochrome-b5 deficiency
- Use in children less than 1 year of age due to an increased risk of extrapyramidal disorders (see section 4.4).

Metoclopramide should not be used during the first three to four days following operations such as pyloroplasty or gut anastomosis as vigorous muscular contractions may not help healing.

### **4.4 Special warnings and precautions for use**

Precautions:

If vomiting persists the patient should be reassessed to exclude the possibility of an underlying disorder e.g. cerebral irritation.

Neurological Disorders

Extrapyramidal disorders may occur, particularly in children and young adults, and/or when high doses are used. These reactions occur usually at the beginning of the treatment and can occur after a single administration. Metoclopramide should be discontinued immediately in the event of extrapyramidal symptoms. These effects are generally completely reversible after treatment discontinuation, but may require a symptomatic treatment (benzodiazepines in children and/or anticholinergic anti-Parkinsonian medicinal products in adults).

The time interval of at least 6 hours specified in the section 4.2 should be respected between each metoclopramide administration, even in case of vomiting and rejection of the dose, in order to avoid overdose.

Prolonged treatment with metoclopramide may cause tardive dyskinesia, potentially irreversible, especially in the elderly. Treatment should not exceed 3 months because of the risk of tardive dyskinesia (see section 4.8). Treatment must be discontinued if clinical signs of tardive dyskinesia appear.

Neuroleptic malignant syndrome has been reported with metoclopramide in combination with neuroleptics as well as with metoclopramide monotherapy (see section 4.8). Metoclopramide should be discontinued immediately in the event of symptoms of neuroleptic malignant syndrome and appropriate treatment should be initiated.

Special care should be exercised in patients with underlying neurological conditions and in patients being treated with other centrally-acting drugs (see section 4.3)

Symptoms of Parkinson's disease may also be exacerbated by metoclopramide.

#### Methaemoglobinaemia

Methemoglobinaemia which could be related to NADH cytochrome b5 reductase deficiency has been reported. In such cases, metoclopramide should be immediately and permanently discontinued and appropriate measures initiated (such as treatment with methylene blue).

#### Cardiac Disorders

There have been reports of serious cardiovascular undesirable effects including cases of circulatory collapse, severe bradycardia, cardiac arrest and QT prolongation following administration of metoclopramide by injection, particularly via the intravenous route (see section 4.8).

Special care should be taken when administering metoclopramide, particularly via the intravenous route to the elderly population, to patients with cardiac conduction disturbances (including QT prolongation), patients with uncorrected electrolyte imbalance, bradycardia and those taking other drugs known to prolong QT interval.

Intravenous doses should be administered as a slow bolus (at least over 3 minutes) in order to reduce the risk of adverse effects (e.g. hypotension, akathisia).

#### Renal and Hepatic Impairment

In patients with renal impairment or with severe hepatic impairment, a dose reduction is recommended (see section 4.2).

Metoclopramide may cause elevation of serum prolactin levels.

#### Contains lactose:

Patients with rare hereditary problems of galactose intolerance, total lactase deficiency or glucose-galactose malabsorption should not take this medicine.

Care should be exercised when using Metoclopramide in patients with a history of atopy (including asthma) or porphyria. Metoclopramide should not be used in the immediate post-operative period (up to 3-4 days) following pyloroplasty or gut anastomosis, as vigorous gastrointestinal contractions may adversely affect healing.

Special care should be taken when administering Metoclopramide intravenously to patients with 'sick sinus syndrome' or other cardiac conduction disturbances. There have been very rare reports of abnormalities of cardiac conduction with intravenous metoclopramide. Metoclopramide should be used with care with other drugs affecting cardiac conduction.

#### **4.5 Interaction with other medicinal products and other forms of interaction**

##### **Contraindicated combination**

Levodopa or dopaminergic agonists and metoclopramide have a mutual antagonism (see section 4.3).

##### **Combination to be avoided**

Alcohol potentiates the sedative effect of metoclopramide.

##### **Combination to be taken into account**

Due to the prokinetic effect of metoclopramide, the absorption of certain drugs may be modified.

##### *Anticholinergics and morphine derivatives*

Anticholinergics and morphine derivatives may have both a mutual antagonism with metoclopramide on the digestive tract motility.

##### *Central nervous system depressants (morphine derivatives, anxiolytics, sedative H1 antihistamines, sedative antidepressants, barbiturates, clonidine and related)*

Sedative effects of Central Nervous System depressants and metoclopramide are potentiated.

##### *Neuroleptics*

Metoclopramide may have an additive effect with other neuroleptics on the occurrence of extrapyramidal disorders.

##### *Serotonergic drugs*

The use of metoclopramide with serotonergic drugs such as SSRIs may increase the risk of serotonin syndrome.

##### *Digoxin*

Metoclopramide may decrease digoxin bioavailability. Careful monitoring of digoxin plasma concentration is required.

##### *Cyclosporine*

Metoclopramide increases cyclosporine bioavailability (C<sub>max</sub> by 46% and exposure by 22%). Careful monitoring of cyclosporine plasma concentration is required. The clinical consequence is uncertain.

*Mivacurium and suxamethonium*

Metoclopramide injection may prolong the duration of neuromuscular block (through inhibition of plasma cholinesterase).

*Strong CYP2D6 inhibitors*

Metoclopramide exposure levels are increased when co-administered with strong CYP2D6 inhibitors such as fluoxetine and paroxetine. Although the clinical significance is uncertain, patients should be monitored for adverse reactions.

Metoclopramide may reduce plasma concentrations of atovaquone.

#### **4.6 Fertility, pregnancy and lactation**

*Pregnancy*

A large amount of data on pregnant women (more than 1000 pregnancy outcomes) indicate no malformative nor feto/neonatal toxicity of Metoclopramide hydrochloride. Metoclopramide can be used during pregnancy if clinically needed. Due to pharmacological properties (as other neuroleptics), in case of metoclopramide administration at the end of pregnancy, extrapyramidal syndrome in newborn cannot be excluded.

Metoclopramide should be avoided at the end of pregnancy. If metoclopramide is used, neonatal monitoring should be undertaken.

*Breast-feeding*

Metoclopramide is excreted in breast milk at low level. Adverse reactions in the breast-fed baby cannot be excluded. Therefore metoclopramide is not recommended during breast-feeding. Discontinuation of metoclopramide in breast-feeding women should be considered.

*Fertility*

No data available.

#### **4.7 Effects on Ability to Drive and Use Machines**

Metoclopramide may cause drowsiness, dizziness, dyskinesia and dystonias which could affect the vision and also interfere with the ability to drive and operate machinery.

#### **4.8 Undesirable effects**

Adverse reactions listed by System Organ Class. Frequencies are defined using the following convention: very common ( $\geq 1/10$ ), common ( $\geq 1/100$  to  $< 1/10$ ), uncommon ( $\geq 1/1,000$  to  $< 1/100$ ), rare ( $\geq 1/10,000$  to  $< 1/1,000$ ), very rare ( $< 1/10,000$ ), not known (cannot be estimated from the available data).

<b>System Organ Class</b>	<b>Frequency</b>	<b>Adverse reactions</b>
<b>Blood and lymphatic system disorders</b>	Not known	Methaemoglobinaemia, which could be related to NADH cytochrome b5 reductase deficiency, particularly in neonates (see section 4.4); Sulphaemoglobinaemia, mainly with concomitant administration of high doses of sulfur-releasing medicinal products
<b>Immune system disorders</b>	Uncommon	Hypersensitivity
	Not known	Anaphylactic reaction (including anaphylactic shock particularly with intravenous formulation)
<b>Endocrine disorders*</b>	Uncommon	Amenorrhoea; Hyperprolactinaemia,
	Rare	Galactorrhoea
	Not known	Gynaecomastia
<b>Psychiatric disorders</b>	Common	Depression
	Uncommon	Hallucination
	Rare	Confusional state
<b>Nervous system disorders</b>	Very common	Somnolence
	Common	Extrapyramidal disorders (particularly in children and young adults and/or when the recommended dose is exceeded, even following administration of a single dose of the drug) (see section 4.4); Parkinsonism; Akathisia
	Uncommon	Dystonia (including visual disturbances and oculogyric crisis); Dyskinesia; Depressed level of consciousness
	Rare	Convulsion especially in epileptic patients
	Not known	Tardive dyskinesia which may be persistent, during or after prolonged treatment, particularly in elderly patients (see section 4.4); Neuroleptic malignant syndrome (see section 4.4)
<b>Cardiac disorders</b>	Uncommon	Bradycardia, particularly with intravenous formulation
	Not known	Cardiac arrest, occurring shortly

		after injectable use, and which can be subsequent to bradycardia (see section 4.4); Atrioventricular block; Sinus arrest particularly with intravenous formulation; Electrocardiogram QT prolonged; Torsade de Pointes
<b>Vascular disorders</b>	Common	Hypotension, particularly with intravenous formulation
	Not known	Shock; Syncope after injectable use; Acute hypertension in patients with phaeochromocytoma (see section 4.3); Transient increase in blood pressure
<b>Gastrointestinal disorders</b>	Common	Diarrhoea
<b>General disorders and administration site conditions</b>	Common	Asthenia

\* Endocrine disorders during prolonged treatment in relation with hyperprolactinaemia (amenorrhoea, galactorrhoea, gynaecomastia).  
The following reactions, sometimes associated, occur more frequently when high doses are used:

- Extrapyramidal symptoms: acute dystonia and dyskinesia, parkinsonian syndrome, akathisia, even following administration of a single dose of the medicinal product, particularly in children and young adults (see section 4.4).
- Drowsiness, decreased level of consciousness, confusion, hallucination.

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the Yellow Card Scheme website: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store.

#### **4.9 Overdose**

##### Symptoms

Extrapyramidal disorders, drowsiness, decreased level of consciousness, confusion, hallucination, and cardio-respiratory arrest may occur.

##### Management

In case of extrapyramidal symptoms related or not to overdose, the treatment is only symptomatic (benzodiazepines in children and/or anticholinergic anti-parkinsonian medicinal products in adults). A symptomatic treatment and a

continuous monitoring of the cardiovascular and respiratory functions should be carried out according to clinical status.

## **5 PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Drugs for functional gastrointestinal disorders, propulsives  
ATC Code: A03FA01

#### Mechanism of action

The action of metoclopramide is closely associated with parasympathetic nervous control of the upper gastro-intestinal tract where it has the effect of encouraging normal peristaltic action. This provides for a fundamental approach to the control of those conditions where disturbed gastrointestinal motility is a common underlying factor.

### **5.2 Pharmacokinetic properties**

Metoclopramide is metabolised in the liver and the predominant route of elimination of metoclopramide and its metabolites is via the kidney.

#### Renal impairment

The clearance of metoclopramide is reduced by up to 70% in patients with severe renal impairment, while the plasma elimination half-life is increased (approximately 10 hours for a creatinine clearance of 10-50 mL/minute and 15 hours for a creatinine clearance <10 mL/minute).

#### Hepatic impairment

In patients with cirrhosis of the liver, accumulation of metoclopramide has been observed, associated with a 50% reduction in plasma clearance.

### **5.3 Preclinical safety data**

No additional data available.

## **6 PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Also contains:

Colloidal silica  
Lactose monohydrate  
Magnesium stearate  
Maize starch  
Microcrystalline cellulose (E460)

### **6.2. Incompatibilities**

None known.

### **6.3 Shelf-life**

Four years from the date of manufacture (PVC blister packs).

Three years from the date of manufacture (polypropylene containers).

### **6.4 Special Precautions for Storage**

Store below 25°C in a dry place. Protect from light.

### **6.5 Nature and contents of container**

The product containers are rigid injection moulded polypropylene containers with snap-on polyethylene lids.

Pack size: 1000s

The product may also be supplied in blister packs and cartons:

a) Carton: Printed carton manufactured from white folding box board.

b) Blister pack: (i) 250µm white rigid PVC. (ii) Surface printed 20µm hard temper aluminium foil.

Pack sizes: 28s, 30s, 56s, 60s, 84s, 90s, 100s, 112s, 120s, 168s, 180s, 250s, 500s, 1000s

Not all pack sizes may be marketed.

### **6.6 Special precautions for disposal and other handling**

Not applicable.

## **7 MARKETING AUTHORISATION HOLDER**

Accord-UK Ltd  
(Trading style: Accord)  
Whiddon Valley  
Barnstaple  
Devon  
EX32 8NS

**8      MARKETING AUTHORISATION NUMBER(S)**

PL 0142/0252

**9      DATE OF FIRST AUTHORISATION/RENEWAL OF THE  
AUTHORISATION**

8.7.87

Renewed: 8.7.92, 8.7.97, 8.7.02

**10     DATE OF REVISION OF THE TEXT**

08/07/2023