

Resolor® 2 mg film-coated tablets

(prucalopride)

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

The name of your medicine is Resolor 2 mg film-coated tablets but will be referred to as Resolor throughout this leaflet.

Please note that the leaflet also contains information about other strengths.

What is in this leaflet

1. What Resolor is and what it is used for
2. What you need to know before you take Resolor
3. How to take Resolor
4. Possible side effects
5. How to store Resolor
6. Contents of the pack and other information

1. What Resolor is and what it is used for

Resolor contains the active substance prucalopride.

Resolor belongs to a group of gut motility enhancing medicines (gastrointestinal prokinetics). It acts on the muscle wall of the gut, helping to restore the normal functioning of the bowel. Resolor is used for the treatment of chronic constipation in adults in whom laxatives do not work well enough.

Not for use in children and adolescents younger than 18 years.

2. What you need to know before you take Resolor

Do not take Resolor:

- if you are allergic to prucalopride or any of the other ingredients of this medicine (listed in section 6).
- if you are on renal dialysis,
- if you suffer from perforation or obstruction of the gut wall, severe inflammation of the intestinal tract, such as Crohn's disease, ulcerative colitis or toxic megacolon/megarectum.

Warnings and precautions

Talk to your doctor before taking Resolor.

Take special care with Resolor and tell your doctor if you:

- suffer from severe kidney disease,
- suffer from severe liver disease,
- are currently under supervision by a doctor for a serious medical problem such as lung or heart disease, nervous system or mental health problems, cancer, AIDS or a hormonal disorder.

If you have very bad diarrhoea, the contraceptive pill may not work properly and the use of an extra method of contraception is recommended. See the instructions in the patient leaflet of the contraceptive pill you are taking.

Other medicines and Resolor

Tell your doctor if you are taking, or have recently taken, or might take any other medicines.

Resolor with food and drink

Resolor can be taken with or without food and drinks, at any time of the day.

Pregnancy and breast-feeding

Resolor is not recommended for use during pregnancy.

- Tell your doctor if you are pregnant or planning to become pregnant.
- Use a reliable method of contraception while you're taking Resolor, to prevent pregnancy.
- If you do become pregnant during treatment with Resolor, tell your doctor.

When breast-feeding, prucalopride can pass into breast milk. Breast-feeding is not recommended during treatment with Resolor. Talk to your doctor about this.

Ask your doctor for advice before taking any medicine.

Driving and using machines

Resolor is unlikely to affect your ability to drive or use machines. However, sometimes Resolor may cause dizziness and tiredness, especially on the first day of treatment, and this may have an effect on driving and use of machines.

Resolor contains lactose

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

3. How to take Resolor

Always take this medicine exactly as described in this leaflet or as your doctor has told you. Check with your doctor or pharmacist if you are not sure. Take Resolor every day for as long as your doctor prescribes it.

The doctor may want to reassess your condition and the benefit of continued treatment after the first 4 weeks and thereafter at regular intervals.

The usual dose of Resolor for most patients is one 2 mg tablet once a day.

If you are older than 65 years or have severe liver disease, the starting dose is one 1 mg tablet once a day, which your doctor may increase to 2 mg once a day if needed.

Your doctor may also recommend a lower dose of one 1 mg tablet daily if you have severe kidney disease.

Taking a higher dose than recommended will not make the product work better.

Resolor is only for adults and should not be taken by children and adolescents up to 18 years.

If you take more Resolor than you should

It is important to keep to the dose as prescribed by your doctor. If you have taken more Resolor than you should, it is possible that you will get diarrhoea, headache and/or nausea. In case of diarrhoea, make sure that you drink enough water.

If you forget to take Resolor

Do not take a double dose to make up for a forgotten tablet. Just take your next dose at the usual time.

If you stop taking Resolor

If you stop taking Resolor, your constipation symptoms may come back again.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them. The side effects mostly occur at the start of treatment and usually disappear within a few days with continued treatment.

The following side effects have been reported very commonly (may affect more than 1 in 10 people): headache, feeling sick, diarrhoea and abdominal pain.

The following side effects have been reported commonly (may affect up to 1 in 10 people): decreased appetite, dizziness, vomiting, disturbed digestion (dyspepsia), windiness, abnormal bowel sounds, tiredness.

The following uncommon side effects have also been seen (may affect up to 1 in 100 people): tremors, pounding heart, rectal bleeding, increase in frequency of passing urine (pollakiuria), fever and feeling unwell. If pounding heart occurs, please tell your doctor.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the **Google Play** or **Apple App Store**. By reporting side effects, you can help provide more information on the safety of this medicine.

5. How to store Resolor

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the blister and carton after EXP. The expiry date refers to the last day of that month.

Store in the original blister package in order to protect from moisture.

If this medicine becomes discoloured or shows any other signs of deterioration, please contact your pharmacist who will advise you on what to do.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information**What Resolor contains**

The active substance is prucalopride.

One film-coated tablet of Resolor 2 mg contains 2 mg prucalopride (as succinate).

The other ingredients are:

Lactose monohydrate (see section 2), microcrystalline cellulose, colloidal silicon dioxide, magnesium stearate, hypromellose, triacetin, titanium dioxide (E171), macrogol. The 2 mg tablet also contains iron oxide red (E172), iron oxide yellow (E172), indigo carmine aluminium lake (E132).

What Resolor looks like and contents of the pack

Resolor 2 mg film-coated tablets are pink round shaped tablets marked "PRU 2" on one side.

Resolor is provided in aluminium/aluminium perforated unit dose blister (calendar marked) containing 7 tablets. Each pack contains 28x1 film-coated tablets.

Product Licence Holder and Manufacturer

Procured from within the EU by the Product Licence holder Orifarm UK Ltd., Ground Floor Suite A2, Breakspear Park, Breakspear Way, Hemel Hempstead, HP2 4TZ, UK

For any information about this medicine, please contact the Product Licence Holder on www.orifarm.com/uk

Or phone: (+44) 1923 204333

Repacked by Orifarm Supply s.r.o., Palouky 1366, 253 01 Hostivice, Czech Republic

Manufactured by Sanico NV, Veedijk 59, B-2300 Turnhout, Belgium

Resolor 2 mg film-coated tablets
PL 45985/0973

POM

Leaflet revision date: 17/10/2025

**Blind or partially sighted?
Is this leaflet hard to see or read?
Call +45 63 95 27 00
to obtain the leaflet in a format suitable for you.**