
170 mm		170 mm		150 mm		
		490 mm		•		
PACKAGE LEAFLET:	• adrenocortical disorder (e.g. Addison's disease)	Addiction can cause withdrawal symptoms when	muscles, including the muscles, that control	If you use Subutex during pregnancy, your		
INFORMATION FOR THE USER	 depression or other conditions that are treated 	you stop taking this medicine. Withdrawal	movement of the eye, agitation, hallucinations,	baby may become dependent and experience	ω	
Subutex [®] 0.4 mg, 2 mg and 8 mg	with antidepressants. The use of these medicines together with Subutex can lead to servitionin	symptoms can include restlessness, difficulty sleeping, irritability, agitation, anxiety, feeling	coma, excessive sweating, tremor, exaggeration of reflexes, increased muscle tension, body	withdrawal symptoms including problems with breathing after the birth which may need to be	7.5	
sublingual tablets	syndrome, a potentially life-threatening condition	your heartbeat (palpitations), increased blood	temperature above 38°C. Contact your doctor	treated. These symptoms may occur several days	H H	
0	(see "Other medicines and Subutex").	pressure, feeling or being sick, diarrhoea, loss of	when experiencing such symptoms.	after birth.	В	
buprenorphine	Important things to be aware of:	appetite, shaking, shivering or sweating. Your	 Other medicines that may make you feel sleepy 	Do not take Subutex while you are breastfeeding as		
lead all of this leaflet carefully before you start	 Misuse, abuse and diversion 	prescriber will discuss with you how to gradually	which are used to treat illnesses such as anxiety,	buprenorphine passes into breast milk and will		
aking this medicine because it contains important + -	This medicine can be a target for people who	- + -reduce your dose before stopping the medicine. It+	– – – sleeplessness, convulsions/seizures, pain. These –	affect your baby.		
nformation for you.	abuse prescription medicines, and should be kept	is important that you do not stop taking the	types of medicines will reduce your alertness	Driving and using machines		
Keep this leaflet. You may need to read it again. If you have any further questions, ask your doctor	in a safe place to protect it from theft. Opioids	medicine suddenly as you will be more likely to	levels making it difficult for you to drive and use	If you feel drowsy or dizzy while taking these tablets		
or pharmacist.	should only be used by those they are prescribed	experience withdrawal symptoms. This product can cause withdrawal symptoms if	machines. They may also cause central nervous system depression, which is very serious. Below	do not use machinery.	37	
This medicine has been prescribed for you only.	for. Do not give your medicine to anyone else.	you take it less than 6 hours after you use a	is a list of examples of these types of medicines:	The medicine can affect your ability to drive as it	UT I I I I I I I I I I I I I I I I I I I	
Do not pass it on to others. It may harm them,	Taking higher doses or more frequent doses of	short-acting opioid (e.g. morphine, heroin) or less	 other opioid containing medicines such as 	may make you sleepy or dizzy.		
even if their signs of illness are the same as	opioid may increase the risk of addiction. Overuse and misuse can lead to overdose and/or death.	than 24 hours after you use a long-acting opioid	methadone, certain pain killers and cough	 Do not drive while taking this medicine until you 		
yours.	 Breathing problems 	such as methadone.	suppressants.	know how it affects you.It is an offence to drive if this medicine affects		
If you get any side effects, talk to your doctor or	Some people have died	Liver damage	- antidepressants (used to treat depression) such	• It is an offence to drive if this medicine affects $-\frac{1}{2}$ -your ability to drive. $ -$		
pharmacist. This includes any possible side	from respiratory failure	Liver damage has been reported after taking	as isocarboxazide, phenelzine, selegeline,	 However, you would not be committing an 	A	
effects not listed in this leaflet. See section 4.	(inability to breathe)	Subutex, especially when the medicine is	tranylcypromine, and valproate may increase	offence if:		
Vhat is in this leaflet:	because they misused this	misused. This could also be due to viral infections (chronic hepatitis C), alcohol abuse, anorexia or	 the effects of this medicine. sedative H, receptor antagonists (used to treat 	o The medicine has been prescribed to treat a		
. What Subutex is and what it is used for . What you need to know before you take Subutex	medicine or took it in combination with other	use of other medicines with the ability to harm	- sedative H_1 receptor antagonists (used to treat allergic reactions) such as diphenhydramine	medical or dental problem and	37	
. How to take Subutex	combination with other	your liver (see section 4). Regular blood tests may	and chlorphenamine.	o You have taken it according to the instructions		
. Possible side effects	depressants, such as	be conducted by your doctor to monitor the	 barbiturates (used to cause sleep or sedation) 	given by the prescriber or in the information	-	
. How to store Subutex	alcohol, benzodiazepines	condition of your liver. Tell your doctor if you	such as phenobarbital, secobarbital.	provided with the medicine and		
. Contents of the pack and other information	(tranguilisers), or other	have any liver problems before you start	 tranquilisers (used to cause sleep or sedation) 	o It was not affecting your ability to drive safely Talk to your doctor or pharmacist if you are not sure		
his medicine contains buprenorphine which is an \perp $_$		treatment with Subutex	such as chloral_hydrate	whether it is safe for you to drive while taking this		
pioid, which can cause addiction. You can get	Tolerance, dependence and addiction	 <u>Blood pressure</u> 	 Naltrexone may prevent Subutex from working. If 	medicine.	Ţ	
vithdrawal symptoms if you stop taking it	This medicine contains buprenorphine which is an	This product may cause your blood pressure to	you take naltrexone whilst you are taking Subutex	Subutex contains lactose and sodium		
uddenly.	opioid medicine. Repeated use of opioids can result	drop suddenly, causing you to feel dizzy if you get up too quickly from sitting or lying down.	you may experience a sudden onset of prolonged and intense withdrawal symptoms.	This medicine contains lactose. If you have been	ω	
WHAT SUBUTEX IS AND WHAT IT IS USED FOR	in the drug being less effective (you become	 Diagnosis of unrelated medical conditions 	 Clonidine (used to treat high blood pressure) may 	told by your doctor that you have an intolerance to	7.5	
ubutex is used to treat dependence on opiate	accustomed to it, known as tolerance). Repeated use	This medicine may mask pain symptoms that	extend the effects of this medicine.	some sugars, contact your doctor before taking this	m	
narcotic) drugs, such as morphine and heroin in	of buprenorphine can also lead to dependence,	could assist in the diagnosis of some diseases. Do	 Anti-retrovirals (used to treat AIDS) such as 	medicine.	3	
rug addicts who have agreed to be treated for their	abuse, and addiction, which may result in life-threatening overdose.	not forget to advise your doctor if you take this	ritonavir, nelfinavir, indinavir may increase the	This medicine contains less than 1 mmol sodium		
ddiction. ubutex is used in adults and adolescents over 16	Dependence or addiction can make you feel that you	medicine.	effects of this medicine.	(23 mg) per tablet, that is to say essentially 'sodium-free'.		
ears of age who are also receiving medical, social	are no longer in control of how much medicine you	Other medicines and Subutex	 - Some antifungal agents (used to treat fungal infections) such as ketoconazole and itraconazole 	1	E E E E E E E E E E E E E E E E E E E	
nd psychological support.	need to take or how often you need to take it.	Tell your doctor or pharmacist if you are taking,	and certain antibiotics (macrolide) may extend the	3. HOW TO TAKE SUBUTEX	E	
his medicine contains buprenorphine which	The risk of becoming dependent or addicted varies	have recently taken or might take any other medicines.	effects of this medicine.	You must place the tablet under your tongue		
elongs to a class of medicines called opioids.	from person to person. You may have a greater risk	Some medicines may increase the side effects of	 medicines used to treat allergies, travel sickness 	(sublingual) and allow it to dissolve, which will take 5 to 10 minutes. This is the only way to take the	37.	
his medicine has been prescribed to you and	of becoming dependent on or addicted to buprenorphine if:	Subutex and may sometimes cause very serious	or nausea (antihistamines or antiemetics);	tablets. Do not chew or swallow them whole, as	5 n	
nould not be given to anyone else. Opioids can	 You or anyone in your family have ever abused or 	reactions. Do not take any other medicines whilst	 medicines to treat psychiatric disorders 	they will not work.	m	
ause addiction and you may get withdrawal ymptoms if you stop taking it suddenly. Your	been dependent on alcohol, prescription	taking Subutex without first talking to your doctor,	(antipsychotics or neuroleptics);	Your doctor will tell you how many tablets to take		
rescriber should have explained how long you will	medicines or illegal drugs ("addiction").	especially:	 muscle relaxants; modicines to treat Parkingen's disease 	and you should always follow this advice.		
e taking it for and when it is appropriate to stop, - + -	– •– You are a smoker – – – – – – – – – – – – – – – – – – –	Benzodiazepines (used to treat anxiety or sleep	 medicines to treat Parkinson's disease. Some medicines may decrease the effect of 	- To avoid sudden withdrawal symptoms, treatment + -		
ow to do this safely.	 You have ever had problems with your mood 	disorders) such as diazepam, temazepam, alprazolam. Concomitant use of Subutex and	Subutex. These include medicines used to treat	with Subutex should be given when there are		
WHAT YOU NEED TO KNOW BEFORE YOU TAKE	(depression, anxiety, or a personality disorder) or	sedative medicines such as benzodiazepines or	epilepsy (such as carbamazepine and phenytoin)	already clear signs of withdrawal symptoms.		
SUBUTEX	have been treated by a psychiatrist for other mental illnesses.	related drugs increases the risk of drowsiness,	and medicines used to treat tuberculosis	Your prescriber should discuss your treatment and	ω,	
o not take Subutex:	If you notice any of the following signs whilst	difficulties in breathing (respiratory depression),	(rifampicin).	whether you need to continue taking tablets at	5	
If you are a child under the age of 16 years.	taking buprenorphine, it could be a sign that you	coma and may be life-threatening. Because of	To get the greatest benefit from taking Subutex, you	regular intervals. If you and your prescriber decide		
If you are allergic to buprenorphine or any of the	have become dependent or addicted:	this, concomitant use should only be considered	must tell your doctor about all the medicines you	to stop treatment, a plan will be put in place to gradually reduce the dose and stop taking the		
other ingredients of this medicine (listed in	 You need to take the medicine for longer than 	when other treatment options are not possible.	are taking, including alcohol, medicines containing alcohol, street drugs, and any prescription medicine	medicine to minimise the risk of		
section 6).	advised by your doctor	However if your doctor does prescribe Subutex	you are taking that has not been prescribed for you	withdrawal effects		
If you have serious breathing problems.	You need to take more than the recommended	- - together with sedative medicines the dose and $-$ - duration of concomitant treatment should be	by your doctor.	Adults and children over the age of 16 years: when		
If you have serious problems with your liver . If you are intoxicated due to alcohol or have	dose • You are using the medicine for reasons other than	limited by your doctor. Please tell your doctor	Subutex with food, drink and alcohol	beginning treatment the dose is between 0.8 to		
trembling, sweating, anxiety confusion or	 You are using the medicine for reasons other than prescribed, for instance, 'to stay calm' or 'help 	about all sedative medicines you are taking, and	Alcohol may increase drowsiness and may increase	4mg, taken once a day.		
hallucinations caused by alcohol.	vou sleep'	follow your doctor's dose recommendation	the risk of respiratory failure (inability to breathe) if	For drug addicts who have not had any withdrawal	37.5	
If you are breast feeding a baby.	 You have made repeated, unsuccessful attempts 	closely. It could be helpful to inform friends or	taken with Subutex. Do not take Subutex together	treatment: one dose of Subutex should be taken at		
/arnings and precautions	to quit or control the use of the medicine	relatives to be aware of the signs and symptoms	with alcohol. Do not swallow or consume food or	least 6 hours after the last use of the opioid (narcotic	3	
alk to your doctor before taking Subutex if you	 When you stop taking the medicine you feel 	stated above. Contact your doctor when experiencing such symptoms.	drink until the tablet is completely dissolved.	such as morphine or heroin), or when the first signs		
ave:	unwell, and you feel better once taking the	 Gabapentin or pregabalin to treat epilepsy or pain 	Pregnancy and breast-feeding If you are pregnant or breast-feeding, think you may	of craving appear. If you take it less than six hours		
seizures, fits or convulsions	medicine again ('withdrawal effects')	due to nerve problems (neuropathic pain).	be pregnant or are planning to have a baby, ask	after you use a narcotic you may get withdrawal	↓ <u>↓</u>	
asthma or other breathing problems	If you notice any of these signs, speak to your doctor to discuss the best treatment pathway for	 Anti-depressants, such as moclobemide, 	your doctor or pharmacist for advice before taking	symptoms.		
any liver disease such as hepatitis low blood pressure	you, including when it is appropriate to stop and	tranylcypromine, citalopram, escitalopram,	this medicine.	For patients taking methadone: before beginning treatment, your doctor should reduce your dose of		
recently suffered head injury or brain disease	how to stop safely (See section 3, If you stop	fluoxetine, fluvoxamine, paroxetine, sertraline,	Do not take Subutex if you are pregnant or think you	methadone to not more than 30mg a day. Subutex	37	
a urinary disorder (especially linked to enlarged	using Subutex).	duloxetine, venlafaxine, amitriptyline, doxepine,	might be pregnant unless you have discussed this	may cause withdrawal symptoms in patients who	.5	
prostate in men)	 Addiction and withdrawal symptoms 	or trimipramine. These medicines may interact	with your prescriber and the benefits of treatment	are dependent on methadone if used within		
any kidney disease	Taking this medicine regularly, particularly for a long time, can lead to addiction.	with Subutex and you may experience symptoms such as involuntary, rhythmic contractions of	are considered to outweigh the potential harm to	24 hours of the last dose of methadone.		
thyroid problems		SUCH AS INVOLUTIARY TRANSFORMER FOR A SUCH AS INVOLUTIARY TRANSFORMER FOR	the baby.			


CUSTOMER INFO: Minimum Point Size = 10.00pt



22 mm

			1		
	During your treatment, your doctor may increase your dose of Subutex, to a maximum single daily dose of 32mg, depending upon your response. Once you have been stable for a while, your doctor will	 very common (affects more than 1 user in 10) common (affects 1 to 10 users in 100) not known (frequency cannot be estimated from the available data). 	away medicines you no longer use. These measures will help to protect the environment. 6. CONTENTS OF THE PACK AND OTHER		
	gradually reduce your dose and it may be possible	Drug Withdrawal	INFORMATION		
	to stop it altogether. Do not suddenly stop taking the tablets, as this may cause withdrawal symptoms.	Side effects reported with Subutex	What Subutex contains The active substance is buprenorphine		
	If you take more Subutex than you should	Very common side effects:	(as buprenorphine hydrochloride).		
	If you or someone else takes too much of this	Drug withdrawal syndrome, headache,	Each sublingual tablet contains 0.4mg, 2mg or 8mg + -		
	medicine, you must go or be taken immediately to	hyperhidrosis (sweating), insomnia (inability to sleep), nausea (feeling sick), pain	of buprenorphine.		
	an emergency centre or hospital as overdose with Subutex may cause serious and life-threatening	Common side effects:	The other ingredients are monohydrated lactose, mannitol, maize starch, povidone K30, citric acid,		
	breathing problems.	Abdominal pain, agitation, anxiety, joint pain,	magnesium stearate and sodium citrate.		
	If you forget to take Subutex	weakness, back pain, bone pain, bronchitis, chest	What Subutex looks like and contents of the pack		
	Tell your doctor as soon as possible if you miss a dose and follow his or her instructions. Do not take	pain, chills, constipation, cough, decreased appetite, depression, diarrhoea, dizziness, dry	Subutex 0.4mg sublingual tablets are uncoated oval		
	a double dose to make up for the forgotten dose.	mouth, painful period, indigestion, shortness of	white tablets of 8 mm x 4 mm, debossed		
	If you stop taking Subutex	 breath, flatulence, gastrointestinal disorder, – – – hostility, increase in muscle tension, infection, 	- with-"04"-on one side		++-
	Do not change the treatment in any way or stop treatment without the agreement of the doctor who	influenza, nervousness, tearing (watery eyes)	Subutex 2mg sublingual tablets are uncoated oval		
	is treating you. Do not suddenly stop taking this	disorder, swollen glands (lymph nodes), malaise,	white tablets of 10 mm x 5 mm, debossed with "B2" on one side.	Arca	
	medicine. If you want to stop taking this medicine,	migraine, muscle spasms, muscle pain, dilation of the pupil, neck pain, palpitations, paranoia,	Subutex 8mg sublingual tablets are uncoated oval	Lice)	
	discuss this with your prescriber first. They will tell you how to do this, usually by reducing the dose	burning or tingling in hands and feet, swelling	white tablets of 14 mm x 7 mm,	ext 1	
	gradually so that any unpleasant withdrawal effects	(hands and feet), runny or stuffy nose, sore throat and painful swallowing, fever, rash, somnolence,	debossed with "B8" on one side.		
	are kept to a minimum. Withdrawal symptoms such as restlessness, difficulty sleeping, irritability,	syncope (fainting), thinking abnormal, tooth	The sublingual tablets come in blister packs		
	agitation, anxiety, feeling your heartbeat	disorder, tremor; flushing, vomiting (being sick), yawning	- containing either 7 or 28 tablets		
	(palpitations), increased blood pressure, feeling or being sick, diarrhoea, shaking, shivering or sweating	Frequency not known:	Marketing Authorisation Holder and Manufacturer		
	may occur if you suddenly stop taking this medicine.	Dental caries, drug dependence and addiction,	Marketing Authorisation Holder: Indivior UK Limited		
	If you have any further questions on the use of this	seizures, drug withdrawal syndrome in newborn,	The Chapleo Building		
	product, ask your doctor or pharmacist. How to remove the tablet from the blister pack	hallucinations (sensing things that are not real), drop in blood pressure on changing position	Henry Boot Way		
		from sitting or lying down to standing, difficulty	Priory Park, Hull		
		in urinating, vertigo Misusing this medicine by injecting it can cause	HU4 7DY		
		withdrawal symptoms, infections, other skin	Manufacturer		
		reactions and potentially serious liver problems	Reckitt Benckiser Healthcare (UK) Limited		
	1 – Remove just one section from the	When you stop taking Subutex, you may experience	Dansom Lane		
	blister pack, tearing it along the	drug withdrawal symptoms, which include restlessness, difficulty sleeping, irritability, agitation,	Hull, HU8 7DS		1 I I I I I I I I I I I I I I I I I I I
	perforated line.	anxiety, feeling your heartbeat (palpitations),			
		increased blood pressure, feeling or being sick, diarrhoea, shaking, shivering or sweating.	For any information about this medicine, please contact the Marketing Authorisation Holder:		
	2 – Starting from the edge where the	If any of the side effects gets serious, or if you notice	Indivior UK Limited		
	seal is lifted, pull back the foil on the back to remove the tablet.	any side effects not listed in this leaflet, please tell your doctor or pharmacist.	Telephone 0808 234 9243		
		Reporting of side effects	PatientSafetyRoW@indivior.com		
	4. POSSIBLE SIDE EFFECTS	If you get any side effects, talk to your doctor,	This leaflet was last revised in		
	Like all medicines, this medicine can cause side effects, although not everybody gets them.	pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report			
	Tell your doctor immediately or seek urgent	side effects directly via the Yellow Card Scheme	I I I I I I I I I I I I I I I I I I I		
	medical attention if you experience side effects	website: www.mhra.gov.uk/yellowcard or search for			
	 such as: sudden wheezing, difficulty breathing, swelling of 	 MHRA Yellow Card in the Google Play or Apple App Store. 	i i i i i i i i i i i i i i i i i i i		
	the eyelids, face, tongue, lips, throat or hands;	By reporting side effects you can help provide more			
	rash or itching especially those covering your whole body. These may be signs of a	information on the safety of this medicine.			
	life-threatening allergic reaction.	5. HOW TO STORE SUBUTEX			
	 if you start to breathe more slowly or weakly than expected (respiratory depression). 	Keep out of the sight and reach of children. Store this medicine in a safe and secure place,			
- I					

expected (respiratory depression).



-6

-8 -6

-2


-22-

-5

-8

20-

-6



- I

1

where other people cannot access it. It can cause