

the lowest dose for the shortest time possible should be used. If taken for more than a few days from 20 weeks of pregnancy onward, Combogesic IV can cause kidney problems in your unborn baby that may lead to low levels of amniotic fluid that surrounds the baby (oligohydramnios) or narrowing of a blood vessel (ductus arteriosus) in the heart of the baby. If you need treatment for longer than a few days, your doctor may recommend additional monitoring.

Breast-feeding

Only small amounts of paracetamol and ibuprofen pass into breast milk. This medicine may be given during breast-feeding, if it is used at the recommended dose and for the shortest possible time.

Fertility

This product may impair female fertility and is not recommended in women attempting to conceive. This effect is reversible on stopping the medicine.

Driving and using machines

Undesirable effects such as dizziness, drowsiness, fatigue and visual disturbances are possible after taking NSAIDs. If affected you should not drive or operate machinery.

Combogesic IV contains sodium

Combogesic IV contains 35 mg sodium (main component of cooking/table salt) in each 100 ml. This is equivalent to 1.75% of the recommended maximum daily dietary intake of sodium for an adult.

3. How to use Combogesic IV

Combogesic IV will be given to you by a healthcare professional by infusion into one of your veins. The infusion should be administered over 15 minutes.

This medicine is for short term use only, maximum 2 days.

The recommended dose is:

For adults who weigh more than 50 kg: 1 vial every 6 hours, as required.

The maximum daily dose is four vials which equals 4000 mg (4 g) paracetamol and 1200 mg ibuprofen.

If you weigh 50 kg or less, are elderly or if you have liver or kidney problems: Your doctor may decide to reduce your dose or increase the time between doses because of the increased risk of side effects.

Do not take more medicine than the label tells you to. If you do not get better, talk to your doctor.

A higher dose than the recommended does not increase pain relief; instead it can lead to serious risks (see also section “**If you are given more Combogesic IV than you should**”). The lowest effective dose should be given for the shortest duration necessary to relieve symptoms. If you have an infection, consult a doctor without delay if symptoms (such as fever and pain) persist or worsen (see section 2).

If you are given more Combogesic IV than you should

Immediately contact a doctor or nurse if you think that you accidentally may have been given too much of this medicine. **Do this even if you feel well.** This is because too much paracetamol can cause delayed, serious liver damage, which may be fatal. Even if there are no signs of discomfort or poisoning, you may need urgent medical attention.

In order to avoid liver damage it is essential to get medical treatment as early as possible. The shorter the interval between intake and initiation of treatment with antidote (as few hours as possible), the greater the likelihood that hepatic injury can be prevented.

Further symptoms can include nausea, stomach pain, vomiting (may be blood streaked), headache, ringing in the ears, confusion and shaky eye movement. At high doses, drowsiness, chest pain, palpitations, loss of consciousness, convulsions (mainly in children), weakness and dizziness, blood in urine, cold body feeling, and breathing problems have been reported.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Stop using Combogesic IV and tell your doctor **immediately** or go to the emergency room at your nearest hospital if you get any of the following side effects:

Uncommon:

- vomiting blood or material that looks like coffee grounds;
- bleeding from the back passage, black sticky bowel motions (stools) or bloody diarrhoea;
- swelling of the face, lips or tongue which may cause difficulty in swallowing or breathing.

Very rare:

- asthma, wheezing, shortness of breath;
- sudden or severe itching, skin rash, hives;
- severe rash with blisters and bleeding in the lips, eyes, mouth, nose and genitals (Steven Johnson Syndrome). Very rare cases of serious skin reactions have been reported;
- worsening of existing severe skin infections (you may notice a rash, blistering and discolouration of the skin, fever, drowsiness, diarrhoea and sickness), or worsening of other infections including chicken pox or shingles or severe infection with destruction (necrosis) of subcutaneous tissue and muscle, blistering and peeling of the skin;
- fever, generally feeling unwell, nausea, stomach ache, headache and stiff neck (symptoms of aseptic meningitis, inflammation of the protective membrane surrounding the brain).

Not known (frequency cannot be estimated from the available data):

- a severe skin reaction known as DRESS syndrome can occur. Symptoms of DRESS include: skin rash, fever, swelling of lymph nodes and an increase of eosinophils (a type of white blood cells).
- a red, scaly widespread rash with bumps under the skin and blisters mainly localised on the skin folds, trunk, and upper extremities accompanied by fever at the initiation of treatment (acute generalised exanthematous pustulosis). See also section 2.
- A serious condition that can make blood more acidic (called metabolic acidosis), in patients with severe illness using paracetamol (see section 2).
- Combogesic IV, especially when taken at higher than recommended doses or for a prolonged period of time, can cause damage to your kidneys and affect them removing acids properly from your blood into the urine (renal tubular acidosis). It can also cause very low levels of potassium in your blood (see section 2). This is a very serious condition and will require immediate treatment. Signs and symptoms include muscle weakness and light-headedness.

Other side effects that may occur:

Common (may affect up to 1 in 10 people):

- nausea or vomiting;
- loss of appetite;
- heartburn or pain the upper part of your stomach;
- stomach cramps, wind, constipation or diarrhoea, slight gastrointestinal blood loss;
- skin rashes, itching of the skin;
- headache;
- dizziness;
- feeling of being nervous;
- ringing or buzzing in the ears;
- unusual weight gain, swelling and fluid retention, swelling of ankles or legs (oedema).

Uncommon (may affect up to 1 in 100 people):

- decrease in red blood cells, nose bleed and heavier periods (menstrual bleeding);
- allergic reactions – skin rash, tiredness, joint pain (e.g. serum sickness, lupus erythematosus syndrome, Henoch-Schönlein vasculitis, angioedema);
- enlargement of breast tissue in men; low blood sugar levels;
- sleeplessness;
- change in mood, for example depression, confusion, nervousness;
- eye problems such as blurred vision (reversible), sore red eyes, itching;
- thickened mucus;
- severe pain or tenderness in the stomach; peptic/gastrointestinal ulcer;
- Bowel inflammation and worsening of inflammation of the colon (colitis) and digestive tract (Crohn's disease) and complications of diverticula of the large bowel (perforation or fistula);
- inability to completely empty the bladder (urinary retention);
- abnormal laboratory test results (blood, liver and kidney enzyme test results).

Rare (may affect up to 1 in 1,000 people):

- tingling of the hands and feet;
- abnormal dreams, seeing things (hallucinations);
- damage of the kidney tissue (particularly in long-term use);
- high level of uric acid in your blood (hyperuricemia).

Very rare (may affect up to 1 in 10,000 people):

- low potassium levels – weakness, fatigue, muscle cramps (hypokalaemia);
- signs of anaemia, such as tiredness, headaches, being short of breath, and looking pale;
- bleeding or bruising more easily than normal, reddish or purplish blotches under the skin;
- severe or persistent headache;
- spinning sensation (vertigo);
- fast or irregular heartbeats, also called palpitations;
- increase in blood pressure and possible heart problems;
- inflammation of the oesophagus;
- yellowing of the skin and /or eyes, also called jaundice;
- liver damage (particularly in long term use);
- loss of hair;
- increase in sweating;
- signs of frequent or worrying infections such as fever, severe chills, sore throat or mouth ulcers;
- nephrotoxicity in various forms, including interstitial nephritis, nephrotic syndrome, and acute and chronic renal failure.

Reporting of side effects

If you get any side effects talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme Website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provided more information on the safety of this medicine.

5. How to store Combogesic IV

Keep this medicine out of the sight and reach of children and adolescents.

Store below 25°C. Do not refrigerate or freeze. Store in the original carton in order to protect from light.

Do not use this medicine after the expiry date which is stated on the carton and on the vial after EXP. The expiry date refers to the last day of that month.

Do not use this medicine if you notice the packaging is torn or shows signs of tampering. Do not use this medicine if you notice any visible particles or discolouration.

This product is for single use only. The product should be used immediately after opening. Any unused solution should be discarded.

Dispose in accordance with local requirements.

6. Contents of the pack and other information

What Combogesic IV contains

The active substances are 10 mg/ml paracetamol and 3 mg/ml ibuprofen (as sodium dihydrate).

The other ingredients are cysteine hydrochloride monohydrate, disodium phosphate dihydrate, mannitol, hydrochloric acid (for pH adjustment), sodium hydroxide (for pH adjustment), and water for injections.

What Combogesic IV looks like and contents of the pack

Combogesic IV is a clear, colourless solution for infusion, free from visible particles. It is supplied in 100 ml clear glass vials, closed with a grey bromobutyl rubber stopper and an aluminium flip-off cap. It comes in a pack size of 10 vials.

Marketing Authorisation Holder and Manufacturer

Marketing Authorisation Holder:	Manufacturer:
AFT Pharma UK Limited,	S.M. Farmaceutici SRL
Olympia House,	Zona Industriale
Unit 13, 2nd Floor, Armitage Road,	85050 Tito (PZ), Italy
London, NW11 8RQ,	
England	

This leaflet was last revised in June 2025

PL 57592/0001



Method of administration

Combogesic IV should be administered as a 15-minute intravenous infusion.

To remove solution, use a 0.8 mm needle (21 gauge needle) and vertically perforate the stopper at the spot specifically indicated.

In patients weighing less than 50 kg for whom a full vial (100 ml) is not required, the correct amount should be infused and the remaining solution discarded.

As for all solutions for infusion presented in glass vials, it should be remembered that close monitoring is needed notably at the end of the infusion, regardless of administration route. This monitoring at the end of the perfusion applies particularly for central route infusion, in order to avoid air embolism.