

Pharma Code

5mm

15mm

300mm

Package leaflet: Information for the user P2192T

Levothyroxine 12.5 micrograms, 25 micrograms, 50 micrograms, 75 micrograms and 100 micrograms Tablets

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

- What Levothyroxine Tablets are and what they are used for
- What you need to know before you take Levothyroxine Tablets
- How to take Levothyroxine Tablets
- Possible side effects
- How to store Levothyroxine Tablets
- Contents of the pack and other information

1. What Levothyroxine Tablets are and what they are used for

Thyroxine is a hormone which is produced naturally in the body by the thyroid gland. Levothyroxine is a synthetic version of this hormone. Thyroxine controls how much energy your body uses. When the thyroid gland does not produce enough thyroxine (a condition known as hypothyroidism), many of the body's functions slow down. Some of the most common symptoms of hypothyroidism are:

- tiredness
- weight gain
- feeling depressed

Levothyroxine Tablets are used to replace the thyroxine that your thyroid gland cannot produce and prevent the symptoms of hypothyroidism. Before starting your treatment your doctor will carry out a blood test to work out how much levothyroxine you need.

2. What you need to know before you take Levothyroxine Tablets

Do not take Levothyroxine Tablets if you:

- Are **allergic** to levothyroxine or any of the other ingredients of this medicine (listed in section 6)
- Suffer from an overactive thyroid gland that produces too much thyroid hormone (thyrotoxicosis)
- Have any condition that affects your adrenal glands (your doctor will be able to advise you if you are not sure).

If any of these apply to you, do not take this medicine and go back to your doctor to discuss your treatment.

Warnings and precautions

Talk to your doctor or pharmacist before taking Levothyroxine Tablets if you:

- have suffered with an **under active thyroid gland** for a long time
- suffer from **heart problems** including **angina, coronary artery disease** or **high blood pressure**
- are being treated for **diabetes**. The dose of your antidiabetic medicine may need to be changed as levothyroxine can raise blood sugar levels
- are over **50 years** of age.

Blood pressure will be regularly monitored when levothyroxine treatment is started in very low birth weight preterm neonates because rapid fall in blood pressure (known as circulatory collapse) may occur.

Blood tests

Before you start taking levothyroxine your doctor will do a blood test to see how much thyroxine your thyroid gland is making and what dose of the medicine you will need. Once you start taking the medicine your doctor will want you to have regular blood tests to see how well the medicine is working.

Other medicines and Levothyroxine Tablets:

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. This includes over the counter medicines, herbal remedies and vitamin supplements.

Many medicines affect the way levothyroxine works. The effects of other drugs may also be affected by levothyroxine.

The following medicines may affect the way that levothyroxine works:

- medicines for epilepsy such as **carbamazepine, phenytoin, primidone and barbiturates**
- sertraline** – used to treat depression and anxiety disorders
- antacids** – used to treat indigestion
- medicines **containing calcium salts**
- cimetidine** – used to reduce excess stomach acid
- proton pump inhibitors** such as omeprazole, lansoprazole and pantoprazole - used to reduce the amount of acid produced by the stomach
- sucralfate** – used to treat and prevent stomach and duodenal ulcers
- cholestyramine and colestipol** – used to treat high level of fat in the blood
- polystyrene sulphone resin** – used to reduce high levels of potassium in the blood
- medicines containing iron that are taken by mouth
- rifampicin** – used to treat infections
- imatinib** – used to treat certain types of cancer
- beta blockers** such as **atenolol** and **sotalol** – used to treat high blood pressure and heart problems
- oestrogen containing medicines** for hormone replacement therapy (HRT) and contraception (the 'pill')
- androgen containing medicines** for male hormone replacement therapy
- corticosteroids** such as **hydrocortisone** and **prednisolone** – used to treat inflammation
- amiodarone** – used to treat an irregular heart beat
- orlistat** – used to treat obesity
- ritonavir** – used to control HIV and chronic hepatitis C virus.

The following medicines may be affected by levothyroxine:

- anticoagulant medicines** to prevent blood clots such as warfarin
- medicines to treat diabetes such as **insulin** and **metformin**
- tricyclic antidepressants** such as **amitriptyline, imipramine** and **dosulepin**
- medicines that stimulate the sympathetic nervous system such as **adrenaline** (used to treat severe allergic reactions) or **phenylephrine** (a decongestant found in many cold and flu treatments)
- digoxin** – used to treat heart problems
- anti-inflammatory medicines such as **phenylbutazone** or **aspirin**
- propranolol** – used to treat high blood pressure and heart problems
- ketamine** – used as an anaesthetic. If you need to have an operation, please tell your doctor or anaesthetist that you are taking levothyroxine.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine. Your doctor will decide if you should continue treatment with levothyroxine whilst you are pregnant, particularly in the first three months of your pregnancy.

Driving and using machines

This medicine should not affect your ability to drive and use machines.

Levothyroxine Tablets contain lactose monohydrate

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

Sodium content

This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium-free'.

3. How to take Levothyroxine Tablets

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

You may be taking this medicine for the rest of your life. Your dose will be decided by your doctor and will depend on the results of your blood tests.

The dose you should take will be on the label attached by your pharmacist.

How and when to take the tablets

Swallow the tablets with plenty of water. You should usually take your tablets before breakfast or your first meal of the day.

Adults:

The recommended starting dose is 50 – 100 micrograms every day.

5mm

92.5mm

5mm

92.5mm

5mm

200mm

5mm

300mm

Your doctor may increase the dose you take every 3 – 4 weeks by 50 micrograms until your thyroxine levels are correct. Your final daily dose may be up to 100 – 200 micrograms daily.

Patients over 50 years of age:

The recommended starting dose will be no more than 50 micrograms every day.

The dose may then be increased by 50 micrograms every 3 – 4 weeks until your thyroxine levels are correct. Your final daily dose will be between 50 – 200 micrograms daily.

Patients over 50 years of age with heart problems:

The recommended starting dose will be 25 micrograms every day or 50 micrograms every other day.

The dose may be increased by 25 micrograms every 4 weeks until your thyroxine levels are correct. Your final daily dose will usually be between 50 – 200 micrograms daily.

Use in children and adolescents:

For young children, your doctor is likely to prescribe Levothyroxine Oral Solution instead of tablets.

Congenital hypothyroidism in infants:

This is a condition where your baby has been born with a thyroid gland that does not produce enough thyroxine.

The starting dose is 10 -15 micrograms/kg bodyweight per day for the first three months. The dose will then be adjusted depending on how your baby responds to the treatment.

Acquired hypothyroidism in children:

This is a condition where your child's thyroid gland stops working properly because it has been attacked by their immune system, e.g. in children with an autoimmune disease or following a viral infection.

The starting dose is 12.5 – 50 micrograms per day. The dose will then be increased every 2 - 4 weeks depending on how your child responds to the medicine.

Juvenile myxoedema:

This is a condition where children and adolescents develop severe hypothyroidism (produce very low levels of thyroid hormones).

The starting dose is 25 micrograms every day. The dose will then be increased by 25 micrograms every 2 – 4 weeks until your child shows mild symptoms of hyperthyroidism (a condition where the thyroid gland produces too much thyroxine). The dose will then be reduced slightly.

The score line is not intended for breaking the tablet.

If you take more Levothyroxine Tablets than you should:

If you (or someone else) swallow a lot of the tablets at the same time, or you think a child may have swallowed some, contact your nearest casualty department or tell your doctor immediately.

Signs of an overdose may include: fever, chest pain (angina), racing or irregular heartbeat, muscle cramps, headache, restlessness, flushing, sweating and diarrhoea. These signs can take up to 5 days to appear.

Take any remaining tablets and this leaflet with you so that the medical staff know exactly what you have taken.

If you forget to take Levothyroxine Tablets:

If you forget to take a dose take it as soon as you remember unless it is nearly time for your next dose. Do not take a double dose to make up for a forgotten dose. If you forget to give your child their dose, contact your doctor or pharmacist for further advice.

If you stop taking Levothyroxine Tablets:

These tablets are for long term use. You may need to take them for the rest of your life.

Do not stop taking the tablets unless your doctor has told you to do so.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Stop taking Levothyroxine Tablets and see a doctor or go to a hospital right away if you have:

- a rare allergic reaction such as swelling of the face, tongue, lips and throat, difficulty breathing, severe itching of your skin with raised lumps, joint pain, sensitivity to the sun, general feeling of being unwell. You may need urgent medical attention.

Some patients may experience a severe reaction to high levels of thyroid hormone.

This is called a “thyroid crisis” and you should contact your doctor immediately if you have any of the following symptoms:

- very high temperature; fast heart rate; irregular heartbeat; low blood pressure; heart failure; jaundice; confusion; fits and coma.

Tell your doctor or pharmacist if any of the following side effects continue, get worse or if you notice any other side effects not listed:

Most of the side effects are similar to the symptoms of hyperthyroidism (where the thyroid gland makes too much thyroxine) and are due to your dose of the medicine being too high. They will usually disappear after reducing the dose or stopping the tablets. **However, you must not change the dose or stop the tablets without talking to your doctor first.**

Not known (frequency cannot be estimated from the available data)

- headache
- flushing
- high temperature, sweating
- weight loss
- tremor, restlessness, excitability, difficulty sleeping (insomnia)
- increased pressure around the brain in children that is not caused by a tumour or other diseases (benign intracranial hypertension)
- chest pain (angina), pounding, irregular or fast heartbeat
- diarrhoea, vomiting
- muscle cramps, muscle weakness
- deformity of the skull in infants caused by the early closure of joints in the skull bone (craniostenosis)
- growth in children may slow or stop due to changes in bone growth
- irregular periods
- intolerance to heat
- temporary hair loss in children.

Reporting of side effects:

If you get any side effects, talk to your doctor, pharmacist, or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Levothyroxine Tablets

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the label and carton after EXP. The expiry date refers to the last date of that month.

Do not store above 25°C.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Levothyroxine Tablets contain:

- The active substance is levothyroxine sodium. Each tablet contains either 12.5, 25, 50, 75 or 100 micrograms of levothyroxine sodium.
- The other ingredients are lactose monohydrate, maize starch, croscarmellose sodium, gelatin and magnesium stearate.

What Levothyroxine Tablets look like and contents of the pack:

Levothyroxine sodium 12.5 micrograms tablets are round and white to off-white coloured. It is plain on one side and debossed on the other side with 12.5.

Levothyroxine sodium 25 micrograms, 50 micrograms, 75 micrograms and 100 micrograms Tablets are round and white to off-white coloured. The tablets have a score line on one side and are debossed on the other side with either:

- 25 (25mcg)
- 50 (50mcg)
- 75 (75mcg)
- 100 (100mcg)

Levothyroxine Tablets are available in packs containing 28 tablets.

Marketing Authorisation Holder and Manufacturer:

Crescent Pharma Ltd.,
Key House, Sarum Hill,
Basingstoke,
RG21 8SR, UK.

This leaflet was last revised in March 2023.

P2192T

5mm

92.5mm

5mm



92.5mm

5mm

200mm



Product:	Levothyroxine tablets - ACME Generics
Item Code:	P2192T
Size:	200mm x 300mm
Min. Point size:	8.5pt
O. Artwork	06/03/23
Proof No.	3
Amendment Date	29/03/23

Colours Used	
	Process Black
	Keyline - does not print