

Dimetrum® 2 mg tablets

(dienogest)

Patient Information Leaflet

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See Section 4.

The name of your medicine is Dimetrum 2 mg tablets but will be referred to as Dimetrum throughout this leaflet.

What is in this leaflet:

- 1 What Dimetrum is and what it is used for
- 2 What you need to know before you take Dimetrum
- 3 How to take Dimetrum
- 4 Possible side effects
- 5 How to store Dimetrum
- 6 Contents of the pack and other information

1 What Dimetrum is and what it is used for

Dimetrum is a preparation for the treatment of endometriosis (painful symptoms due to displaced tissue of the lining of the womb). Dimetrum contains a hormone, the progestogen dienogest.

2 What you need to know before you take Dimetrum

Do not take Dimetrum if you:

- * are suffering from a blood clot (thromboembolic disorder) in your veins. This may occur, for example, in the blood vessels of the legs (deep vein thrombosis) or the lungs (pulmonary embolism). See also "*Dimetrum and venous blood clots*" below
- * have or have ever had a **severe arterial disease**, including cardiovascular disease, such as a **heart attack, stroke** or **heart disease** which causes a reduced blood supply (angina pectoris). See also "*Dimetrum and arterial blood clots*" below
- * have **diabetes** with blood vessel damage
- * have or have ever had **severe liver disease** (and your liver function values have not returned to normal). Symptoms of liver disease may be yellowing of the skin and/or itching of the whole body,
- * have or have ever had a **benign or malignant liver tumour**
- * suffer or have ever suffered, or if it is suspected that you suffer from a **malignant** sex-hormone dependent tumour such as cancer of the breast or the genital organs
- * have any unexplained **vaginal bleeding**
- * are **allergic (hypersensitive)** to dienogest or any of the other ingredients of this medicine (see *Section 6, and end of Section 2*).

If any of these conditions appear for the first time while using Dimetrum, stop taking it at once and consult your doctor.

Warnings and precautions

You must not use hormonal contraceptives of any form (tablet, patch, intrauterine system) while taking Dimetrum.

Dimetrum is NOT a contraceptive. If you want to prevent pregnancy, you should use condoms or other non-hormonal contraceptive precautions.

In some situations you need to take special care while using Dimetrum, and your doctor may need to examine you regularly. Tell your doctor if any of the following conditions applies to you:

If you:

- have ever had a **blood clot** (venous thromboembolism) or anyone in your immediate family has had a blood clot at a relatively early age
- have a close relative who has had **breast cancer**
- have ever suffered from **depression**
- have **high blood pressure** or develop high blood pressure while taking Dimetrum
- develop a **liver disease** while taking Dimetrum. Symptoms may include yellowing of the skin or eyes or itching all over your body. Inform your doctor also if such symptoms occurred during a previous pregnancy
- have diabetes or had **diabetes** temporarily during previous pregnancy
- have ever had **chloasma** (golden-brown patches on the skin, particularly of the face); if so, avoid too much exposure to the sun or ultraviolet radiation
- suffer from **pain in your lower abdomen** while taking Dimetrum.

While taking Dimetrum your chance of becoming pregnant is reduced because Dimetrum may affect ovulation.

If you become pregnant while taking Dimetrum you are at a **slightly increased risk** of having an extrauterine pregnancy (the embryo develops outside the womb). Tell your doctor before you start taking Dimetrum, if you had an extrauterine pregnancy in the past or have an impaired function of the Fallopian tubes.

Dimetrum and serious uterine bleeding

Uterine bleeding, for example in women with a condition where the mucous membrane of your uterus (endometrium) grows into the muscle layer of your uterus, called adenomyosis uteri or **benign tumours** of the womb sometimes called uterine fibroids (uterine leiomyomata), may become worse with the use of Dimetrum. If bleeding is heavy and continuous over time, this may lead to low red blood cell levels (anemia), which may be severe in some cases. In the event of anemia, you should discuss with your doctor if you should stop taking Dimetrum.

Dimetrum and changes in bleeding pattern

Most women treated with Dimetrum experience changes in their menstrual bleeding pattern (see section 4, possible side effects).

Dimetrum and venous blood clots

Some studies indicate that there may be a slight, but not statistically significant, increased risk of a **blood clot in the legs (venous thromboembolism)** associated with the use of preparations with progestagens like Dimetrum. Very rarely, blood clots may cause serious permanent disabilities or may even be fatal.

The risk of a **venous blood** clot increases:

- * with increasing age
- * if you are overweight
- * if you or one of your close relatives had a blood clot in the leg (thrombosis), lung (pulmonary embolism), or other organ at a young age
- * if you must have surgery, if you have had a serious accident or if you are immobilized for a long time. It is important to tell your doctor in advance that you are using Dimetrum as the treatment may have to be stopped. Your doctor will tell you when to start Dimetrum again. This is usually about two weeks after you are back on your feet.

Dimetrum and arterial blood clots

There is little evidence for an association between preparations with progestagens like Dimetrum and an increased risk of a blood clot in, for example, the bloodvessels of the heart (heart attack) or the brain (stroke). In women with hypertension the risk of stroke may be slightly enhanced by these preparations.

The risk of an **arterial blood** clot increases:

- * **if you smoke. You are strongly advised to stop smoking when you use Dimetrum, especially if you are older than 35 years.**
- * if you are overweight
- * if one of your close relatives had a heart attack or stroke at a young age
- * if you have high blood pressure

Talk to your doctor before taking Dimetrum

Stop taking Dimetrum and contact your doctor immediately if you notice possible signs of a blood clot, such as:

- * severe pain and/or swelling in one of your legs
- * sudden severe pain in the chest which may reach the left arm
- * sudden breathlessness
- * sudden cough without an obvious cause
- * any unusual, severe or long-lasting headache or worsening of migraine
- * partial or complete blindness or double vision
- * difficulty in speaking or inability to speak
- * giddiness or fainting
- * weakness, strange feeling, or numbness in any part of the body.

Dimetrum and cancer

It is not clear from the data currently available whether or not Dimetrum increases the risk of breast cancer. Breast cancer has been observed slightly more often in women taking hormones compared to those not taking hormones, but it is not known whether this is caused by the treatment. For example, it may be that more tumours are detected and detected earlier in women taking hormones because they are examined by their doctor more often. The occurrence of breast tumours becomes gradually less after stopping the hormone treatment. **It is important to regularly check your breasts** and you should contact your doctor if you feel any lump. In rare cases, benign liver tumours, and in even fewer cases malignant liver tumours have been reported in women taking hormones. Contact your doctor if you have unusually severe stomach pain.

Dimetrum and osteoporosis

Changes in bone mineral density (BMD)

The use of Dimetrum may affect the strength of the bone of adolescents (12 to under 18 years). If you are under 18 your doctor will, therefore, carefully weigh the benefits and risks of using Dimetrum for you as an individual patient, taking into account possible risk factors for bone loss (osteoporosis).

If you use Dimetrum, it will help your bones if you have an adequate intake of calcium and vitamin D either via your food or via supplements.

If you have an increased risk of getting osteoporosis (weakening of bones due to loss of bone minerals), your doctor will carefully weigh the risks and benefits of treatment with Dimetrum because Dimetrum has a moderate suppressing effect on the production of oestrogen (another type of female hormone) by your body.

Other medicines and Dimetrum

Always tell your doctor which medicines or herbal products you are already using. Also tell any other doctor or dentist who prescribes another medicine (or the pharmacist) that you are taking Dimetrum.

Some medicines can have an influence on the blood levels of Dimetrum and can make it less effective, or can cause undesirable effects.

These include:

- * medicines used for the treatment of:
 - * **epilepsy** (e.g. phenytoin, barbiturates, primidone, carbamazepine, oxcarbazepine, topiramate, felbamate)
 - * **tuberculosis** (e.g. rifampicin)
 - * **HIV and Hepatitis C Virus infections** (so-called protease inhibitors and non-nucleoside reverse transcriptase inhibitors such as ritonavir, nevirapine, efavirenz)
 - * **fungal infections** (griseofulvin, ketoconazole)
- * the herbal remedy **St. John's wort**.

Dimetrum® 2 mg tablets

Ref: 2581/010425/1/B

(dienogest)

Patient Information Leaflet (continued)

Ask your doctor or pharmacist for advice before taking any medicine.

Dimetrum with food and drink

During Dimetrum treatment, you should avoid drinking grapefruit juice, because this may increase the levels of Dimetrum in your blood. This may increase the risk of getting side effects.

Dimetrum can be taken with or without food.

Laboratory tests

If you need a blood test, tell your doctor or the laboratory staff that you are taking Dimetrum, because Dimetrum can affect the results of some tests.

Pregnancy, breast-feeding and fertility

Do not take Dimetrum if you are pregnant or breast-feeding.

Driving and using machines

No effects on the ability to drive and use machines have been observed in users of Dimetrum.

Dimetrum contains lactose

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking Dimetrum.

Children and adolescents

Dimetrum is not for use in girls before menarche (first menstrual bleeding). The use of Dimetrum may affect the strength of the bone of adolescents (12 to under 18 years). If you are under 18 your doctor will, therefore, carefully weigh the benefits and risks of using Dimetrum for you as an individual patient, taking into account possible risk factors for bone loss (osteoporosis).

3 How to take Dimetrum

Always take Dimetrum exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure. For adults, the usual dose is 1 tablet per day.

The following statements apply to Dimetrum unless otherwise prescribed by your doctor. Please follow these instructions, otherwise you will not fully benefit from Dimetrum.

You can start treatment with Dimetrum on any day of your natural cycle.

Adults: take one tablet every day, preferably at the same time with some liquid as needed. When a pack is finished the next one should be started without interruption. Continue to take the tablets also on days of menstrual bleeding.

If you take more Dimetrum than you should

There have been no reports of serious harmful effects from taking too many Dimetrum at one time. However, if you are concerned, contact your doctor.

If you forget to take Dimetrum or suffer from vomiting or diarrhoea

Dimetrum will be less effective if you miss a tablet. If you miss one or more tablets, take one tablet only as soon as you remember, and then continue next day taking the tablet at your usual time.

If you vomit within 3-4 hours of taking Dimetrum or you have severe diarrhoea, there is a risk that the active substance in the tablet will not be taken up by your body. The situation is almost the same as forgetting a tablet. After vomiting or diarrhoea within 3-4 hours of taking Dimetrum, you should take another tablet as soon as possible.

Do not take a double dose to make up for a forgotten tablet.

If you stop taking Dimetrum

If you stop taking Dimetrum, your original endometriosis symptoms may return.

4 Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them. These effects are more common during the first months after start of intake of Dimetrum and usually disappear with continued use. You may also experience changes in your bleeding pattern, such as spotting, irregular bleeding or your periods may stop completely.

Common (affecting between 1 and 10 in every 100 users)

- weight gain
- depressed mood, problems sleeping, nervousness, loss of interest in sex, or changed mood
- headache or migraine
- nausea, abdominal pain, wind, swollen tummy or vomiting
- acne or hair loss
- back pain
- breast discomfort, ovarian cyst or hot flushes
- uterine/vaginal bleeding including spotting
- weakness or irritability.

Uncommon (affecting between 1 and 10 in every 1,000 users)

- anemia
- weight loss or increase in appetite
- anxiety, depression or mood swings
- imbalance in the autonomic nervous system (controls unconscious bodily functions, e.g. perspiration) or disturbed attention
- dry eye
- tinnitus
- unspecific circulatory problems or uncommon palpitations
- low blood pressure

- shortness of breath
- diarrhoea, constipation, abdominal discomfort, inflammation the stomach and intestines, (gastrointestinal inflammation), inflammation of the gums (gingivitis)
- dry skin, excessive sweating, severe itching of the whole body, male pattern hair growth (hirsutism), brittle nails, dandruff, dermatitis, abnormal hair growth, hypersensitive response to light or problems with skin pigmentation
- pains in your bones, muscle spasms, pains and/or a sensation of heaviness in your arms and hands or legs and feet
- urinary tract infection
- vaginal thrush, dryness of the genital area, vaginal discharge, pelvic pain, atrophic inflammation of the genitals with discharge (atrophic vulvovaginitis), or a lump or lumps in the breast
- swelling due to fluid retention.

Additional side effects in adolescents (12 to under 18 years): loss of bone density.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard. By reporting side effects you can help provide more information on the safety of this medicine.

5 How to store Dimetrum

Keep this medicine out of the sight and reach of children.

Store in the outer carton box packaging to protect from light.

Do not use this medicine after the expiry date which is stated on the carton and blister after EXP. The expiry date refers to the last day of that month.

If the tablet becomes discoloured or shows any other signs of deterioration, you should seek the advice of your pharmacist who will tell you what to do.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6 Contents of the pack and other information

What Dimetrum contains

- The active substance is dienogest. Each tablet contains 2 mg dienogest.
- The other ingredients are lactose monohydrate, povidone K30, pregelatinized maize starch, microcrystalline cellulose, crospovidone, colloidal anhydrous silica, magnesium stearate.

What Dimetrum looks like and contents of the pack

Dimetrum tablets are white to slightly yellowish round tablets marked with 'D2' on one side and without marking on the other side, with a diameter of approximately 7 mm.

They are supplied in a blister pack containing 28 tablets.

Manufacturer and Licence Holder

This medicine is manufactured by CYNDEA PHARMA SL, Poligono Industrial Emiliano Revilla Sanz, Avenida de Agreda 31, Olvega 42110 (Soria), Spain and is procured from within the EU and repackaged by the Product Licence Holder: Lexon (UK) Limited, Unit 18, Oxleasow Road, East Moons Moat, Redditch, Worcestershire, B98 0RE.

POM

PL 15184/2581

Dimetrum 2 mg tablets

Dimetrum is a registered trademark of Besins Healthcare Luxembourg S.A.R.L

Leaflet revision date: 01/04/25

Blind or partially sighted?

Is this leaflet hard to see or read?

Phone Lexon (UK) Limited,

Tel: 01527 505414 to obtain the leaflet

in a format suitable for you

Dienogest 2 mg tablets

Patient Information Leaflet

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See Section 4.

The name of your medicine is Dienogest 2 mg tablets but will be referred to as Dienogest throughout this leaflet.

What is in this leaflet:

- 1 What Dienogest is and what it is used for
- 2 What you need to know before you take Dienogest
- 3 How to take Dienogest
- 4 Possible side effects
- 5 How to store Dienogest
- 6 Contents of the pack and other information

1 What Dienogest is and what it is used for

Dienogest is a preparation for the treatment of endometriosis (painful symptoms due to displaced tissue of the lining of the womb). Dienogest contains a hormone, the progestogen dienogest.

2 What you need to know before you take Dienogest

Do not take Dienogest if you:

- * are suffering from a blood clot (thromboembolic disorder) in your veins. This may occur, for example, in the blood vessels of the legs (deep vein thrombosis) or the lungs (pulmonary embolism). See also "*Dienogest and venous blood clots*" below
- * have or have ever had a **severe arterial disease**, including cardiovascular disease, such as a **heart attack, stroke** or **heart disease** which causes a reduced blood supply (angina pectoris). See also "*Dienogest and arterial blood clots*" below
- * have **diabetes** with blood vessel damage
- * have or have ever had **severe liver disease** (and your liver function values have not returned to normal). Symptoms of liver disease may be yellowing of the skin and/or itching of the whole body,
- * have or have ever had a **benign or malignant liver tumour**
- * suffer or have ever suffered, or if it is suspected that you suffer from a **malignant** sex-hormone dependent tumour such as cancer of the breast or the genital organs
- * have any unexplained **vaginal bleeding**
- * are **allergic (hypersensitive)** to dienogest or any of the other ingredients of this medicine (see *Section 6, and end of Section 2*).

If any of these conditions appear for the first time while using Dienogest, stop taking it at once and consult your doctor.

Warnings and precautions

You must not use hormonal contraceptives of any form (tablet, patch, intrauterine system) while taking Dienogest.

Dienogest is NOT a contraceptive. If you want to prevent pregnancy, you should use condoms or other non-hormonal contraceptive precautions.

In some situations you need to take special care while using Dienogest, and your doctor may need to examine you regularly. Tell your doctor if any of the following conditions applies to you:

If you:

- have ever had a **blood clot** (venous thromboembolism) or anyone in your immediate family has had a blood clot at a relatively early age
- have a close relative who has had **breast cancer**
- have ever suffered from **depression**
- have **high blood pressure** or develop high blood pressure while taking Dienogest
- develop a **liver disease** while taking Dienogest. Symptoms may include yellowing of the skin or eyes or itching all over your body. Inform your doctor also if such symptoms occurred during a previous pregnancy
- have diabetes or had **diabetes** temporarily during previous pregnancy
- have ever had **chloasma** (golden-brown patches on the skin, particularly of the face); if so, avoid too much exposure to the sun or ultraviolet radiation
- suffer from **pain in your lower abdomen** while taking Dienogest.

While taking Dienogest your chance of becoming pregnant is reduced because Dienogest may affect ovulation.

If you become pregnant while taking Dienogest you are at a **slightly increased risk** of having an extrauterine pregnancy (the embryo develops outside the womb). Tell your doctor before you start taking Dienogest, if you had an extrauterine pregnancy in the past or have an impaired function of the Fallopian tubes.

Dienogest and serious uterine bleeding

Uterine bleeding, for example in women with a condition where the mucous membrane of your uterus (endometrium) grows into the muscle layer of your uterus, called adenomyosis uteri or **benign tumours** of the womb sometimes called uterine fibroids (uterine leiomyomata), may become worse with the use of Dienogest. If bleeding is heavy and continuous over time, this may lead to low red blood cell levels (anemia), which may be severe in some cases. In the event of anemia, you should discuss with your doctor if you should stop taking Dienogest.

Dienogest and changes in bleeding pattern

Most women treated with Dienogest experience changes in their menstrual bleeding pattern (see section 4, possible side effects).

Dienogest and venous blood clots

Some studies indicate that there may be a slight, but not statistically significant, increased risk of a **blood clot in the legs (venous thromboembolism)** associated with the use of preparations with progestagens like Dienogest. Very rarely, blood clots may cause serious permanent disabilities or may even be fatal.

The risk of a **venous blood** clot increases:

- * with increasing age
- * if you are overweight
- * if you or one of your close relatives had a blood clot in the leg (thrombosis), lung (pulmonary embolism), or other organ at a young age
- * if you must have surgery, if you have had a serious accident or if you are immobilized for a long time. It is important to tell your doctor in advance that you are using Dienogest as the treatment may have to be stopped. Your doctor will tell you when to start Dienogest again. This is usually about two weeks after you are back on your feet.

Dienogest and arterial blood clots

There is little evidence for an association between preparations with progestagens like Dienogest and an increased risk of a blood clot in, for example, the bloodvessels of the heart (heart attack) or the brain (stroke). In women with hypertension the risk of stroke may be slightly enhanced by these preparations.

The risk of an **arterial blood** clot increases:

- * **if you smoke. You are strongly advised to stop smoking when you use Dienogest, especially if you are older than 35 years.**
- * if you are overweight
- * if one of your close relatives had a heart attack or stroke at a young age
- * if you have high blood pressure

Talk to your doctor before taking Dienogest

Stop taking Dienogest and contact your doctor immediately if you notice possible signs of a blood clot, such as:

- * severe pain and/or swelling in one of your legs
- * sudden severe pain in the chest which may reach the left arm
- * sudden breathlessness
- * sudden cough without an obvious cause
- * any unusual, severe or long-lasting headache or worsening of migraine
- * partial or complete blindness or double vision
- * difficulty in speaking or inability to speak
- * giddiness or fainting
- * weakness, strange feeling, or numbness in any part of the body.

Dienogest and cancer

It is not clear from the data currently available whether or not Dienogest increases the risk of breast cancer. Breast cancer has been observed slightly more often in women taking hormones compared to those not taking hormones, but it is not known whether this is caused by the treatment. For example, it may be that more tumours are detected and detected earlier in women taking hormones because they are examined by their doctor more often. The occurrence of breast tumours becomes gradually less after stopping the hormone treatment. **It is important to regularly check your breasts** and you should contact your doctor if you feel any lump. In rare cases, benign liver tumours, and in even fewer cases malignant liver tumours have been reported in women taking hormones. Contact your doctor if you have unusually severe stomach pain.

Dienogest and osteoporosis

Changes in bone mineral density (BMD)

The use of Dienogest may affect the strength of the bone of adolescents (12 to under 18 years). If you are under 18 your doctor will, therefore, carefully weigh the benefits and risks of using Dienogest for you as an individual patient, taking into account possible risk factors for bone loss (osteoporosis).

If you use Dienogest, it will help your bones if you have an adequate intake of calcium and vitamin D either via your food or via supplements.

If you have an increased risk of getting osteoporosis (weakening of bones due to loss of bone minerals), your doctor will carefully weigh the risks and benefits of treatment with Dienogest because Dienogest has a moderate suppressing effect on the production of oestrogen (another type of female hormone) by your body.

Other medicines and Dienogest

Always tell your doctor which medicines or herbal products you are already using. Also tell any other doctor or dentist who prescribes another medicine (or the pharmacist) that you are taking Dimetrum.

Some medicines can have an influence on the blood levels of Dienogest and can make it less effective, or can cause undesirable effects.

These include:

- * medicines used for the treatment of:
 - * **epilepsy** (e.g. phenytoin, barbiturates, primidone, carbamazepine, oxcarbazepine, topiramate, felbamate)
 - * **tuberculosis** (e.g. rifampicin)
 - * **HIV and Hepatitis C Virus** infections (so-called protease inhibitors and non-nucleoside reverse transcriptase inhibitors such as ritonavir, nevirapine, efavirenz)
 - * **fungal infections** (griseofulvin, ketoconazole)
- * the herbal remedy **St. John's wort**.

Dienogest 2 mg tablets

Patient Information Leaflet (continued)

Ask your doctor or pharmacist for advice before taking any medicine.

Dienogest with food and drink

During Dienogest treatment, you should avoid drinking grapefruit juice, because this may increase the levels of Dienogest in your blood. This may increase the risk of getting side effects.

Dienogest can be taken with or without food.

Laboratory tests

If you need a blood test, tell your doctor or the laboratory staff that you are taking Dienogest, because Dienogest can affect the results of some tests.

Pregnancy, breast-feeding and fertility

Do not take Dienogest if you are pregnant or breast-feeding.

Driving and using machines

No effects on the ability to drive and use machines have been observed in users of Dienogest.

Dienogest contains lactose

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking Dienogest.

Children and adolescents

Dienogest is not for use in girls before menarche (first menstrual bleeding). The use of Dienogest may affect the strength of the bone of adolescents (12 to under 18 years). If you are under 18 your doctor will, therefore, carefully weigh the benefits and risks of using Dienogest for you as an individual patient, taking into account possible risk factors for bone loss (osteoporosis).

3 How to take Dienogest

Always take Dienogest exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure. For adults, the usual dose is 1 tablet per day.

The following statements apply to Dienogest unless otherwise prescribed by your doctor. Please follow these instructions, otherwise you will not fully benefit from Dienogest.

You can start treatment with Dienogest on any day of your natural cycle.

Adults: take one tablet every day, preferably at the same time with some liquid as needed. When a pack is finished the next one should be started without interruption. Continue to take the tablets also on days of menstrual bleeding.

If you take more Dienogest than you should

There have been no reports of serious harmful effects from taking too many Dienogest at one time. However, if you are concerned, contact your doctor.

If you forget to take Dienogest or suffer from vomiting or diarrhoea

Dienogest will be less effective if you miss a tablet. If you miss one or more tablets, take one tablet only as soon as you remember, and then continue next day taking the tablet at your usual time.

If you vomit within 3-4 hours of taking Dienogest or you have severe diarrhoea, there is a risk that the active substance in the tablet will not be taken up by your body. The situation is almost the same as forgetting a tablet. After vomiting or diarrhoea within 3-4 hours of taking Dienogest, you should take another tablet as soon as possible.

Do not take a double dose to make up for a forgotten tablet.

If you stop taking Dienogest

If you stop taking Dienogest, your original endometriosis symptoms may return.

4 Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them. These effects are more common during the first months after start of intake of Dienogest and usually disappear with continued use. You may also experience changes in your bleeding pattern, such as spotting, irregular bleeding or your periods may stop completely.

Common (affecting between 1 and 10 in every 100 users)

- weight gain
- depressed mood, problems sleeping, nervousness, loss of interest in sex, or changed mood
- headache or migraine
- nausea, abdominal pain, wind, swollen tummy or vomiting
- acne or hair loss
- back pain
- breast discomfort, ovarian cyst or hot flushes
- uterine/vaginal bleeding including spotting
- weakness or irritability.

Uncommon (affecting between 1 and 10 in every 1,000 users)

- anemia
- weight loss or increase in appetite
- anxiety, depression or mood swings
- imbalance in the autonomic nervous system (controls unconscious bodily functions, e.g. perspiration) or disturbed attention
- dry eye
- tinnitus
- unspecific circulatory problems or uncommon palpitations
- low blood pressure

- shortness of breath
- diarrhoea, constipation, abdominal discomfort, inflammation the stomach and intestines, (gastrointestinal inflammation), inflammation of the gums (gingivitis)
- dry skin, excessive sweating, severe itching of the whole body, male pattern hair growth (hirsutism), brittle nails, dandruff, dermatitis, abnormal hair growth, hypersensitive response to light or problems with skin pigmentation
- pains in your bones, muscle spasms, pains and/or a sensation of heaviness in your arms and hands or legs and feet
- urinary tract infection
- vaginal thrush, dryness of the genital area, vaginal discharge, pelvic pain, atrophic inflammation of the genitals with discharge (atrophic vulvovaginitis), or a lump or lumps in the breast
- swelling due to fluid retention.

Additional side effects in adolescents (12 to under 18 years): loss of bone density.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard. By reporting side effects you can help provide more information on the safety of this medicine.

5 How to store Dienogest

Keep this medicine out of the sight and reach of children.

Store in the outer carton box packaging to protect from light.

Do not use this medicine after the expiry date which is stated on the carton and blister after EXP. The expiry date refers to the last day of that month.

If the tablet becomes discoloured or shows any other signs of deterioration, you should seek the advice of your pharmacist who will tell you what to do.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6 Contents of the pack and other information

What Dienogest contains

- The active substance is dienogest. Each tablet contains 2 mg dienogest.
- The other ingredients are lactose monohydrate, povidone K30, pregelatinized maize starch, microcrystalline cellulose, crospovidone, colloidal anhydrous silica, magnesium stearate.

What Dienogest looks like and contents of the pack

Dienogest tablets are white to slightly yellowish round tablets marked with 'D2' on one side and without marking on the other side, with a diameter of approximately 7 mm.

They are supplied in a blister pack containing 28 tablets.

Manufacturer and Licence Holder

This medicine is manufactured by CYNDEA PHARMA SL, Poligono Industrial Emiliano Revilla Sanz, Avenida de Agreda 31, Olvega 42110 (Soria), Spain and is procured from within the EU and repackaged by the Product Licence Holder: Lexon (UK) Limited, Unit 18, Oxleasow Road, East Moons Moat, Redditch, Worcestershire, B98 0RE.

POM

PL 15184/2581

Dienogest 2 mg tablets

Leaflet revision date: 01/04/25

Blind or partially sighted?

Is this leaflet hard to see or read?

Phone Lexon (UK) Limited,

Tel: 01527 505414 to obtain the leaflet

in a format suitable for you