

than 50 “dips”. (Only a small area or small number of warts should be treated at any one time).



5. Firmly replace the top on the bottle and wash your hands.
6. Steps 1-5 should be repeated 12 hours later on the same day, and then twice a day (once in the morning and once in the evening) for the next 2 days so that you **have 6 treatments in 3 days. No more treatments should be used in the first week.**

Extended treatment period:

If your warts have not disappeared or improved after the one week treatment, your doctor may advise you to repeat the series of 6 treatments in 3 days for a second time (starting on the same day of the week as when you first used Condylone). If necessary, this treatment programme may need to be repeated a third, fourth and fifth week.

If the solution gets into your eyes, immediately bathe them with water or with an appropriate eye cleaning solution if available. Do not keep bathing the eyes as this may cause irritation. In any case, inform your doctor straight away.

If you use more Condylone than you should

If you accidentally apply too much medicine or get it in the wrong place, wash the affected area well with soap and water. There is a risk of systemic toxicity if you use excessive amounts of Condylone for a long time. If you have done this, you should tell your doctor. If someone happens to drink the solution, contact their doctor or nearest Accident & Emergency Department immediately.

If you forget to use Condylone

If you miss a treatment, apply it as soon as you remember. If it is nearly time for your next application, skip the missed dose and continue using Condylone as your doctor has directed.

If you have any further questions on the use of this product, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, Condylone can cause side effects, although not everybody gets them.

Side effects of Condylone include:

Not known (frequency cannot be estimated from the available data)

- Inflammation of the foreskin and head of the penis
- Local irritation of the vagina
- Irritation of your warts shortly after applying the solution. This may include itching, burning, pain, redness or lesions (abnormal growths) on the skin
- Although it is unlikely, some people may experience an allergic reaction to the active ingredient podophyllotoxin – this may be severe and painful redness or irritation, or any other unexpected pain.

Reporting of side effects

If you get any side effects, talk to your doctor. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Condylone

Keep this medicine out of the sight and reach of children.

The product should be stored below 25°C, away from naked flames.

Do not use Condylone after 6 weeks from the date you first opened the bottle.

Do not use this medicine after the expiry date which is stated on the carton

and bottle after ‘EXP’. The expiry date refers to the last day of that month.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to dispose of medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Condylone contains

-The **active substance** is: podophyllotoxin 5 mg/ml.

-The **other substances** are: lactic acid, sodium lactate and alcohol (96%).

(See Section 2 ‘Condylone contains alcohol’.)

What Condylone looks like and contents of the pack

Each 5ml pack of Condylone contains 3.5ml of clear, colourless solution in a glass bottle fitted with a child resistant closure, as well as a supply of special plastic applicator sticks.

Marketing Authorisation Holder

Takeda UK Limited
1 Kingdom Street,
London,
W2 6BD,
United Kingdom

Manufacturer

Takeda Austria GmbH
St Peter-Strasse 25
4020 Linz
Austria

This leaflet was last revised in November 2020.

1128342
2011 
Code 736