Package leaflet: Information for the patient

P3819T

Fexofenadine Hydrochloride 120mg Film-coated Tablets

(Fexofenadine hydrochloride)

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.
- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

The name of your medicine is Fexofenadine Hydrochloride 120mg Film-coated Tablets (referred to as Fexofenadine Tablets throughout this leaflet).

What is in this leaflet

1. What Fexofenadine Tablets is and what it is used for
2. What you need to know before you take Fexofenadine Tablets
3. How to take Fexofenadine Tablets
4. Possible side effects
5. How to store Fexofenadine Tablets
6. Contents of the pack and other information

1. What Fexofenadine Tablets is and what it is used for

Fexofenadine Tablets contains fexofenadine hydrochloride, which is a non-drowsy antihistamine.

Fexofenadine Tablets are used in adults and adolescents of 12 years and older to relieve the symptoms associated with long term allergic skin reactions (chronic idiopathic urticaria) such as itching, swelling and rashes.

2. What you need to know before you take Fexofenadine

Tablets Do not take Fexofenadine Tablets:
- if you are allergic to fexofenadine or any of the other ingredients of this medicine (listed in section 6).

Warnings and precautions

Talk to your doctor or pharmacist before taking Fexofenadine Tablets if:
- you have problems with your liver or kidneys
- you have or ever had heart disease, since this kind of medicine may lead to a fast or irregular heart beat
- you are elderly

If any of these apply to you, or if you are not sure, tell your doctor before taking Fexofenadine Tablets.

Other medicines and Fexofenadine Tablets

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

If you are taking apalutamide (a medicine to treat prostate cancer), as the effect of fexofenadine may be decreased.

Indigestion remedies containing aluminium and magnesium may affect the action of Fexofenadine Tablets, by lowering the amount of medicinal product absorbed.

It is recommended that you leave about 2 hours between the time that you take Fexofenadine Tablets and your indigestion remedy.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

Do not take Fexofenadine Tablets if you are pregnant, unless necessary.

Fexofenadine Tablets is not recommended during breast-feeding.

Driving and using machines

Fexofenadine Tablets is unlikely to affect your ability to drive or operate machinery. However, you should check that these tablets do not make you feel sleepy or dizzy before driving or operating machinery.

Fexofenadine Tablets contains lactose

Fexofenadine Tablets contains lactose; if you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

3. How to use Fexofenadine Tablets

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

For adults and children aged 12 years and over

The recommended dose is one tablet (120mg) daily.

Take your tablet with water before a meal.

This medicine starts to relieve your symptoms within 1 hour and lasts for 24 hours.

If you take more Fexofenadine Tablets than you should

If you take too many tablets, contact your doctor or the nearest hospital emergency department immediately. Symptoms of an overdose in adults are dizziness, drowsiness, fatigue and dry mouth.

If you forget to take Fexofenadine Tablets

Do not take a double dose to make up for a forgotten tablet.

Take the next dose at the usual time as prescribed by your doctor.

Crescent PHARMA LIMITED

Product:	Fexofenadine Hydrochloride 120mg Film-coated Tablets
Item Code:	P3819T
Size:	140 x 285 mm
Min. Point size:	9pt
O. Artwork	19/02/26
Proof No.	2
Amendment Date:	19/02/26

Colours Used	
	Process Black
	Keyline - Does not print

If you stop taking Fexofenadine Tablets
Tell your doctor if you want to stop taking Fexofenadine Tablets before you have finished your course of treatment.

If you stop taking Fexofenadine Tablets earlier than planned, your symptoms may return.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Tell your doctor immediately and stop taking Fexofenadine Tablets if you experience:
- swelling of the face, lips, tongue or throat and difficulty breathing, as these may be signs of a serious allergic reaction.

The following undesirable effects have been reported in clinical trials, with an incidence similar to those observed in patient who did not receive the drug (placebo).

Common side effects (may affect up to 1 in 10 people):
- headache
- drowsiness
- feeling sick (nausea)
- dizziness.

Uncommon side effects (may affect up to 1 in 100 people):
- tiredness/sleepiness.

Additional side effects (frequency not known: cannot be estimated from the available data) which may occur are:
- difficulty sleeping (insomnia)
- sleeping disorders
- bad dreams
- nervousness
- fast or irregular heart beat
- diarrhoea
- skin rash and itching
- hives
- serious allergic reactions which can cause swelling of the face, lips, tongue or throat, flushing, chest tightness and difficulty breathing.
- Blurred vision

Reporting of side effects
If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Fexofenadine Tablets

- Keep this medicine out of the sight and reach of children.
- Do not use this medicine after the expiry date which is stated on the carton after EXP. The expiry date refers to the last day of that month.
- This medicinal product does not require any special storage conditions.
- Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

6. Contents of the pack and other information

What Fexofenadine Tablets contains
The active substance is fexofenadine hydrochloride.

Each film-coated tablet contains 120mg of fexofenadine hydrochloride.

The other ingredients are:
Lactose monohydrate, Low-substituted-Hydroxypropyl Cellulose (E463), Pregelatinised starch, Colloidal anhydrous silica (E551), Microcrystalline cellulose (E460), Croscarmellose sodium (E468), Magnesium stearate (E572), Hypromellose, Povidone, Titanium dioxide (E171), Iron oxide Red (E172), Iron oxide Yellow (E172) and Macrogol 400.

What Fexofenadine Tablets looks like and contents of the pack
Fexofenadine 120mg Tablets: Peach coloured capsule shaped film-coated tablets, debossed as '120' on one side and 'FX'on other side.

Fexofenadine 120mg tablets are provided in blisters of Alu-PVC/PVDC or Alu /PVC/PE/ACLAR in pack size of 7, 10, 15, 20, 30, 50, and 100 tablets.

Fexofenadine 120mg Tablets are provided in white opaque HDPE container with screw neck and child resistant white opaque polypropylene outer closure in pack size of 500 and 1000 tablets.

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer
Crescent Pharma Ltd., Key House, Sarum Hill, Basingstoke, RG21 8SR, UK.

This leaflet was last revised in February 2026.

pharmacode

P3819T

Crescent PHARMA LIMITED

Product:	Fexofenadine Hydrochloride 120mg Film-coated Tablets
Item Code:	P3819T
Size:	140 x 285 mm
Min. Point size:	9pt
O. Artwork	19/02/26
Proof No.	2
Amendment Date:	19/02/26

Colours Used	
	Process Black
	Keyline - Does not print

Package leaflet: Information for the patient

P3817T

Fexofenadine Hydrochloride 120mg Film-coated Tablets

(Fexofenadine hydrochloride)

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.
- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

The name of your medicine is Fexofenadine Hydrochloride 120mg Film-coated Tablets (referred to as Fexofenadine Tablets throughout this leaflet).

What is in this leaflet

1. What Fexofenadine Tablets is and what it is used for
2. What you need to know before you take Fexofenadine Tablets
3. How to take Fexofenadine Tablets
4. Possible side effects
5. How to store Fexofenadine Tablets
6. Contents of the pack and other information

1. What Fexofenadine Tablets is and what it is used for

Fexofenadine Tablets contains fexofenadine hydrochloride, which is a non-drowsy antihistamine.

Fexofenadine Tablets are used in adults and adolescents of 12 years and older to relieve the symptoms associated with long term allergic skin reactions (chronic idiopathic urticaria) such as itching, swelling and rashes.

2. What you need to know before you take Fexofenadine

Tablets Do not take Fexofenadine Tablets:
- if you are allergic to fexofenadine or any of the other ingredients of this medicine (listed in section 6).

Warnings and precautions

Talk to your doctor or pharmacist before taking Fexofenadine Tablets if:
- you have problems with your liver or kidneys
- you have or ever had heart disease, since this kind of medicine may lead to a fast or irregular heart beat
- you are elderly

If any of these apply to you, or if you are not sure, tell your doctor before taking Fexofenadine Tablets.

Other medicines and Fexofenadine Tablets

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

If you are taking apalutamide (a medicine to treat prostate cancer), as the effect of fexofenadine may be decreased.

Indigestion remedies containing aluminium and magnesium may affect the action of Fexofenadine Tablets, by lowering the amount of medicinal product absorbed.

It is recommended that you leave about 2 hours between the time that you take Fexofenadine Tablets and your indigestion remedy.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

Do not take Fexofenadine Tablets if you are pregnant, unless necessary.

Fexofenadine Tablets is not recommended during breast-feeding.

Driving and using machines

Fexofenadine Tablets is unlikely to affect your ability to drive or operate machinery. However, you should check that these tablets do not make you feel sleepy or dizzy before driving or operating machinery.

Fexofenadine Tablets contains lactose

Fexofenadine Tablets contains lactose; if you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

3. How to use Fexofenadine Tablets

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

For adults and children aged 12 years and over

The recommended dose is one tablet (120mg) daily.

Take your tablet with water before a meal.

This medicine starts to relieve your symptoms within 1 hour and lasts for 24 hours.

If you take more Fexofenadine Tablets than you should

If you take too many tablets, contact your doctor or the nearest hospital emergency department immediately. Symptoms of an overdose in adults are dizziness, drowsiness, fatigue and dry mouth.

If you forget to take Fexofenadine Tablets

Do not take a double dose to make up for a forgotten tablet.

Take the next dose at the usual time as prescribed by your doctor.

Crescent PHARMA LIMITED

Product:	Fexofenadine 120mg Tablets
Item Code:	P3817T
Size:	140 x 280 mm
Min. Point size:	9pt
O. Artwork	19/02/26
Proof No.	2
Amendment Date:	20/02/26

Colours Used	
	Process Black
	Keyline - Does not print

If you stop taking Fexofenadine Tablets
Tell your doctor if you want to stop taking Fexofenadine Tablets before you have finished your course of treatment.

If you stop taking Fexofenadine Tablets earlier than planned, your symptoms may return.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Tell your doctor immediately and stop taking Fexofenadine Tablets if you experience:
- swelling of the face, lips, tongue or throat and difficulty breathing, as these may be signs of a serious allergic reaction.

The following undesirable effects have been reported in clinical trials, with an incidence similar to those observed in patient who did not receive the drug (placebo).

Common side effects (may affect up to 1 in 10 people):
- headache
- drowsiness
- feeling sick (nausea)
- dizziness.

Uncommon side effects (may affect up to 1 in 100 people):
- tiredness/sleepiness.

Additional side effects (frequency not known: cannot be estimated from the available data) which may occur are:
- difficulty sleeping (insomnia)
- sleeping disorders
- bad dreams
- nervousness
- fast or irregular heart beat
- diarrhoea
- skin rash and itching
- hives
- serious allergic reactions which can cause swelling of the face, lips, tongue or throat, flushing, chest tightness and difficulty breathing.
- Blurred vision

Reporting of side effects
If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Fexofenadine Tablets

- Keep this medicine out of the sight and reach of children.
- Do not use this medicine after the expiry date which is stated on the carton after EXP. The expiry date refers to the last day of that month.
- This medicinal product does not require any special storage conditions.
- Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

6. Contents of the pack and other information

What Fexofenadine Tablets contains
The active substance is fexofenadine hydrochloride.

Each film-coated tablet contains 120mg of fexofenadine hydrochloride.

The other ingredients are:
Lactose monohydrate, Low-substituted-Hydroxypropyl Cellulose (E463), Pregelatinised starch, Colloidal anhydrous silica (E551), Microcrystalline cellulose (E460), Croscarmellose sodium (E468), Magnesium stearate (E572), Hypromellose, Povidone, Titanium dioxide (E171), Iron oxide Red (E172), Iron oxide Yellow (E172) and Macrogol 400.

What Fexofenadine Tablets looks like and contents of the pack
Fexofenadine 120mg Tablets: Peach coloured capsule shaped film-coated tablets, debossed as '120' on one side and 'FX'on other side.

Fexofenadine 120mg tablets are provided in blisters of Alu-PVC/PVDC or Alu /PVC/ PE/ACLAR in pack size of 7, 10, 15, 20, 30, 50, and 100 tablets.

Fexofenadine 120mg Tablets are provided in white opaque HDPE container with screw neck and child resistant white opaque polypropylene outer closure in pack size of 500 and 1000 tablets.

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer
Crescent Pharma Ltd., Key House, Sarum Hill, Basingstoke, RG21 8SR, UK.

This leaflet was last revised in February 2026.

P3817T

Crescent PHARMA LIMITED

Product:	Fexofenadine 120mg Tablets
Item Code:	P3817T
Size:	140 x 280 mm
Min. Point size:	9pt
O. Artwork	19/02/26
Proof No.	2
Amendment Date:	20/02/26

Colours Used	
	Process Black
	Keyline - Does not print

Package leaflet: Information for the patient

P3966T

Fexofenadine Hydrochloride 120mg Film-coated Tablets

(Fexofenadine hydrochloride)

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.
- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

The name of your medicine is Fexofenadine Hydrochloride 120mg Film-coated Tablets (referred to as Fexofenadine Tablets throughout this leaflet).

What is in this leaflet

1. What Fexofenadine Tablets is and what it is used for
2. What you need to know before you take Fexofenadine Tablets
3. How to take Fexofenadine Tablets
4. Possible side effects
5. How to store Fexofenadine Tablets
6. Contents of the pack and other information

1. What Fexofenadine Tablets is and what it is used for

Fexofenadine Tablets contains fexofenadine hydrochloride, which is a non-drowsy antihistamine.

Fexofenadine Tablets are used in adults and adolescents of 12 years and older to relieve the symptoms associated with long term allergic skin reactions (chronic idiopathic urticaria) such as itching, swelling and rashes.

2. What you need to know before you take Fexofenadine

Tablets Do not take Fexofenadine Tablets:
- if you are allergic to fexofenadine or any of the other ingredients of this medicine (listed in section 6).

Warnings and precautions

Talk to your doctor or pharmacist before taking Fexofenadine Tablets if:
- you have problems with your liver or kidneys
- you have or ever had heart disease, since this kind of medicine may lead to a fast or irregular heart beat
- you are elderly

If any of these apply to you, or if you are not sure, tell your doctor before taking Fexofenadine Tablets.

Other medicines and Fexofenadine Tablets

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

If you are taking apalutamide (a medicine to treat prostate cancer), as the effect offexofenadine may be decreased.

Indigestion remedies containing aluminium and magnesium may affect the action of Fexofenadine Tablets, by lowering the amount of medicinal product absorbed.

It is recommended that you leave about 2 hours between the time that you take Fexofenadine Tablets and your indigestion remedy.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

Do not take Fexofenadine Tablets if you are pregnant, unless necessary.

Fexofenadine Tablets is not recommended during breast-feeding.

Driving and using machines

Fexofenadine Tablets is unlikely to affect your ability to drive or operate machinery. However, you should check that these tablets do not make you feel sleepy or dizzy before driving or operating machinery.

Fexofenadine Tablets contains lactose

Fexofenadine Tablets contains lactose; if you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

3. How to use Fexofenadine Tablets

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

For adults and children aged 12 years and over

The recommended dose is one tablet (120mg) daily.

Take your tablet with water before a meal.

This medicine starts to relieve your symptoms within 1 hour and lasts for 24 hours.

If you take more Fexofenadine Tablets than you should

If you take too many tablets, contact your doctor or the nearest hospital emergency department immediately. Symptoms of an overdose in adults are dizziness, drowsiness, fatigue and dry mouth.

If you forget to take Fexofenadine Tablets

Do not take a double dose to make up for a forgotten tablet.

Take the next dose at the usual time as prescribed by your doctor.

If you stop taking Fexofenadine Tablets

Tell your doctor if you want to stop taking Fexofenadine Tablets before you have finished your course of treatment.

If you stop taking Fexofenadine Tablets earlier than planned, your symptoms

Crescent PHARMA LIMITED

Product:	Fexofenadine Hydrochloride 120mg Film-coated Tablets (Jingxin Pharma)
Item Code:	P3966T
Size:	140 x 285 mm
Min. Point size:	9pt
O. Artwork	06/05/26
Proof No.	1
Amendment Date	

Colours Used	
	Process Black
	Keyline - Does not print

may return.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Tell your doctor immediately and stop taking Fexofenadine Tablets if you experience:

- swelling of the face, lips, tongue or throat and difficulty breathing, as these may be signs of a serious allergic reaction.

The following undesirable effects have been reported in clinical trials, with an incidence similar to those observed in patient who did not receive the drug (placebo).

Common side effects (may affect up to 1 in 10 people):

- headache
- drowsiness
- feeling sick (nausea)
- dizziness.

Uncommon side effects (may affect up to 1 in 100 people):

- tiredness/sleepiness.

Additional side effects (frequency not known: cannot be estimated from the available data) which may occur are:

- difficulty sleeping (insomnia)
- sleeping disorders
- bad dreams
- nervousness
- fast or irregular heart beat
- diarrhoea
- skin rash and itching
- hives
- serious allergic reactions which can cause swelling of the face, lips, tongue or throat, flushing, chest tightness and difficulty breathing.
- Blurred vision

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Fexofenadine Tablets

- Keep this medicine out of the sight and reach of children.
- Do not use this medicine after the expiry date which is stated on the carton after EXP. The expiry date refers to the last day of that month.
- This medicinal product does not require any special storage conditions.
- Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

6. Contents of the pack and other information

What Fexofenadine Tablets contains

The active substance is fexofenadine hydrochloride.

Each film-coated tablet contains 120mg of fexofenadine hydrochloride.

The other ingredients are:
Lactose monohydrate,
Low-substituted-Hydroxypropyl Cellulose (E463), Pregelatinised starch,
Colloidal anhydrous silica (E551),
Microcrystalline cellulose (E460),
Croscarmellose sodium (E468),
Magnesium stearate (E572),
Hypromellose,
Povidone, Titanium dioxide (E171),
Iron oxide Red (E172),
Iron oxide Yellow (E172) and
Macrogol 400.

What Fexofenadine Tablets looks like and contents of the pack

Fexofenadine 120mg Tablets: Peach coloured capsule shaped film-coated tablets, debossed as '120' on one side and 'FX'on other side.

Fexofenadine 120mg tablets are provided in blisters of Alu-PVC/PVDC or Alu /PVC/ PE/ACLAR in pack size of 7, 10, 15, 20, 30, 50, and 100 tablets.

Fexofenadine 120mg Tablets are provided in white opaque HDPE container with screw neck and child resistant white opaque polypropylene outer closure in pack size of 500 and 1000 tablets.

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer

Crescent Pharma Ltd., Key House, Sarum Hill, Basingstoke, RG21 8SR, UK.

This leaflet was last revised in May 2026.

P3966T

Crescent PHARMA LIMITED

Product:	Fexofenadine Hydrochloride 120mg Film-coated Tablets (Jingxin Pharma)
Item Code:	P3966T
Size:	140 x 285 mm
Min. Point size:	9pt
O. Artwork	06/05/26
Proof No.	1
Amendment Date	

Colours Used	
	Process Black
	Keyline - Does not print