

SUMMARY OF PRODUCT CHARACTERISTICS

1 NAME OF THE MEDICINAL PRODUCT

Atropine 1% w/v eye drops, solution

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

One ml eye drops contain 10 mg atropine sulfate.

Excipients with known effect

Benzalkonium chloride 0.2mg/ml is included as preservative

Each 1 ml of this medicinal product contains 9.36mg sodium dihydrogen phosphate dihydrate and 0.950mg disodium phosphate anhydrous.

For the full list of excipients, see section 6.1.

3 PHARMACEUTICAL FORM

Eye drops.

Clear, colourless, solution.

4 CLINICAL PARTICULARS

4.1 Therapeutic indications

Atropine sulfate is an antimuscarinic agent used as a cycloplegic and mydriatic. The eye drops are used in the treatment of iritis and uveitis to immobilise the iris and ciliary muscle and to prevent or break down adhesions.

Since it is a powerful cycloplegic it is used in the determination of refraction in children below six years and children with convergent strabismus

4.2 Posology and method of administration

For topical ocular use.

The depth of the angle of the anterior chamber should be assessed before the product is used.

Adults

Refraction: One or two drops to be instilled into the eye(s) one hour before refracting
Uveitis / iritis: One or two drops to be instilled into the eye(s) to a maximum of 4 times daily.

Elderly

Mydriatics and cycloplegics should only be used with caution in the elderly and others who may have raised intra ocular pressure

Children

Refraction: One drop to be instilled into each eye twice daily for 1 - 3 days prior to the examination

Uveitis / iritis: One drop to be instilled into each eye to a maximum of 3 times daily

4.3 Contraindications

For topical ocular use.

The depth of the angle of the anterior chamber should be assessed before the product is used.

Adults

Refraction: One or two drops to be instilled into the eye(s) one hour before refracting
Uveitis / iritis: One or two drops to be instilled into the eye(s) to a maximum of 4 times daily.

Elderly

Mydriatics and cycloplegics should only be used with caution in the elderly and others who may have raised intra ocular pressure

Children

Refraction: One drop to be instilled into each eye twice daily for 1 - 3 days prior to the examination

Uveitis / iritis: One drop to be instilled into each eye to a maximum of 3 times daily

4.4 Special warnings and precautions for use

It should only be used with caution in patients who may have raised intra ocular pressure.

The product contains benzalkonium chloride solution which may cause eye irritation and soft contact lenses must not be worn during the period of use.

Patients should be warned that antimuscarinic eye drops will temporarily impair vision.

Patients should wash hands after using the eye drops and great care should be taken to

avoid getting the product into the mouth.

Due to the risk of provoking hyperpyrexia, atropine should only be used with great caution when the ambient temperature is high or the patient has a fever.

Care is also required in patients with conditions characterised by tachycardia.

Darkly pigmented iris is more resistant to pupillary dilation and caution should be exercised to avoid overdosage.

The eye drops should be discarded 30 days after first opening.

During use, care should be taken not to touch the dropper nozzle on to the eyelid or any other surface.

The product is for external use only and should be stored out of the sight and reach of children.

This medicine contains less than 1 mmol sodium (23 mg) per drop, that is to say essential 'sodium-free'.

The medicine contains 0,633mg phosphates.

4.5 Interaction with other medicinal products and other forms of interaction

The effects of atropine may be enhanced by the concomitant use of other drugs with antimuscarinic properties. Studies have indicated that the absorption of atropine sulphate appears to be delayed by solutions of higher osmolarity.

4.6 Fertility, Pregnancy and lactation

The safety for use in pregnancy and lactation has not been established, therefore, use only when directed by a physician. No studies have been conducted to evaluate the effect of topical ophthalmic atropine on fertility. No effects on fertility have been reported from rat studies.

4.7 Effects on ability to drive and use machines

May cause transient blurring of vision on instillation. Warn patients not to drive or operate hazardous machinery until vision is clear.

4.8 Undesirable effects

Patients may experience photophobia and eyes should be protected from bright light while the pupils are dilated.

Prolonged use of atropine eye drops may lead to local irritation, transient stinging, hyperaemia, oedema and conjunctivitis. An increase in intra-ocular pressure may occur, especially in patients with closed angle glaucoma.

Hypersensitivity to atropine is not uncommon and may appear as a skin rash or conjunctivitis.

Systemic toxicity may be produced by the instillation of the eye drops especially in infants and the elderly. Reported symptoms include severe ataxia, restlessness, excitement and hallucinations.

Other adverse effects may include a dry mouth with difficulty in swallowing and talking, flushing and a dry skin, transient bradycardia followed by tachycardia, palpitations and arrhythmias, reduced bronchial secretions, urinary urgency and retention and constipation.

Side effects that occur occasionally include confusion (particularly in the elderly), nausea, vomiting and giddiness

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or [search for MHRA Yellow Card in the Google Play or Apple App Store](#).

4.9 Overdose

Symptoms

Systemic reactions to topical atropine are unlikely at normal doses. Symptoms which can occur following an overdose, however, include anticholinergic effects (as listed in section 4.8 above), cardiovascular changes (tachycardia, atrial arrhythmias, atrio-ventricular dissociation) and central nervous system effects (confusion, ataxia, restlessness, hallucination, convulsions).

Treatment

Supportive therapy should be given as required.

5 PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: anticholinergic agents

ATC code: S01FA01

Dilation of the pupil normally occurs within half an hour following local application and lasts for seven days or longer. Paralysis of accommodation in one to three hours with recovery in three to seven days.

5.2 Pharmacokinetic properties

Atropine is readily absorbed from the gastro-intestinal tract and mucous membranes, it is also absorbed from the eye.

It is incompletely metabolised in the liver and is excreted in the urine as unchanged drug and metabolites.

5.3 Preclinical safety data

No additional pre-clinical data of relevance to the prescriber.

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Sodium dihydrogen phosphate dihydrate

Disodium phosphate anhydrous

Benzalkonium chloride solution

Sulfuric acid

Water for injection

6.2 Incompatibilities

Incompatible with alkali, bromide, iodine, mercury and tannic acid

6.3 Shelf life

36 months (unopened)
30 days (once opened)

6.4 Special precautions for storage

This medicinal product does not require any special storage condition

6.5 Nature and contents of container

A white low density polyethylene (LDPE), non-transparent plastic vial (10mL), fitted with a white nozzle and a white plastic cap.

6.6 Special precautions for disposal

None stated

7 MARKETING AUTHORISATION HOLDER

Macure Pharma UK Ltd.
3 Waterhouse Square,
138-142 Holborn, London
EC1N 2SW
United Kingdom

8 MARKETING AUTHORISATION NUMBER(S)

PL 54594/0008

9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

19/01/2022

10 DATE OF REVISION OF THE TEXT

11/04/2025