

Magnaspartate® 243 mg powder for oral solution
magnesium

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

1. What MAGNASPARTATE is and what it is used for
2. What you need to know before you take MAGNASPARTATE
3. How to take MAGNASPARTATE
4. Possible side effects
5. How to store MAGNASPARTATE
6. Contents of the pack and other information

Magnaspartate is a mineral preparation and contains the active ingredient magnesium aspartate dihydrate equivalent to 243 mg (10 mmol) of magnesium. Magnaspartate is used in the treatment and prevention of magnesium deficiency in adults and children aged 2 years and above.

2. What you need to know before you take MAGNASPARTATE

- if you are allergic to magnesium aspartate dihydrate or any of the other ingredients of this medicine (listed in section 6).
- if you have severe kidney problems
- if you have a very slow heart rhythm (called 'bradycardia')

If you have switched from another magnesium medicine to ensure that treatment with Magnaspartate does not make you feel unwell and that you can continue to receive the correct dose of magnesium.

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. This is because Magnaspartate can affect the way some other medicines work. Also, some other medicines can affect the way Magnaspartate works.

- **Cellulose sodium phosphate**, used to prevent kidney stones
- **EDTA**, used in hospital for high calcium levels
- **Flourides**, used for preventing tooth decay
- **Tetracyclines, Nitrofurantoin, Nitroloxline**, antibiotic to treat infections
- **Aminoquinolines**, medicines to treat malaria

This medicine contains less than 1 mmol sodium (23 mg) per sachet, that is to say essentially 'sodium-free'.

3. How to take MAGNASPARTATE

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

The recommended dose is
Adults (over 18 years) 1-2 sachets daily (equal to 243-486 mg magnesium).

Use in children and adolescents
Children and Adolescents (10 to 18 years) One sachet daily (equal to 243 mg magnesium) dissolved in 50-200ml of water, tea, or orange juice.
Children (4 to 10 years) One sachet daily (equal to 243 mg magnesium) dissolved in 50-200ml of water, tea, or orange juice. Alternatively, for a half dosage, use half the volume of 1 sachet dissolved in 50-200ml of water (equal to 121.5 mg magnesium).
Children (2 to 4 years) Half the volume of 1 sachet dissolved in 50-200ml water (equal to 121.5 mg magnesium).

Renal patients:
No dose adjustment is required in mild to moderate renal failure.
The blood magnesium levels of children and adolescents (12 to 18 years) and renal patients need to be monitored at regular intervals.
The duration of magnesium supplementation will depend upon the clinical needs of each patient.

Directions for use:
Dissolve one sachet of Magnaspartate in 50-200ml water, tea or orange juice. Stir until the solution in water is cloudy to transparent. In orange juice or tea inactive particles will be visible.
Drink immediately or within 24 hours if made with bottled water and stored below 25°C.
If you are unable to swallow and have a gastric, duodenal or nasal feeding tube, Magnaspartate in 200ml water can be administered through the tube immediately after preparation or within 24 hours if made with bottled water and stored below 25°C.

If you take more MAGNASPARTATE than you should
No side effects are expected, if your kidneys work properly. Any excessive amount of magnesium will be excreted through your kidneys.
If you have taken more of this medicine than directed, or if a person with severe kidney problems or if a child accidentally has taken this medicine, please contact your doctor or local hospital casualty department for judgement of the risk and advice.

If you forget to take MAGNASPARTATE
Do not take a double dose to make up for a forgotten dose.
If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects
Like all medicines, this medicine can cause side effects, although not everybody gets them.
Uncommon (may affect 1 to 10 users in 1,000):
Soft stools to diarrhoea at the beginning of treatment (harmless and will usually decrease in frequency as the treatment advances).
Very rare (may affect less than 1 user in 10,000):
Fatigue if Magnaspartate is used for a longer period of time.
If you get any side effects such as diarrhoea, you should temporarily interrupt the treatment. After the symptoms improve you can restart treatment, with a reduced dosage.

Reporting of side effects
If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via (see details below).

United Kingdom (Northern Ireland)
Yellow Card Scheme
Website: www.mhra.gov.uk/yellowcard
or search for MHRA Yellow Card in the Google Play or Apple App Store

By reporting side effects, you can help provide more information on the safety of this medicine.

5. How to store MAGNASPARTATE
Do not store above 25°C.
When Magnaspartate has been dissolved in water, orange juice or tea, the solution should be consumed immediately or within 24 hours when dissolved in bottled water and stored below 25°C.
Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the sachet and carton after EXP. The expiry date refers to the last day of that month.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information
What MAGNASPARTATE contains
• The active substance is magnesium. One sachet (6.5 g) contains magnesium aspartate dihydrate 3246 mg; magnesium content: 243 mg.
• The other ingredients are sucrose, citric acid monohydrate, peach/apricot flavour, saccharin sodium, silica, colloidal anhydrous

What MAGNASPARTATE looks like and contents of the pack
White powder with a peach/apricot-like flavour.
Magnaspartate is available in packs of 10 or 20 sachets. Not all pack sizes may be marketed.

Marketing Authorisation Holder
Kora Corporation Ltd t/a Kora Healthcare
20 Harcourt Street, Dublin 2, D02 H364, Ireland.

Manufacturer
Lusomedicamenta Sociedade Tecnica Farmaceutica, S.A.
Estrada Consiglieri Pedroso, 69 - B, Queluz, Barcarena, 2730-055, Portugal

This medicine is authorised in the Member States of the European Economic Area and in the United Kingdom under the following names:
United Kingdom (NI&GB):
Magnaspartate 243 mg powder for oral solution
Belgium, Netherlands, Ireland:
Magnence 243 mg powder for oral solution.

This leaflet was last revised in April 2024

