

Slow-K® 600mg prolonged-release tablets

(potassium chloride)

PATIENT INFORMATION LEAFLET

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

Always take this medicine exactly as described in this leaflet or as your doctor or pharmacist has told you.

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

This medicine is available using above names but will be referred to as Slow-K throughout this leaflet.

What is in this leaflet:

1. What Slow-K is and what it is used for
2. What you need to know before you take Slow-K
3. How to take Slow-K
4. Possible side effects
5. How to store Slow-K
6. Contents of the pack and other information

1. WHAT SLOW-K IS AND WHAT IT IS USED FOR

Slow-K belongs to a group of medicines called potassium supplements.

Slow-K contains potassium and is used to treat or prevent low levels of potassium in your body. Slow-K tablets are prolonged-release tablets which means that the potassium is released slowly from the tablet.

Whilst on this medicine your doctor will want to measure the potassium levels in your blood on a regular basis.

2. WHAT YOU NEED TO KNOW BEFORE YOU TAKE SLOW-K

Do not take Slow-K:

- if you are allergic to potassium chloride or any of the other ingredients of this medicine (listed in section 6)
- if you are allergic to potassium intake due to a genetic disorder such as congenital paramyotonia
- if you have been told by your doctor that you have kidney failure
- if you have Addison's disease (which is a condition where your adrenal gland is not producing enough steroids)
- if you have recently suffered from severe burns
- if you suffer from digestive problems or have difficulty swallowing (due to a narrowing or blockage of your gullet (food pipe) or intestines)
- if you have been told you have metabolic acidosis (a condition caused by increased acid levels in the blood)
- if you are dehydrated (you may feel thirsty with a dry mouth)
- if you have high blood potassium levels (which can cause an abnormal heartbeat)
- if you suffer from a condition called hyporeninaemic hypoaldosteronism (where your body has low levels of an enzyme called renin and a hormone called aldosterone which normally helps to control your blood pressure)
- if you are taking certain types of diuretics (water tablets); either potassium-sparing diuretics (e.g. triamterene and amiloride) or aldosterone antagonists (e.g. spironolactone and eplerenone).

Warnings and precautions

Talk to your doctor or pharmacist before taking Slow-K if you:

- have a stoma or have had any of your bowel removed
- suffer from any condition which affects your bowel movements
- suffer from heart disease (which may cause chest pain, shortness of breath or ankle swelling)
- suffer from kidney problems
- currently have or have previously had a stomach ulcer.

Slow-K is not recommended for use in children.

If any of the above applies to you, or if you are not sure, speak to your doctor or pharmacist before you take Slow-K.

3145
16.03.24[6]

Other medicines and Slow-K

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

In particular, tell your doctor or pharmacist if you are taking:

- Potassium-sparing diuretics such as triamterene and amiloride
- “water tablets” for high blood pressure and certain heart diseases
- Aldosterone antagonists such as spironolactone and eplerenone
- “water tablets” for heart failure and certain liver and kidney problems

If you are taking potassium-sparing diuretics or aldosterone antagonists, stop taking Slow-K. See also the section ‘Do not take Slow-K’.

Tell your doctor or pharmacist if you are taking any of the following as they may increase the risk of high blood potassium levels:

- ACE inhibitors or beta blockers such as captopril or atenolol
- for high blood pressure or heart problems
- Angiotensin-II-receptor antagonists such as losartan or valsartan
- for treatment of high blood pressure, kidney damage due to diabetes and congestive heart failure
- Ciclosporin
- for transplants, rheumatic disease or skin complaints
- Non-steroidal anti-inflammatory medicines (NSAIDs) such as aspirin or naproxen
- for pain relief or rheumatism
- Heparin
- used to thin the blood
- Digoxin
- for an irregular heartbeat
- Direct renin inhibitors such as aliskerin
- for high blood pressure
- Proton pump inhibitors
- for indigestion

Tell your doctor or pharmacist if you are taking anticholinergics such as atropine sulfate or hyoscine butylbromide for stomach spasms or cramps as they can slow down gastrointestinal transit.

Pregnancy and breast-feeding

Your doctor will only give you Slow-K if you really need it. Slow-K is a prolonged-release tablet which means it takes a long time to dissolve. When you are pregnant your digestive system works more slowly and so Slow-K tablets will take longer to pass through your system. This could mean you do not receive the correct amount of potassium from your Slow-K tablets.

Slow-K should not be taken whilst breast-feeding unless your doctor considers that the benefits for you are greater than the risks to the child.

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

Slow-K contains sucrose

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially ‘sodium-free’.

3. HOW TO TAKE SLOW-K

Always take this medicine exactly as described in this leaflet or as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

The recommended dose is 2 to 3 tablets a day. Your doctor will decide when and how to treat you with Slow-K. Your doctor may tell you to take up to 12 tablets a day.

- The tablets should not all be taken at once but spread out during the day.
- Take your tablets with fluid during a meal and whilst sitting upright.
- Swallow them whole.
- Do not break or chew them. This is important as the tablets release potassium slowly. If you chew the tablets, this could damage the tablet and let the potassium out too quickly and this could cause stomach problems.

If you take more Slow-K than you should, you may get stomach ache and feel or be sick. In more severe cases you may suffer from low blood pressure, shock, heart problems and convulsions. If you accidentally take too many tablets, or someone else takes your medicine, you should tell your doctor at once or contact your nearest accident and emergency department. Take this leaflet and / or the container with you so the doctor will know what has been taken.

If you forget to take Slow-K, take it as soon as possible, unless it is almost time to take the next dose and then go on as before. Do not take a double dose to make up for a forgotten dose.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. POSSIBLE SIDE EFFECTS

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Serious side effects

If you experience any of the following side effects, contact your doctor or pharmacist IMMEDIATELY:

- Signs of intestinal bleeding or ulcers such as coughing up blood (however small the amount), blood in your vomit and black stools
- Severe sickness (feeling or being sick)
- Constipation and / or wind
- Swelling of the stomach
- Diarrhoea
- Severe stomach pains.

Other side effects:

- High blood levels of potassium (which can cause an abnormal heartbeat)
- Itching and skin rashes (nettle rash).

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme Website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the **Google Play** or **Apple App Store**. By reporting side effects, you can help provide more information on the safety of this medicine.

5. HOW TO STORE SLOW-K

Keep out of the sight and reach of children.
Do not use this medicine after the expiry date which is stated on the tablet container after EXP. The expiry date refers to the last day of that month.

Do not store above 25°C.

Store in the original package in order to protect from light.

Do not use this medicine if you notice the appearance has changed in any way.

If your doctor decides to stop your treatment, return any unused medicine to the pharmacist. Only keep it if your doctor tells you to.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. CONTENTS OF THE PACK AND OTHER INFORMATION

What Slow-K contains

The active substance is potassium chloride.
Each tablet contains 600mg potassium chloride equivalent to 315mg potassium or approximately 8mmol potassium ion (K⁺).
The other ingredients are cetostearyl alcohol, gelatin, magnesium stearate, acacia, yellow iron oxide (E172), red iron oxide (E172), talc, titanium dioxide (E171), carnauba wax and sucrose.

What Slow-K looks like and contents of the pack

Slow-K is a pale orange, round, biconvex tablet with no marking on either side. They are prolonged-release tablets with a sugar-coating.

Slow-K is available in aluminium and PVC blisters in pack sizes of 30, 40 and 100 tablets.

MANUFACTURER AND PRODUCT LICENCE HOLDER

Manufactured by
Gebro Pharma GmbH, Bahnhofbichl 13, A-6391 Fieberbrunn, Austria.

Procured from within the EU by Product Licence holder:
Star Pharmaceuticals Ltd., 5 Sandridge Close, Harrow, Middlessex, HA1 1XD.
Repackaged by Servipharma Ltd.

P

PL 20636/3145

Leaflet revision and issue date (Ref) 16.03.24[6]

Slow-K is a trademark of Novartis AG.

Blind or partially sighted?
Is this leaflet hard to see or read?
Call 020 8423 2111 to obtain the
leaflet in a format suitable for you.