

## Pentasa® 1g suppositories

(mesalazine)

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

The name of your medicine is Pentasa 1g suppositories but will be referred to as Pentasa throughout this leaflet.

### What is in this leaflet:

1. What Pentasa is and what it is used for
2. What you need to know before you use Pentasa
3. How to use Pentasa
4. Possible side effects
5. How to store Pentasa
6. Contents of the pack and other information

## 1. WHAT PENTASA IS AND WHAT IT IS USED FOR

Each Pentasa contains mesalazine 1g as the active ingredient. Mesalazine belongs to a group of medicines called salicylates.

Pentasa is used to treat inflammation in the gut caused by conditions such as ulcerative colitis. Sometimes ulcerative colitis only affects the back passage (rectum), this is known as ulcerative proctitis. Pentasa are used to treat ulcerative proctitis and are intended for rectal use only.

The suppositories release the active ingredient slowly which then acts locally to reduce the inflammation and help relieve or stop the pain.

## 2. WHAT YOU NEED TO KNOW BEFORE YOU USE PENTASA

### Do Not use Pentasa if you:

- are **allergic (hypersensitive) to mesalazine** or any of the other ingredients of this medicine (see section 6)
- are **allergic to other salicylates e.g. acetylsalicylic acid**
- have **severe liver and/or kidney** problems

### Warnings and precautions

#### Talk to your doctor or pharmacist before using these suppositories if you:

- are allergic to **sulphasalazine** (risk of allergy to salicylates)
- currently have or have previously had **liver** or **kidney** disease
- have a medical condition that can make you prone to **bleeding**
- are on medication that may affect kidney function e.g. Non-steroidal anti-inflammatory drugs (NSAIDs) such as aspirin
- have lung problems, in particular asthma
- suddenly develop abdominal cramps, abdominal pain, fever, severe headache and rash. In such circumstances you should stop taking Pentasa immediately.
- Kidney stones may develop with use of mesalazine. Symptoms may include pain in sides of abdomen and blood in urine. Take care to drink sufficient amount of liquid during treatment with mesalazine
- Have ever developed a severe skin rash or skin peeling, blistering and/or mouth sores after using mesalazine.

Mesalazine may produce red-brown urine discoloration after contact with sodium hypochlorite bleach in the toilet water. It concerns a chemical reaction between mesalazine and bleach and is harmless.

### Take special care with mesalazine:

Serious skin reactions including Drug reaction with eosinophilia and systemic symptoms (DRESS), Stevens-Johnson syndrome (SJS) and toxic epidermal necrolysis (TEN) have been reported in association with mesalazine treatment. Stop using Pentasa and seek medical attention immediately if you notice any of the symptoms related to these serious skin reactions described in section 4.

If you experience strong or recurrent headache, disturbed vision, or ringing or buzzing in the ears contact your doctor immediately.

**While you are on treatment with this medicine, your doctor will normally arrange blood and urine tests to check your kidney function especially at the beginning of treatment.**

You should make sure that you don't become dehydrated, while you are using this medicine. This can occur after severe or prolonged attacks of vomiting and/or diarrhoea, high fever or heavy sweating. If this occurs, you should talk to your doctor or a pharmacist for advice as soon as possible.

### Other medicines and Pentasa

**Tell your doctor or pharmacist if you are taking or have recently taken any other medicines**, including medicines obtained without a prescription. This is especially important if you are taking any of the following:

- **azathioprine** (used after transplantations or to treat auto-immune diseases)
- **6-mercaptopurine or thioguanine** (chemotherapy, used to treat leukaemia)
- certain agents that inhibit blood clotting (medicines for thrombosis or to thin your blood, e.g. warfarin)

### Pregnancy, breast-feeding and fertility

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before using this medicine.

There is limited experience with the use of mesalazine during pregnancy and breast-feeding. Blood disorders have been reported in newborns of mothers being treated with this medicine. The newborn may develop allergic reactions after breast-feeding, e.g. diarrhoea. If the newborn develops diarrhoea, breast-feeding should be discontinued.

### Driving and using machines

This medicine is not known to affect the ability to drive and/or use machines.

## 3. HOW TO USE PENTASA

For Adult use only, not recommended in children.

Always use Pentasa exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure. The suppository should be inserted into the back passage (rectum) only. It is intended to be left in place for as long as possible and is therefore best inserted before going to sleep. To treat an attack of proctitis your doctor will usually prescribe one suppository to be used daily for two to four weeks. To help prevent further attacks it may be used for longer.

### How to insert the suppository

You should make sure that you empty your bowels before inserting the suppository.

- a. Tear along the perforation to remove one suppository blister from a blister strip
- b. Put a protector on the finger with which you will insert the suppository
- c. Remove the suppository from the blister before insertion. This is done by pushing the suppository through the printed foil
- d. Insertion may be made easier if you moisten the suppository with water first
- e. Insert the suppository gently and fully into the back passage. It may help to lie down on one side and raise the other knee.
- f. Remain still for a few minutes after insertion to help keep the suppository in place. If it comes out within 10 minutes, a new one should be inserted
- g. Dispose of the empty blister and used finger protector safely and wash your hands.

### If you use more Pentasa than you should

If you accidentally use too many suppositories, you should go to your nearest emergency department or contact your doctor immediately. Take the pack and any remaining suppositories with you.

### If you forget to use Pentasa

If you forget to use a dose, use the next dose as soon as you remember, unless it is less than 6 hours until your next dose. Do not use a double dose to make up for the forgotten one.

If you have any further questions on the use of this product, ask your doctor or pharmacist.

## 4. POSSIBLE SIDE EFFECTS

Like all medicines, Pentasa can cause side effects, although not everyone gets them. Following rectal administration local reactions such as itching, rectal discomfort and urge may occur.

**STOP using Pentasa and seek medical attention immediately if you notice any of the following symptoms:**

- reddish non-elevated, target-like or circular patches on the trunk, often with central blisters, skin peeling, ulcers of mouth, throat, nose, genitals and eyes, widespread rash, fever and enlarged lymph nodes. These serious skin rashes can be preceded by fever and flu-like symptoms.
- itching, skin rashes, swelling of the face, lips or throat, difficulty in breathing or wheeziness (signs of an allergic reaction).
- unexplained bleeding, bruising, skin rashes, fever or sore throat (signs of blood disorder).
- a change in the colour or amount of urine produced (signs of kidney problems).

If you experience any of the above side effects, you should contact your doctor or go to the nearest hospital emergency department immediately.

**The following common side effects** may affect up to 1 in 10 people:

- diarrhoea
- abdominal pain
- nausea
- vomiting
- headache
- rash
- flatulence (passing wind)
- rectal discomfort,
- irritation at the site of application

**The following rare side effects** may affect up to 1 in 1,000 people:

- inflammation of some areas of the heart (myocarditis and pericarditis) which can cause shortness of breath and chest pain or palpitations (rapid or irregular heart beats).
- inflammation of the pancreas (symptoms include back and/or stomach pain).
- increased amylase (enzyme that helps digest carbohydrates).
- dizziness.
- increased sensitivity of your skin to sun and ultraviolet light (photosensitivity).

**The following very rare side effects** may affect up to 1 in 10,000 people:

- anaemia and other blood disorders (decrease in the numbers of certain blood cells, which can cause unexplained bleeding, bruising, fever or sore throat)
- liver disorders (symptoms include jaundice (yellowing of the skin and/or eyes) and/or pale bowel motions)
- kidney disorders (symptoms include blood in the urine, and/or oedema (swelling due to build-up of fluid))
- peripheral neuropathy (a condition affecting the nerves of the hands and feet symptoms include tingling and numbness)
- allergic and fibrotic lung reactions, inflammation of the lining of the lungs or lung scarring (symptoms include coughing, bronchospasm, chest discomfort or pain on breathing, breathing difficulties, bloody and/or excessive phlegm)
- pancolitis (a kind of inflammatory bowel disorder (IBD) that affects the entire internal lining of the large bowel)
- hair loss (this is reversible)
- muscle or joint pain
- inflammation which can affect different parts of the body such as joints, skin, kidneys, heart etc. (symptoms include painful joints, fatigue, fever, abnormal or unexplained bleeding (e.g. nose bleeds), bruising, purple discoloration of the skin (including severe skin erosions and severe blistering that may affect the skin as the protective barrier of the body))
- semen with a low concentration of sperm (oligospermia) (this is reversible)
- severe diarrhoea and abdominal pain because of an allergic reaction to this medicine within the bowel
- allergic reactions and fever may occasionally occur.

**Not known** (frequency cannot be estimated from the available data)

- kidney stones and associated kidney pain (see also section 2)
- change in urine colour.
- If you experience strong or recurrent headache, disturbed vision, or ringing or buzzing in the ears. These could be symptoms of increased pressure within your skull (idiopathic intracranial hypertension).

**Reporting of side effects**

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet.

You can also report side effects directly via the Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the **Google Play** or **Apple App Store**.

By reporting side effects, you can help provide more information on the safety of this medicine.

**5. HOW TO STORE PENTASA**

Keep out of the sight and reach of children.

Do not store above 25°C.

Store in the original package in order to protect from light.

Do not use the suppositories after the expiry date which is stated on the carton and blister labels after 'Exp'. The date refers to the last day of that month.

If the suppositories become discoloured or show any signs of deterioration, seek the advice of your pharmacist.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

**6. CONTENTS OF THE PACK AND OTHER INFORMATION**

**What Pentasa contains**

The active ingredient in Pentasa is mesalazine.

Each suppository contains 1g mesalazine.

The other ingredients are magnesium stearate, talc, povidone, and macrogol 6,000.

**What Pentasa looks like and the contents of the pack**

Pentasa are oblong, compressed, white to light tan speckled suppositories.

Each pack contains 28 suppositories (4 blister x 7 suppositories), with a cap in aluminum foil and Finger cots.

**Manufactured by:** Ferring GmbH, Wittland 11, D-24109 Kiel, Germany.

**Procured from within the EU and repackaged by the Product Licence holder:** B&S Healthcare, Unit 4, Bradfield Road, Ruislip, Middlesex, HA4 0NU, UK.

**Pentasa® 1g suppositories; PL 18799/2923**

Leaflet date: 28.08.2025

**POM**

Pentasa is the registered trademark of Ferring B.V.

**Blind or partially sighted?  
Is this leaflet hard to see or read?  
Call 0208 515 3763 to obtain the  
leaflet in a format suitable for you.**

## Mesalazine 1g suppositories

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

The name of your medicine is Mesalazine 1g suppositories but will be referred to as Mesalazine throughout this leaflet.

### What is in this leaflet:

1. What Mesalazine is and what it is used for
2. What you need to know before you use Mesalazine
3. How to use Mesalazine
4. Possible side effects
5. How to store Mesalazine
6. Contents of the pack and other information

## 1. WHAT MESALAZINE IS AND WHAT IT IS USED FOR

Each Mesalazine contains mesalazine 1g as the active ingredient. Mesalazine belongs to a group of medicines called salicylates.

Mesalazine is used to treat inflammation in the gut caused by conditions such as ulcerative colitis. Sometimes ulcerative colitis only affects the back passage (rectum), this is known as ulcerative proctitis. Mesalazine are used to treat ulcerative proctitis and are intended for rectal use only.

The suppositories release the active ingredient slowly which then acts locally to reduce the inflammation and help relieve or stop the pain.

## 2. WHAT YOU NEED TO KNOW BEFORE YOU USE MESALAZINE

### Do Not use Mesalazine if you:

- are **allergic (hypersensitive) to mesalazine** or any of the other ingredients of this medicine (see section 6)
- are **allergic to other salicylates e.g. acetylsalicylic acid**
- have **severe liver and/or kidney** problems

### Warnings and precautions

#### Talk to your doctor or pharmacist before using these suppositories if you:

- are allergic to **sulphasalazine** (risk of allergy to salicylates)
- currently have or have previously had **liver** or **kidney** disease
- have a medical condition that can make you prone to **bleeding**
- are on medication that may affect kidney function e.g. Non-steroidal anti-inflammatory drugs (NSAIDs) such as aspirin
- have lung problems, in particular asthma
- suddenly develop abdominal cramps, abdominal pain, fever, severe headache and rash. In such circumstances you should stop taking Mesalazine immediately.
- Kidney stones may develop with use of mesalazine. Symptoms may include pain in sides of abdomen and blood in urine. Take care to drink sufficient amount of liquid during treatment with mesalazine
- Have ever developed a severe skin rash or skin peeling, blistering and/or mouth sores after using mesalazine.

Mesalazine may produce red-brown urine discoloration after contact with sodium hypochlorite bleach in the toilet water. It concerns a chemical reaction between mesalazine and bleach and is harmless.

### Take special care with mesalazine:

Serious skin reactions including Drug reaction with eosinophilia and systemic symptoms (DRESS), Stevens-Johnson syndrome (SJS) and toxic epidermal necrolysis (TEN) have been reported in association with mesalazine treatment. Stop using Mesalazine and seek medical attention immediately if you notice any of the symptoms related to these serious skin reactions described in section 4.

If you experience strong or recurrent headache, disturbed vision, or ringing or buzzing in the ears contact your doctor immediately.

**While you are on treatment with this medicine, your doctor will normally arrange blood and urine tests to check your kidney function especially at the beginning of treatment.**

You should make sure that you don't become dehydrated, while you are using this medicine. This can occur after severe or prolonged attacks of vomiting and/or diarrhoea, high fever or heavy sweating. If this occurs, you should talk to your doctor or a pharmacist for advice as soon as possible.

### Other medicines and Mesalazine

**Tell your doctor or pharmacist if you are taking or have recently taken any other medicines**, including medicines obtained without a prescription. This is especially important if you are taking any of the following:

- **azathioprine** (used after transplantations or to treat auto-immune diseases)
- **6-mercaptopurine or thioguanine** (chemotherapy, used to treat leukaemia)
- certain agents that inhibit blood clotting (medicines for thrombosis or to thin your blood, e.g. warfarin)

### Pregnancy, breast-feeding and fertility

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before using this medicine.

There is limited experience with the use of mesalazine during pregnancy and breast-feeding. Blood disorders have been reported in newborns of mothers being treated with this medicine. The newborn may develop allergic reactions after breast-feeding, e.g. diarrhoea. If the newborn develops diarrhoea, breast-feeding should be discontinued.

### Driving and using machines

This medicine is not known to affect the ability to drive and/or use machines.

## 3. HOW TO USE MESALAZINE

For Adult use only, not recommended in children.

Always use Mesalazine exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure. The suppository should be inserted into the back passage (rectum) only. It is intended to be left in place for as long as possible and is therefore best inserted before going to sleep. To treat an attack of proctitis your doctor will usually prescribe one suppository to be used daily for two to four weeks. To help prevent further attacks it may be used for longer.

### How to insert the suppository

You should make sure that you empty your bowels before inserting the suppository.

- a. Tear along the perforation to remove one suppository blister from a blister strip
- b. Put a protector on the finger with which you will insert the suppository
- c. Remove the suppository from the blister before insertion. This is done by pushing the suppository through the printed foil
- d. Insertion may be made easier if you moisten the suppository with water first
- e. Insert the suppository gently and fully into the back passage. It may help to lie down on one side and raise the other knee.
- f. Remain still for a few minutes after insertion to help keep the suppository in place. If it comes out within 10 minutes, a new one should be inserted
- g. Dispose of the empty blister and used finger protector safely and wash your hands.

### If you use more Mesalazine than you should

If you accidentally use too many suppositories, you should go to your nearest emergency department or contact your doctor immediately. Take the pack and any remaining suppositories with you.

### If you forget to use Mesalazine

If you forget to use a dose, use the next dose as soon as you remember, unless it is less than 6 hours until your next dose. Do not use a double dose to make up for the forgotten one.

If you have any further questions on the use of this product, ask your doctor or pharmacist.

## 4. POSSIBLE SIDE EFFECTS

Like all medicines, Mesalazine can cause side effects, although not everyone gets them. Following rectal administration local reactions such as itching, rectal discomfort and urge may occur.

**STOP using Mesalazine and seek medical attention immediately if you notice any of the following symptoms:**

- reddish non-elevated, target-like or circular patches on the trunk, often with central blisters, skin peeling, ulcers of mouth, throat, nose, genitals and eyes, widespread rash, fever and enlarged lymph nodes. These serious skin rashes can be preceded by fever and flu-like symptoms.
- itching, skin rashes, swelling of the face, lips or throat, difficulty in breathing or wheeziness (signs of an allergic reaction).
- unexplained bleeding, bruising, skin rashes, fever or sore throat (signs of blood disorder).
- a change in the colour or amount of urine produced (signs of kidney problems).

If you experience any of the above side effects, you should contact your doctor or go to the nearest hospital emergency department immediately.

**The following common side effects** may affect up to 1 in 10 people:

- diarrhoea
- abdominal pain
- nausea
- vomiting
- headache
- rash
- flatulence (passing wind)
- rectal discomfort,
- irritation at the site of application

**The following rare side effects** may affect up to 1 in 1,000 people:

- inflammation of some areas of the heart (myocarditis and pericarditis) which can cause shortness of breath and chest pain or palpitations (rapid or irregular heart beats).
- inflammation of the pancreas (symptoms include back and/or stomach pain).
- increased amylase (enzyme that helps digest carbohydrates).
- dizziness.
- increased sensitivity of your skin to sun and ultraviolet light (photosensitivity).

**The following very rare side effects** may affect up to 1 in 10,000 people:

- anaemia and other blood disorders (decrease in the numbers of certain blood cells, which can cause unexplained bleeding, bruising, fever or sore throat)
- liver disorders (symptoms include jaundice (yellowing of the skin and/or eyes) and/or pale bowel motions)
- kidney disorders (symptoms include blood in the urine, and/or oedema (swelling due to build-up of fluid))
- peripheral neuropathy (a condition affecting the nerves of the hands and feet symptoms include tingling and numbness)
- allergic and fibrotic lung reactions, inflammation of the lining of the lungs or lung scarring (symptoms include coughing, bronchospasm, chest discomfort or pain on breathing, breathing difficulties, bloody and/or excessive phlegm)
- pancolitis (a kind of inflammatory bowel disorder (IBD) that affects the entire internal lining of the large bowel)
- hair loss (this is reversible)
- muscle or joint pain
- inflammation which can affect different parts of the body such as joints, skin, kidneys, heart etc. (symptoms include painful joints, fatigue, fever, abnormal or unexplained bleeding (e.g. nose bleeds), bruising, purple discoloration of the skin (including severe skin erosions and severe blistering that may affect the skin as the protective barrier of the body))
- semen with a low concentration of sperm (oligospermia) (this is reversible)
- severe diarrhoea and abdominal pain because of an allergic reaction to this medicine within the bowel
- allergic reactions and fever may occasionally occur.

**Not known** (frequency cannot be estimated from the available data)

- kidney stones and associated kidney pain (see also section 2)
- change in urine colour.
- If you experience strong or recurrent headache, disturbed vision, or ringing or buzzing in the ears. These could be symptoms of increased pressure within your skull (idiopathic intracranial hypertension).

**Reporting of side effects**

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet.

You can also report side effects directly via the Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the **Google Play** or **Apple App Store**.

By reporting side effects, you can help provide more information on the safety of this medicine.

**5. HOW TO STORE MESALAZINE**

Keep out of the sight and reach of children.

Do not store above 25°C.

Store in the original package in order to protect from light.

Do not use the suppositories after the expiry date which is stated on the carton and blister labels after 'Exp'. The date refers to the last day of that month.

If the suppositories become discoloured or show any signs of deterioration, seek the advice of your pharmacist.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

**6. CONTENTS OF THE PACK AND OTHER INFORMATION**

**What Mesalazine contains**

The active ingredient in Mesalazine is mesalazine.

Each suppository contains 1g mesalazine.

The other ingredients are magnesium stearate, talc, povidone, and macrogol 6,000.

**What Mesalazine looks like and the contents of the pack**

Mesalazine are oblong, compressed, white to light tan speckled suppositories.

Each pack contains 28 suppositories (4 blister x 7 suppositories), with a cap in aluminum foil and Finger cots.

**Manufactured by:** Ferring GmbH, Wittland 11, D-24109 Kiel, Germany.

**Procured from within the EU and repackaged by the Product Licence holder:** B&S Healthcare, Unit 4, Bradfield Road, Ruislip, Middlesex, HA4 0NU, UK.

**Mesalazine 1g suppositories; PL 18799/2923**

Leaflet date: 28.08.2025

**POM**

**Blind or partially sighted?  
Is this leaflet hard to see or read?  
Call 0208 515 3763 to obtain the  
leaflet in a format suitable for you.**