

Package Leaflet: Information for the User
Streflam 8.75 mg lozenges
Flurbiprofen 8.75 mg

Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

- Always use this medicine exactly as described in this leaflet or as your doctor or pharmacist has told you.
- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Streflam 8.75 mg lozenges are and what they are used for
2. What you need to know before you take Streflam 8.75 mg lozenges
3. How to take Streflam 8.75 mg lozenges
4. Possible side effects
5. How to store Streflam 8.75 mg lozenges
6. Contents of the pack and other information

1. What Streflam 8.75 mg lozenges are and what they are used for

Streflam 8.75 mg lozenges contain flurbiprofen. Flurbiprofen is a non-steroidal anti-inflammatory drug (NSAID) which has analgesic, antipyretic and anti-inflammatory properties. Streflam 8.75 mg lozenges are used for the short-term relief of symptoms of sore throat such as throat pain, soreness and swelling, and difficulty in swallowing in adults and children over the age of 12 years.

2. What you need to know before you use Streflam 8.75 mg lozenges

Do not take these lozenges if you:

- are allergic (hypersensitive) to Flurbiprofen or to any of the other ingredients of this medicine (see Section 6).
- have ever had asthma, unexpected wheezing or shortness of breath, a runny nose, facial swelling or itchy rash (hives) after taking aspirin or any other NSAID medicine.
- currently have or have ever had two or more episodes of a stomach ulcer, intestinal ulcers, or gastrointestinal bleeding.
- have had gastrointestinal bleeding or perforation, severe colitis (inflammation of the bowel) or bleeding disorders when taking NSAID medicines in the past.
- are taking high dose aspirin or another NSAID medicine (such as celecoxib, ibuprofen, diclofenac sodium, etc.)
- are in the last 3 months of pregnancy.
- have severe heart, severe kidney or severe liver failure.

Warnings and precautions

Talk to your doctor or pharmacist before taking Streflam 8.75 mg lozenges if you:

- have ever had asthma or suffer from allergies.
- have tonsillitis (inflamed tonsils) or think you may have a bacterial throat infection (as you may need antibiotics)
- you have an infection - please see heading "Infections" below

- have heart, kidney or liver problems.
- have had a stroke.
- have a history of bowel disease (ulcerative colitis, Crohn's disease)
- suffer from chronic autoimmune disease such as systemic lupus erythematosus and mixed connective tissue disease.
- are elderly, as you are more likely to experience the side effects listed in this leaflet.
- are in the first 6 months of pregnancy or breastfeeding.
- have been told by your doctor that you have intolerance to some sugars.

Whilst using Streflam 8.75 mg lozenges

- At the first sign of any skin reaction (rash, peeling, blistering) or other sign of an allergic reaction, stop using the lozenge and consult a doctor at once.
- Report any unusual abdominal symptoms (especially bleeding) to your doctor.
- If you do not get better, you get worse, or develop new symptoms, talk to a doctor.
- Medicines such as Streflam 8.75 mg lozenges may be associated with a small increased risk of heart attack (myocardial infarction) or stroke. Any risk is more likely with high doses and prolonged treatment. Do not exceed the recommended dose or duration of treatment (3 days).

Children

This medicine should not be used by children under the age of 12 years.

Other medicines and Streflam 8.75 mg lozenges

Please tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicine. In particular, tell them if you are taking:

- low dose aspirin (up to 75 mg daily)
- medicines for high blood pressure or heart failure (antihypertensives, cardiac glycosides)
- water tablets (diuretics, including potassium sparing drugs)
- medicines for thinning the blood (anticoagulants, antiplatelet agents)
- medicines for gout (probenecid, sulfinpyrazone)
- Other NSAID medicines or corticosteroids (such as celecoxib, ibuprofen, diclofenac sodium or prednisolone)
- mifepristone (a medicine used for pregnancy termination)
- quinolone antibiotics (such as ciprofloxacin)
- cyclosporine or tacrolimus (to suppress the immune system)
- phenytoin (to treat epilepsy)
- methotrexate (to treat autoimmune diseases or cancer)
- lithium or SSRIs (for depression)
- oral antidiabetics (to treat diabetes)
- zidovudine (to treat HIV)

Taking Streflam 8.75 mg lozenges with food, drink and alcohol.

Alcohol should be avoided during treatment with Streflam 8.75 mg lozenges as it increases the risk of bleeding in the stomach or intestines.

Pregnancy, breast-feeding and fertility

Do not take this medicine if you are **in the last 3 months of pregnancy**. If you are in the first 6 months of pregnancy or are breast-feeding, speak to your doctor before taking these lozenges.

Flurbiprofen belongs to a group of medicines which may impair fertility in women. This effect is reversible on stopping the medicine. It is unlikely that the lozenges, used occasionally, will affect your chances of becoming pregnant; however, tell your doctor before taking this medicine if you have problems becoming pregnant.

Infections

Non-steroidal anti-inflammatory drugs (NSAIDs) may hide signs of infections such as fever and pain. This may delay appropriate treatment of infection, which may lead to an increased risk of complications. If you take this medicine while you have an infection and your symptoms of the infection persist or worsen, consult a doctor or pharmacist without delay.

Driving and using machines

No studies on the effects on the ability to drive and use of machines have been performed. However, dizziness and visual disturbances are possible side effects after taking NSAIDs. If affected, do not drive or operate machinery.

Important information about some of the ingredients of Streflam 8.75 mg lozenges

Streflam 8.75 mg lozenges contains 1.069 mg of glucose and 1.407 mg of sucrose per dose. This should be taken into account in patients with diabetes mellitus.

If you have been told by your doctor that you have intolerance to some sugars, contact your doctor before taking this medicinal product.

This medicine contains only very low levels of gluten (from wheat starch). It is regarded as 'glutenfree' and is very unlikely to cause problems if you have coeliac disease.

One lozenge contains no more than 21.38 micrograms of gluten. If you have wheat allergy (different from coeliac disease) you should not take this medicine.

This medicine contains fragrance with Citral, Citronellol, d-Limonene, Farnesol, Geraniol and Linalool. Citral, Citronellol, d-Limonene, Farnesol, Geraniol and Linalool may cause allergic reactions.

Sulphur Dioxide (E220) may rarely cause severe hypersensitivity reactions and bronchospasm. Butylated hydroxyanisole (E320) may cause local skin reactions (e.g. contact dermatitis), or irritation to the eyes and mucous membranes.

3. How to take Streflam 8.75 mg lozenges

Always use this medicine exactly as described in this leaflet or as your doctor or pharmacist has told you.

Check with your doctor or pharmacist if you are not sure.

The recommended dose: Adults and children over the age of 12 years:

- Take one lozenge by mouth and suck slowly.
- Always move the lozenge around whilst sucking.
- The lozenges should start to work within 30 minutes.
- Then take one lozenge every 3-6 hours, if needed.
- **Do not take more than 5 lozenges in 24 hours.**

Do not use this product in children under 12 years.

These lozenges are for short-term use only. The lowest effective dose should be used for the shortest duration necessary to relieve symptoms. If you have an infection, consult a doctor or pharmacist without delay if symptoms (such as fever and pain) persist or worsen (see section 2). If mouth irritation occurs, flurbiprofen treatment should be withdrawn.

Do not take Streflam 8.75 mg lozenges for more than 3 days unless your doctor tells you to. If you do not get better, you get worse, or if you develop new symptoms, talk to a doctor or pharmacist.

If you take more Streflam 8.75 mg lozenges than you should

Talk to a doctor or pharmacist or go to your nearest hospital straight away. Symptoms of overdose may include: feeling sick or being sick, stomach ache or, more rarely, diarrhoea. Ringing in the ears, headache and gastrointestinal bleeding is also possible.

If you have any questions on the use of this product, ask your doctor or pharmacist.

If you forget to take Streflam 8.75 mg lozenges

If you forget to take this product, do not 'double dose' to make up for a forgotten dose.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

STOP TAKING this medicine and contact a doctor immediately if you develop:

- signs of an allergic reaction such as asthma, unexplained wheezing or shortness of breath, itchiness, runny nose, skin rashes, etc.
- swelling of the face, tongue or throat causing difficulty in breathing, racing heart and drop in blood pressure leading to shock (These can happen even on the first use of the medicine).
- severe skin reactions such as peeling, blistering or flaking skin.

Tell your doctor or pharmacist if you notice any of the following effects or any effects not listed:

Common (*may affect up to 1 in 10 people*)

- dizziness, headache
- throat irritation
- mouth ulcers or pain in the mouth
- throat pain
- discomfort or unusual sensation in the mouth (such as warmth, burning, tingling, prickling, etc.)
- nausea and diarrhoea
- prickling and itching sensation in skin

Uncommon (*may affect up to 1 in 100 people*)

- drowsiness
- blistering in the mouth or throat, numbness in the throat
- stomach bloating, abdominal pain, wind, constipation, indigestion, vomiting.
- dry mouth
- burning sensation in the mouth, altered sense of taste
- skin rashes, itchy skin
- fever, pain
- feeling sleepy or difficulty in falling asleep
- worsening of asthma, wheezing, shortness of breath
- reduced sensation in the throat

Rare (*may affect up to 1 in 1000 people*)

- anaphylactic reaction

Frequency not known (cannot be estimated from the available data)

- anaemia, thrombocytopenia (low platelet count in the blood that can give rise to bruising and bleeding)
- swelling (oedema), high blood pressure, heart failure or attack
- severe forms of skin reactions such as bullous reactions, including Stevens-Johnson syndrome and Lyell's syndrome and toxic epidermal necrolysis.
- hepatitis (inflammation of the liver)

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store.

By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Streflam 8.75 mg lozenges

- Keep out of the sight and reach of children
- Do not use this medicine after the expiry date, which is stated on the pack after EXP. The expiry date refers to the last date of that month.
- Store in the original carton.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

6. Contents of the pack and further information

What Streflam 8.75 mg lozenges contains:

The active ingredient (which makes the medicine work) is flurbiprofen 8.75 mg. The other ingredients are:

macrogol 300, potassium hydroxide (E525), lemon flavor (containing butylated hydroxyanisole (E320)), levomenthol, invert sugar (honey), liquid glucose (containing wheat starch and sulphur dioxide (E220)) and liquid sucrose.

What Streflam 8.75 mg lozenges look like and contents of the pack

The lozenges are circular in shape with an embossed brand logo and an opaque pale yellow colour. The pack contains 8, 16, 24 or 36 lozenges. Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer:

Marketing Authorisation Holder:

Reckitt Benckiser Healthcare (UK) Ltd, Slough, SL1 3UH

Manufacturer:

RECKITT BENCKISER HEALTHCARE INTERNATIONAL Ltd,
Nottingham, NG90 2DB
UK

This medicinal product is authorised in the Member States of the EEA under the following names:

United Kingdom Streflam 8.75mg lozenges
Austria Strepfen 8,75 mg - Lutschtabletten
Germany Dobendan Direkt Flurbiprofen 8,75 mg Lutschtabletten
Portugal Strepfen 8,75 mg Mel e Limao Flurbiprofeno Pastilhas
Belgium Strepfen 8,75 mg zuigtablet
Cyprus Strefen 8,75 mg Τροχίσκοι
Denmark Strefen 8,75mg Sugetabletter
Estonia Strepsils Intensive, 8,75 mg, Loseng
Greece Strepfen 8,75 mg Τροχίσκοι
Spain Strepfen 8,75 mg pastillas para chupar sabor miel y limón
Finland Strefen 8,75mg Imeskelytabletti
Iceland Strefen 8,75mg Munnsogstafla
Lithuania Strepsils Intensive, 8,75 mg, Kietoji pastilė
Luxembourg Strepfen 8,75 mg pastille
Netherlands Strepsils Strepfen, zuigtabletten 8,75 mg
Sweden Strefen 8,75mg Sugtablett
Slovenia Strepfen 8,75mg pastile

This leaflet was last revised in

Oct 2023