

(clotrimazole)

## Patient Information Leaflet

Read all of this leaflet carefully because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not give it to anyone else under any circumstances.
- If you have any unusual effects after using this product, tell your doctor.

## In this leaflet:

- 1 What is Canesten 500mg Pessary and what is it used for?
- 2 Before you use Canesten 500mg Pessary
- 3 How to use Canesten 500mg Pessary
- 4 Possible side effects
- 5 How to store Canesten 500mg Pessary
- 6 Further information

### 1 What is Canesten 500mg Pessary and what is it used for?

Canesten 500mg Pessary is a full course of treatment for vaginal thrush (candidiasis). It is inserted into the vagina for treatment at the site of infection.

The active substance in Canesten 500mg Pessary is clotrimazole. Clotrimazole belongs to a group of medicines called azoles and is an antifungal agent which fights the cause of infections such as vaginal thrush.

### 2 Before you use Canesten 500mg Pessary

#### Do not use Canesten 500mg Pessary:

- If you are allergic (hypersensitive) to clotrimazole or any of the other ingredients of Canesten 500mg Pessary (see Section 6. Further Information).
- During your period as it may be less effective.

#### Special precautions:

As with other pessaries, this product may reduce the effectiveness of rubber contraceptives, such as condoms or diaphragms. Consequently, you should use alternative precautions for at least five days after using this product.

Do not use tampons, intravaginal douches, spermicides or other vaginal products while using this product.

Avoid vaginal intercourse while you have thrush and during use of this product because your partner could become infected.

#### Using other medicines:


Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription. Inform your doctor if you are taking tacrolimus or sirolimus (used to reduce the immune response to prevent rejection after an organ transplant).

#### Pregnancy and breastfeeding:

Canesten 500mg Pessary can be used in pregnancy and breastfeeding. If you are pregnant, breastfeeding or trying for a baby, tell your doctor or midwife before using Canesten 500mg Pessary. If you have informed your doctor or midwife, follow his/her instructions carefully. To treat internal thrush, your doctor may recommend that you use the pessary without the help of an applicator.


### 3 How to use Canesten 500mg Pessary

Unless directed otherwise by your doctor, the pessary should be inserted as high as possible into the vagina, preferably before going to sleep at night for convenient and comfortable treatment. Wash your hands before handling the applicator and the foil blister pack and again afterwards when you have used the applicator.




1. Image 1 shows all the components included in the Canesten 500mg Pessary pack.

Remove the applicator from the packaging and pull out the plunger (with the red lock attached) from the applicator.



2. Remove the pessary from the foil blister pack and place into the open end of the applicator with the curved edge of the pessary facing down. Push the plunger and lock into the applicator until you feel a click.



3. Once you have felt the click, remove the lock from the plunger.



4. Carefully insert the applicator as deep as is comfortable into the vagina (this is easiest when lying on your back with your knees bent up) up to the patterned grip zone. Hold the applicator at the patterned grip zone. Carefully push the plunger all the way until it stops to dispense the vaginal pessary.

5. Remove the applicator. Dispose of the applicator in a safe place, out of the reach of children.

The applicator cannot be flushed down the toilet.

As the pessary dissolves in the vagina, it may be helpful to wear a panty liner. It may leave a white chalky residue after using the pessary, which does not mean that the treatment has not worked. However, if you notice pieces of undissolved pessary, speak to your doctor or a pharmacist, as the treatment may not have worked properly.

The symptoms of thrush should disappear within three days of treatment.

If no improvement is seen after seven days you must tell your doctor.

**Canesten 500mg Pessary is for use in the vagina only:**

**Do not put the pessary in your mouth or swallow it.**

If the pessary is swallowed accidentally, tell your doctor straight away or contact the Accident and Emergency Department of your nearest hospital.

#### 4 Possible side effects

Like all medicines, Canesten 500mg Pessary can cause side effects, although not everybody gets them.

As with all medicines, some people may be allergic to the pessary. If you are allergic, a reaction will occur soon after you have used the medicine. If you experience an allergic reaction or the redness, burning, pain, itching or swelling get worse, stop using this product and tell your doctor straight away or contact the Accident and Emergency Department of your nearest hospital. Signs of an allergic reaction may include:

- Rash.
- Swallowing or breathing problems.
- Swelling of your lips, face, throat or tongue.
- Weakness, feeling dizzy or faint.
- Nausea.

After you use the pessary you might experience:

- Itching, rash, swelling, redness, discomfort, burning, irritation, vaginal peeling, discharge or bleeding.
- Pain (eg. abdominal pain, vulvovaginal pain).
- Nausea.

If you experience any of the above effects tell your doctor immediately.

#### Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the **Google Play** or **Apple App Store**. By reporting side effects you can help provide more information on the safety of this medicine.

#### 5 How to store Canesten 500mg Pessary

**Keep this medicine out of the sight and reach of children.**

Do not store above 25°C.

Do not use Canesten 500mg Pessary after the expiry date which is stated on the carton and the foil blister strip after EXP. The expiry date refers to the last day of that month.

If your doctor tells you to stop using this medicine, return any unused Canesten 500mg Pessary to your pharmacist (chemist) for safe disposal. Only keep this medicine if your doctor tells you to.

If the pessary becomes discoloured or shows any other signs of deterioration, you should seek the advice of your pharmacist who will tell you what to do.

Medicines should not be disposed of via wastewater or household waste.

Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

#### 6 Further Information

**What Canesten 500mg Pessary contains:**

- The active substance is clotrimazole at a strength of 500mg.
- The other ingredients are lactose monohydrate, cellulose microcrystalline, crospovidone, hypromellose 15 CPS, lactic acid, calcium lactate pentahydrate, maize starch, magnesium stearate, colloidal anhydrous silica and purified water.

**What Canesten 500mg Pessary looks like and contents of the pack:**

Canesten 500mg Pessary contains a white pessary held inside a foil blister and an applicator for insertion of the pessary into the vagina.

#### Manufacturer and Licence Holder

Manufactured by GP Grenzach Produktions, GmbH, Germany and is procured from within the EU and repackaged by the Product Licence Holder: Lexon (UK) Limited, Unit 18, Oxleasow Road, East Moons Moat, Redditch, Worcestershire, B98 0RE.

Canesten is a registered trademark of Bayer Intellectual Property GmbH.

**POM** PL 15184/2502 Canesten 500mg Pessary

Leaflet revision date: 17/10/2025

**Blind or partially sighted?  
Is this leaflet hard to see  
or read? Phone Lexon  
(UK) Limited, Tel: 01527  
505414 to obtain the  
leaflet in a format suitable  
for you**

**Remember: This medicine has been prescribed for you.**

**Do not give it to anyone else under any circumstances.**

**If you have any doubts about using Canesten 500mg Pessary correctly, seek the advice of your doctor or pharmacist.**

**Further information about vaginal thrush:**

Vaginal thrush (candidiasis) is a common infection that most women suffer from at some time in their lives and is not caused by lack of personal hygiene.

Thrush is caused by a yeast (fungus) called Candida which lives harmlessly in the vagina and other parts of the body, without you even noticing it. However, the natural balance that keeps Candida under control can be upset by many factors such as hormonal changes (menstruation, contraceptive pill, pregnancy, menopause), poor health, antibiotics, perfumed soaps, bath additives and tight clothing.

If the natural pH balance is altered, the level of yeast increases and can develop into a thrush infection causing any of the following symptoms: persistent burning and/or itching around the vagina and vulva, redness, swelling and soreness of the tissues of the vagina and vulva and a whitish, odourless discharge from the vagina. Not everybody who has thrush has all these symptoms; you may have only one of them.

**How to avoid future recurrences:**

- ✓ Wear cotton knickers and loose clothing.
- ✓ Wash daily.
- ✓ After going to the toilet, wipe yourself from the front to back as a thrush infection may be transferred from the bowel.
- ✓ Change your sanitary protection regularly.
- X Try to avoid wearing tights, nylon knickers and close fitting jeans.
- X Try to avoid washing with perfumed soaps or using vaginal deodorants.
- X Do not wash or rub yourself hard with sponges or flannels and avoid hot baths with strong perfumed oils.

If you are still worried or have any questions about the symptoms or the treatment of thrush, do not hesitate to ask your doctor or pharmacist for advice.

For UK residents only: if you have any questions or would like more information, call our Canesten Advice Line on 0845 758 5030. Calls charged at local rate.

(clotrimazole)

## Patient Information Leaflet

Read all of this leaflet carefully because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not give it to anyone else under any circumstances.
- If you have any unusual effects after using this product, tell your doctor.

## In this leaflet:

- 1 What is Canesten 500mg Pessary and what is it used for?
- 2 Before you use Canesten 500mg Pessary
- 3 How to use Canesten 500mg Pessary
- 4 Possible side effects
- 5 How to store Canesten 500mg Pessary
- 6 Further information

### 1 What is Canesten 500mg Pessary and what is it used for?

Canesten 500mg Pessary is a full course of treatment for vaginal thrush (candidiasis). It is inserted into the vagina for treatment at the site of infection.

The active substance in Canesten 500mg Pessary is clotrimazole. Clotrimazole belongs to a group of medicines called azoles and is an antifungal agent which fights the cause of infections such as vaginal thrush.

### 2 Before you use Canesten 500mg Pessary

#### Do not use Canesten 500mg Pessary:

- If you are allergic (hypersensitive) to clotrimazole or any of the other ingredients of Canesten 500mg Pessary (see Section 6. Further Information).
- During your period as it may be less effective.

#### Special precautions:

As with other pessaries, this product may reduce the effectiveness of rubber contraceptives, such as condoms or diaphragms. Consequently, you should use alternative precautions for at least five days after using this product.

Do not use tampons, intravaginal douches, spermicides or other vaginal products while using this product.

Avoid vaginal intercourse while you have thrush and during use of this product because your partner could become infected.

#### Using other medicines:


Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription. Inform your doctor if you are taking tacrolimus or sirolimus (used to reduce the immune response to prevent rejection after an organ transplant).

#### Pregnancy and breastfeeding:

Canesten 500mg Pessary can be used in pregnancy and breastfeeding. If you are pregnant, breastfeeding or trying for a baby, tell your doctor or midwife before using Canesten 500mg Pessary. If you have informed your doctor or midwife, follow his/her instructions carefully. To treat internal thrush, your doctor may recommend that you use the pessary without the help of an applicator.


### 3 How to use Canesten 500mg Pessary

Unless directed otherwise by your doctor, the pessary should be inserted as high as possible into the vagina, preferably before going to sleep at night for convenient and comfortable treatment. Wash your hands before handling the applicator and the foil blister pack and again afterwards when you have used the applicator.




1. Image 1 shows all the components included in the Canesten 500mg Pessary pack.

Remove the applicator from the packaging and pull out the plunger (with the red lock attached) from the applicator.



2. Remove the pessary from the foil blister pack and place into the open end of the applicator with the curved edge of the pessary facing down. Push the plunger and lock into the applicator until you feel a click.



3. Once you have felt the click, remove the lock from the plunger.



4. Carefully insert the applicator as deep as is comfortable into the vagina (this is easiest when lying on your back with your knees bent up) up to the patterned grip zone. Hold the applicator at the patterned grip zone. Carefully push the plunger all the way until it stops to dispense the vaginal pessary.

5. Remove the applicator. Dispose of the applicator in a safe place, out of the reach of children.

The applicator cannot be flushed down the toilet.

As the pessary dissolves in the vagina, it may be helpful to wear a panty liner. It may leave a white chalky residue after using the pessary, which does not mean that the treatment has not worked. However, if you notice pieces of undissolved pessary, speak to your doctor or a pharmacist, as the treatment may not have worked properly.

The symptoms of thrush should disappear within three days of treatment.

If no improvement is seen after seven days you must tell your doctor.

**Canesten 500mg Pessary is for use in the vagina only:**

**Do not put the pessary in your mouth or swallow it.**

If the pessary is swallowed accidentally, tell your doctor straight away or contact the Accident and Emergency Department of your nearest hospital..

#### 4 Possible side effects

Like all medicines, Canesten 500mg Pessary can cause side effects, although not everybody gets them.

As with all medicines, some people may be allergic to the pessary. If you are allergic, a reaction will occur soon after you have used the medicine. If you experience an allergic reaction or the redness, burning, pain, itching or swelling get worse, stop using this product and tell your doctor straight away or contact the Accident and Emergency Department of your nearest hospital. Signs of an allergic reaction may include:

- Rash.
- Swallowing or breathing problems.
- Swelling of your lips, face, throat or tongue.
- Weakness, feeling dizzy or faint.
- Nausea.

After you use the pessary you might experience:

- Itching, rash, swelling, redness, discomfort, burning, irritation, vaginal peeling, discharge or bleeding.
- Pain (eg. abdominal pain, vulvovaginal pain).
- Nausea.

If you experience any of the above effects tell your doctor immediately.

#### Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the **Google Play** or **Apple App Store**. By reporting side effects you can help provide more information on the safety of this medicine.

#### 5 How to store Canesten 500mg Pessary Keep this medicine out of the sight and reach of children.

Do not store above 25°C.

Do not use Canesten 500mg Pessary after the expiry date which is stated on the carton and the foil blister strip after EXP. The expiry date refers to the last day of that month.

If your doctor tells you to stop using this medicine, return any unused Canesten 500mg Pessary to your pharmacist (chemist) for safe disposal. Only keep this medicine if your doctor tells you to.

If the pessary becomes discoloured or shows any other signs of deterioration, you should seek the advice of your pharmacist who will tell you what to do.

Medicines should not be disposed of via wastewater or household waste.

Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

#### 6 Further Information

##### What Canesten 500mg Pessary contains:

- The active substance is clotrimazole at a strength of 500mg.
- The other ingredients are lactose monohydrate, cellulose microcrystalline, crospovidone, hypromellose 15 CPS, lactic acid, calcium lactate pentahydrate, maize starch, magnesium stearate, colloidal anhydrous silica and purified water.

##### What Canesten 500mg Pessary looks like and contents of the pack:

Canesten 500mg Pessary contains a white pessary held inside a foil blister and an applicator for insertion of the pessary into the vagina.

#### Manufacturer and Licence Holder

Manufactured by Bayer AG, Leverkusen, Germany and is procured from within the EU and repackaged by the Product Licence Holder: Lexon (UK) Limited, Unit 18, Oxleasow Road, East Moons Moat, Redditch, Worcestershire, B98 0RE.

Canesten is a registered trademark of Bayer Intellectual Property GmbH.

**POM** PL 15184/2502 Canesten 500mg Pessary

Leaflet revision date: 17/10/2025

## Blind or partially sighted? Is this leaflet hard to see or read? Phone Lexon (UK) Limited, Tel: 01527 505414 to obtain the leaflet in a format suitable for you

**Remember: This medicine has been prescribed for you.**

**Do not give it to anyone else under any circumstances.**

**If you have any doubts about using Canesten 500mg Pessary correctly, seek the advice of your doctor or pharmacist.**

#### Further information about vaginal thrush:

Vaginal thrush (candidiasis) is a common infection that most women suffer from at some time in their lives and is not caused by lack of personal hygiene.

Thrush is caused by a yeast (fungus) called Candida which lives harmlessly in the vagina and other parts of the body, without you even noticing it. However, the natural balance that keeps Candida under control can be upset by many factors such as hormonal changes (menstruation, contraceptive pill, pregnancy, menopause), poor health, antibiotics, perfumed soaps, bath additives and tight clothing.

If the natural pH balance is altered, the level of yeast increases and can develop into a thrush infection causing any of the following symptoms: persistent burning and/or itching around the vagina and vulva, redness, swelling and soreness of the tissues of the vagina and vulva and a whitish, odourless discharge from the vagina. Not everybody who has thrush has all these symptoms; you may have only one of them.

#### How to avoid future recurrences:

- ✓ Wear cotton knickers and loose clothing.
- ✓ Wash daily.
- ✓ After going to the toilet, wipe yourself from the front to back as a thrush infection may be transferred from the bowel.
- ✓ Change your sanitary protection regularly.
- X Try to avoid wearing tights, nylon knickers and close fitting jeans.
- X Try to avoid washing with perfumed soaps or using vaginal deodorants.
- X Do not wash or rub yourself hard with sponges or flannels and avoid hot baths with strong perfumed oils.

If you are still worried or have any questions about the symptoms or the treatment of thrush, do not hesitate to ask your doctor or pharmacist for advice.

For UK residents only: if you have any questions or would like more information, call our Canesten Advice Line on 0845 758 5030. Calls charged at local rate.

## Patient Information Leaflet

### Read all of this leaflet carefully because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not give it to anyone else under any circumstances.
- If you have any unusual effects after using this product, tell your doctor

### In this leaflet:

- 1 What is Clotrimazole 500mg Pessary and what is it used for?
- 2 Before you use Clotrimazole 500mg Pessary
- 3 How to use Clotrimazole 500mg Pessary
- 4 Possible side effects
- 5 How to store Clotrimazole 500mg Pessary
- 6 Further information

### 1 What is Clotrimazole 500mg Pessary and what is it used for?

Clotrimazole 500mg Pessary is a full course of treatment for vaginal thrush (candidiasis). It is inserted into the vagina for treatment at the site of infection.

The active substance in Clotrimazole 500mg Pessary is clotrimazole. Clotrimazole belongs to a group of medicines called azoles and is an antifungal agent which fights the cause of infections such as vaginal thrush.

### 2 Before you use Clotrimazole 500mg Pessary

#### Do not use Clotrimazole 500mg Pessary:

- If you are allergic (hypersensitive) to clotrimazole or any of the other ingredients of Clotrimazole 500mg Pessary (see Section 6. Further Information).
- During your period as it may be less effective.

#### Special precautions:

As with other pessaries, this product may reduce the effectiveness of rubber contraceptives, such as condoms or diaphragms. Consequently, you should use alternative precautions for at least five days after using this product.

Do not use tampons, intravaginal douches, spermicides or other vaginal products while using this product.

Avoid vaginal intercourse while you have thrush and during use of this product because your partner could become infected.

#### Using other medicines:

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription.


Inform your doctor if you are taking tacrolimus or sirolimus (used to reduce the immune response to prevent rejection after an organ transplant).

#### Pregnancy and breastfeeding:

Clotrimazole 500mg Pessary can be used in pregnancy and breastfeeding. If you are pregnant, breastfeeding or trying for a baby, tell your doctor or midwife before using Clotrimazole 500mg Pessary. If you have informed your doctor or midwife, follow his/her instructions carefully. To treat internal thrush, your doctor may recommend that you use the pessary without the help of an applicator.


### 3 How to use Clotrimazole 500mg Pessary

Unless directed otherwise by your doctor, the pessary should be inserted as high as possible into the vagina, preferably before going to sleep at night for convenient and comfortable treatment. Wash your hands before handling the applicator and the foil blister pack and again afterwards when you have used the applicator.




1. Image 1 shows all the components included in the Clotrimazole 500mg Pessary pack.

Remove the applicator from the packaging and pull out the plunger (with the red lock attached) from the applicator.



2. Remove the pessary from the foil blister pack and place into the open end of the applicator with the curved edge of the pessary facing down. Push the plunger and lock into the applicator until you feel a click.



3. Once you have felt the click, remove the lock from the plunger.



4. Carefully insert the applicator as deep as is comfortable into the vagina (this is easiest when lying on your back with your knees bent up) up to the patterned grip zone. Hold the applicator at the patterned grip zone. Carefully push the plunger all the way until it stops to dispense the vaginal pessary.

5. Remove the applicator. Dispose of the applicator in a safe place, out of the reach of children.

The applicator cannot be flushed down the toilet.

As the pessary dissolves in the vagina, it may be helpful to wear a pantyliner. It may leave a white chalky residue after using the pessary, which does not mean that the treatment has not worked. However, if you notice pieces of undissolved pessary, speak to your doctor or a pharmacist, as the treatment may not have worked properly.

The symptoms of thrush should disappear within three days of treatment.

If no improvement is seen after seven days you must tell your doctor.

**Clotrimazole 500mg Pessary is for use in the vagina only:**

**Do not put the pessary in your mouth or swallow it.**

If the pessary is swallowed accidentally, tell your doctor straight away or contact the Accident and Emergency Department of your nearest hospital.

#### **4 Possible side effects**

Like all medicines, Clotrimazole 500mg Pessary can cause side effects, although not everybody gets them.

As with all medicines, some people may be allergic to the pessary. If you are allergic, a reaction will occur soon after you have used the medicine. If you experience an allergic reaction or the redness, burning, pain, itching or swelling get worse, stop using this product and tell your doctor straight away or contact the Accident and Emergency Department of your nearest hospital. Signs of an allergic reaction may include:

- Rash.
- Swallowing or breathing problems.
- Swelling of your lips, face, throat or tongue.
- Weakness, feeling dizzy or faint.
- Nausea.

After you use the pessary you might experience:

- Itching, rash, swelling, redness, discomfort, burning, irritation, vaginal peeling, discharge or bleeding.
- Pain (eg. abdominal pain, vulvovaginal pain).
- Nausea.

If you experience any of the above effects tell your doctor immediately.

#### **Reporting of side effects**

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the **Google Play** or **Apple App Store**. By reporting side effects you can help provide more information on the safety of this medicine.

#### **5 How to store Clotrimazole 500mg Pessary**

**Keep this medicine out of the sight and reach of children.**

Do not store above 25°C.

Do not use Clotrimazole 500mg Pessary after the expiry date which is stated on the carton and foil blister strip after EXP. The expiry date refers to the last day of that month.

If your doctor tells you to stop using this medicine, return any unused Clotrimazole 500mg Pessary to your pharmacist (chemist) for safe disposal. Only keep this medicine if your doctor tells you to.

If the pessary becomes discoloured or shows any other signs of deterioration, you should seek the advice of your pharmacist who will tell you what to do.

Medicines should not be disposed of via wastewater or household waste.

Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

#### **6 Further Information**

**What Clotrimazole 500mg Pessary contains:**

- The active substance is clotrimazole at a strength of 500mg.
- The other ingredients are lactose monohydrate, cellulose microcrystalline, crospovidone, hypromellose 15 CPS, lactic acid, calcium lactate pentahydrate, maize starch, magnesium stearate, colloidal anhydrous silica and purified water.

**What Clotrimazole 500mg Pessary looks like and contents of the pack:**

Clotrimazole 500mg Pessary contains a white pessary held inside a foil blister and an applicator for insertion of the pessary into the vagina.

#### **Manufacturer and Licence Holder**

Manufactured by GP Grenzach Produktions, GmbH, Germany and is procured from within the EU and repackaged by the Product Licence Holder: Lexon (UK) Limited, Unit 18, Oxleasow Road, East Moons Moat, Redditch, Worcestershire, B98 0RE.

**POM** PL 15184/2502 Clotrimazole 500mg Pessary

Leaflet revision date: 17/10/2025

**Blind or partially sighted?  
Is this leaflet hard to see  
or read? Phone Lexon  
(UK) Limited, Tel: 01527  
505414 to obtain the  
leaflet in a format suitable  
for you**

**Remember: This medicine has been prescribed for you.**

**Do not give it to anyone else under any circumstances.**

**If you have any doubts about using Clotrimazole 500mg Pessary correctly, seek the advice of your doctor or pharmacist.**

#### **Further information about vaginal thrush:**

Vaginal thrush (candidiasis) is a common infection that most women suffer from at some time in their lives and is not caused by lack of personal hygiene.

Thrush is caused by a yeast (fungus) called Candida which lives harmlessly in the vagina and other parts of the body, without you even noticing it. However, the natural balance that keeps Candida under control can be upset by many factors such as hormonal changes (menstruation, contraceptive pill, pregnancy, menopause), poor health, antibiotics, perfumed soaps, bath additives and tight clothing.

If the natural pH balance is altered, the level of yeast increases and can develop into a thrush infection causing any of the following symptoms: persistent burning and/or itching around the vagina and vulva, redness, swelling and soreness of the tissues of the vagina and vulva and a whitish, odourless discharge from the vagina. Not everybody who has thrush has all these symptoms; you may have only one of them.

#### **How to avoid future recurrences:**

- ✓ Wear cotton knickers and loose clothing.
- ✓ Wash daily.
- ✓ After going to the toilet, wipe yourself from the front to back as a thrush infection may be transferred from the bowel.
- ✓ Change your sanitary protection regularly.
- X Try to avoid wearing tights, nylon knickers and close fitting jeans.
- X Try to avoid washing with perfumed soaps or using vaginal deodorants.
- X Do not wash or rub yourself hard with sponges or flannels and avoid hot baths with strong perfumed oils.

If you are still worried or have any questions about the symptoms or the treatment of thrush, do not hesitate to ask your doctor or pharmacist for advice.

For UK residents only: if you have any questions or would like more information, call our Canesten Advice Line on 0845 758 5030. Calls charged at local rate.

## Patient Information Leaflet

### Read all of this leaflet carefully because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not give it to anyone else under any circumstances.
- If you have any unusual effects after using this product, tell your doctor

### In this leaflet:

- 1 What is Clotrimazole 500mg Pessary and what is it used for?
- 2 Before you use Clotrimazole 500mg Pessary
- 3 How to use Clotrimazole 500mg Pessary
- 4 Possible side effects
- 5 How to store Clotrimazole 500mg Pessary
- 6 Further information

### 1 What is Clotrimazole 500mg Pessary and what is it used for?

Clotrimazole 500mg Pessary is a full course of treatment for vaginal thrush (candidiasis). It is inserted into the vagina for treatment at the site of infection.

The active substance in Clotrimazole 500mg Pessary is clotrimazole. Clotrimazole belongs to a group of medicines called azoles and is an antifungal agent which fights the cause of infections such as vaginal thrush.

### 2 Before you use Clotrimazole 500mg Pessary

#### Do not use Clotrimazole 500mg Pessary:

- If you are allergic (hypersensitive) to clotrimazole or any of the other ingredients of Clotrimazole 500mg Pessary (see Section 6. Further Information).
- During your period as it may be less effective.

#### Special precautions:

As with other pessaries, this product may reduce the effectiveness of rubber contraceptives, such as condoms or diaphragms. Consequently, you should use alternative precautions for at least five days after using this product.

Do not use tampons, intravaginal douches, spermicides or other vaginal products while using this product.

Avoid vaginal intercourse while you have thrush and during use of this product because your partner could become infected.

#### Using other medicines:

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription.


Inform your doctor if you are taking tacrolimus or sirolimus (used to reduce the immune response to prevent rejection after an organ transplant).

#### Pregnancy and breastfeeding:

Clotrimazole 500mg Pessary can be used in pregnancy and breastfeeding. If you are pregnant, breastfeeding or trying for a baby, tell your doctor or midwife before using Clotrimazole 500mg Pessary. If you have informed your doctor or midwife, follow his/her instructions carefully. To treat internal thrush, your doctor may recommend that you use the pessary without the help of an applicator.


### 3 How to use Clotrimazole 500mg Pessary

Unless directed otherwise by your doctor, the pessary should be inserted as high as possible into the vagina, preferably before going to sleep at night for convenient and comfortable treatment. Wash your hands before handling the applicator and the foil blister pack and again afterwards when you have used the applicator.




1. Image 1 shows all the components included in the Clotrimazole 500mg Pessary pack.

Remove the applicator from the packaging and pull out the plunger (with the red lock attached) from the applicator.



2. Remove the pessary from the foil blister pack and place into the open end of the applicator with the curved edge of the pessary facing down. Push the plunger and lock into the applicator until you feel a click.



3. Once you have felt the click, remove the lock from the plunger.



4. Carefully insert the applicator as deep as is comfortable into the vagina (this is easiest when lying on your back with your knees bent up) up to the patterned grip zone. Hold the applicator at the patterned grip zone. Carefully push the plunger all the way until it stops to dispense the vaginal pessary.

5. Remove the applicator. Dispose of the applicator in a safe place, out of the reach of children.

The applicator cannot be flushed down the toilet.

As the pessary dissolves in the vagina, it may be helpful to wear a pantyliner. It may leave a white chalky residue after using the pessary, which does not mean that the treatment has not worked. However, if you notice pieces of undissolved pessary, speak to your doctor or a pharmacist, as the treatment may not have worked properly.

The symptoms of thrush should disappear within three days of treatment.

If no improvement is seen after seven days you must tell your doctor.

**Clotrimazole 500mg Pessary is for use in the vagina only:**

**Do not put the pessary in your mouth or swallow it.**

If the pessary is swallowed accidentally, tell your doctor straight away or contact the Accident and Emergency Department of your nearest hospital.

#### 4 Possible side effects

Like all medicines, Clotrimazole 500mg Pessary can cause side effects, although not everybody gets them.

As with all medicines, some people may be allergic to the pessary. If you are allergic, a reaction will occur soon after you have used the medicine. If you experience an allergic reaction or the redness, burning, pain, itching or swelling get worse, stop using this product and tell your doctor straight away or contact the Accident and Emergency Department of your nearest hospital. Signs of an allergic reaction may include:

- Rash.
- Swallowing or breathing problems.
- Swelling of your lips, face, throat or tongue.
- Weakness, feeling dizzy or faint.
- Nausea.

After you use the pessary you might experience:

- Itching, rash, swelling, redness, discomfort, burning, irritation, vaginal peeling, discharge or bleeding.
- Pain (eg. abdominal pain, vulvovaginal pain).
- Nausea.

If you experience any of the above effects tell your doctor immediately.

#### Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the **Google Play** or **Apple App Store**. By reporting side effects you can help provide more information on the safety of this medicine.

#### 5 How to store Clotrimazole 500mg Pessary

**Keep this medicine out of the sight and reach of children.**

Do not store above 25°C.

Do not use Clotrimazole 500mg Pessary after the expiry date which is stated on the carton and foil blister strip after EXP. The expiry date refers to the last day of that month.

If your doctor tells you to stop using this medicine, return any unused Clotrimazole 500mg Pessary to your pharmacist (chemist) for safe disposal. Only keep this medicine if your doctor tells you to.

If the pessary becomes discoloured or shows any other signs of deterioration, you should seek the advice of your pharmacist who will tell you what to do.

Medicines should not be disposed of via wastewater or household waste.

Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

#### 6 Further Information

**What Clotrimazole 500mg Pessary contains:**

- The active substance is clotrimazole at a strength of 500mg.
- The other ingredients are lactose monohydrate, cellulose microcrystalline, crospovidone, hypromellose 15 CPS, lactic acid, calcium lactate pentahydrate, maize starch, magnesium stearate, colloidal anhydrous silica and purified water.

**What Clotrimazole 500mg Pessary looks like and contents of the pack:**

Clotrimazole 500mg Pessary contains a white pessary held inside a foil blister and an applicator for insertion of the pessary into the vagina.

#### Manufacturer and Licence Holder

Manufactured by Bayer AG, Leverkusen, Germany, Germany and is procured from within the EU and repackaged by the Product Licence Holder: Lexon (UK) Limited, Unit 18, Oxleasow Road, East Moons Moat, Redditch, Worcestershire, B98 0RE.

**POM** PL 15184/2502 Clotrimazole 500mg Pessary

Leaflet revision date: 17/10/2025

**Blind or partially sighted?  
Is this leaflet hard to see  
or read? Phone Lexon  
(UK) Limited, Tel: 01527  
505414 to obtain the  
leaflet in a format suitable  
for you**

**Remember: This medicine has been prescribed for you.**

**Do not give it to anyone else under any circumstances.**

**If you have any doubts about using Clotrimazole 500mg Pessary correctly, seek the advice of your doctor or pharmacist.**

#### Further information about vaginal thrush:

Vaginal thrush (candidiasis) is a common infection that most women suffer from at some time in their lives and is not caused by lack of personal hygiene.

Thrush is caused by a yeast (fungus) called Candida which lives harmlessly in the vagina and other parts of the body, without you even noticing it. However, the natural balance that keeps Candida under control can be upset by many factors such as hormonal changes (menstruation, contraceptive pill, pregnancy, menopause), poor health, antibiotics, perfumed soaps, bath additives and tight clothing.

If the natural pH balance is altered, the level of yeast increases and can develop into a thrush infection causing any of the following symptoms: persistent burning and/or itching around the vagina and vulva, redness, swelling and soreness of the tissues of the vagina and vulva and a whitish, odourless discharge from the vagina. Not everybody who has thrush has all these symptoms; you may have only one of them.

#### How to avoid future recurrences:

- ✓ Wear cotton knickers and loose clothing.
- ✓ Wash daily.
- ✓ After going to the toilet, wipe yourself from the front to back as a thrush infection may be transferred from the bowel.
- ✓ Change your sanitary protection regularly.
- X Try to avoid wearing tights, nylon knickers and close fitting jeans.
- X Try to avoid washing with perfumed soaps or using vaginal deodorants.
- X Do not wash or rub yourself hard with sponges or flannels and avoid hot baths with strong perfumed oils.

If you are still worried or have any questions about the symptoms or the treatment of thrush, do not hesitate to ask your doctor or pharmacist for advice.

For UK residents only: if you have any questions or would like more information, call our Canesten Advice Line on 0845 758 5030. Calls charged at local rate.