

PACKAGE LEAFLET: INFORMATION FOR THE USER
IMIGRAN® 10 mg NASAL SPRAY
(sumatriptan)

This medicine is available as the above name but will be referred to as Imigran throughout this leaflet. Please note that the leaflet also contains information about other strength IMIGRAN® 20 mg NASAL SPRAY.

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

Keep this leaflet. You may need to read it again.

If you have any further questions, ask your doctor or pharmacist.

This medicine has been prescribed for you only. Do not pass it on to others - it may harm them, even if their signs of illness are the same as yours.

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Imigran is and what it is used for
2. What you need to know before you use Imigran
3. How to use Imigran
4. Possible side effects
5. How to store Imigran
6. Contents of the pack and other information

1. What Imigran is and what it is used for

Each Imigran contains a single dose of sumatriptan, which belongs to a group of medicines called triptans (*also known as 5-HT₁ receptor agonists*).

Imigran is used to treat migraine headache.

Migraine symptoms may be caused by the temporary widening of blood vessels in the head. Imigran is believed to reduce the widening of these blood vessels. This in turn helps to take away the headache and relieve other symptoms of a migraine attack, such as feeling or being sick (nausea or vomiting) and sensitivity to light and sound.

2. What you need to know before you use Imigran

Don't use Imigran:

- **If you're allergic** to sumatriptan, or any of the other ingredients of this medicine (listed in section 6).
- **If you have a heart problem** such as narrowing of the arteries (*Ischaemic heart disease*) or chest pains (*angina*) or have already had a heart attack.
- **If you have circulation problems in your legs** that cause cramp-like pains when you walk (*peripheral vascular disease*).
- **If you have had a stroke** or a mini-stroke (*also called a transient ischaemic attack or TIA*).
- **If you have high blood pressure.** You may be able to use Imigran if your high blood pressure is mild and is being treated.
- **If you have serious liver disease.**
- **With other migraine medicines**, including those which contain ergotamine, or similar medicines such as methysergide or any triptan/5-HT₁ receptor agonist (medicines which are also used for treating migraine).
- **With anti-depressants called MAOIs** (*monoamine oxidase inhibitors*), or if you have taken these medicines in the last 2 weeks.

If any of these apply to you:

→ **Tell your doctor, and don't use Imigran.**

Take special care with Imigran

Talk to your doctor or pharmacist before using Imigran.

If you have any extra risk factors

- **If you are a heavy smoker**, or are using **nicotine replacement therapy**, and especially
- **If you are a man aged over 40**, or
- **If you are a woman who has been through the menopause.**

In very rare cases, people have developed serious heart conditions after using Imigran, even though they had no signs of heart disease before. If any of the points above applies to you it could mean you have a greater risk of developing heart disease - so:

→ **Tell your doctor so that your heart function can be checked** before Imigran is prescribed for you.

If you have a history of fits (*seizures*)

Or if you have other conditions which might make it more likely that you'll have a fit - for example, a head injury or alcoholism.

→ **Tell your doctor so that you can be supervised more closely.**

If you have liver or kidney disease

→ **Tell your doctor so that you can be supervised more closely.**

If you are allergic to antibiotics called sulphonamides

If so, you may also be allergic to Imigran. If you know you are allergic to an antibiotic but you are not sure whether it is a sulphonamide:

→ **Tell your doctor or pharmacist before using Imigran.**

If you are taking anti-depressants called SSRIs

(*Selective Serotonin Reuptake Inhibitors*) or SNRIs (*Serotonin Noradrenaline Reuptake Inhibitors*)

→ **Tell your doctor or pharmacist before using Imigran.**

Also see *Other medicines and Imigran*, below.

If you use Imigran frequently

Using Imigran too often may make your headaches worse.

→ **Tell your doctor if this applies to you.** He or she may recommend you stop using Imigran.

If you feel pain or tightness in your chest after you use Imigran

These effects may be intense but they usually pass quickly. If they don't pass quickly, or they become severe:

→ **Get medical help immediately.** Section 4 of this leaflet has more information about these possible side effects.

Other medicines and Imigran

Tell your doctor or pharmacist if you're taking, have recently taken or might take any other medicines.

This includes any herbal products or medicines you've bought without a prescription.

Some medicines must not be taken with Imigran and others may cause adverse effects if they're taken with Imigran. **You must tell your doctor if you are taking:**

- **ergotamine** also used to treat **migraine**, or similar medicines such as methysergide (see Section 2). Don't use Imigran at the same time as these medicines. Stop taking these medicines at least 24 hours before using Imigran. Don't take any medicines which contain ergotamine or compounds similar to ergotamine again for at least 6 hours after using Imigran.
- **other triptans/5-HT₁ receptor agonists** (such as naratriptan, rizatriptan, zolmitriptan), also used to treat **migraine**, (see Section 2). Don't use Imigran at the same time as these medicines. Stop taking these medicines at least 24 hours before using Imigran. Don't take another triptan/5-HT₁ receptor agonist again for at least 24 hours after using Imigran.
- **SSRIs** (*Selective Serotonin Reuptake Inhibitors*) or **SNRIs** (*Serotonin Noradrenaline Reuptake Inhibitors*) used to treat **depression**. Using Imigran with these medicines can cause serotonin syndrome (a collection of symptoms which can include restlessness, confusion, sweating, hallucinations, increased reflexes, muscle spasms, shivering, increased heartbeat and shaking). Tell your doctor immediately if you are affected in this way.
- **MAOIs** (*monoamine oxidase inhibitors*) used to treat **depression**. Don't use Imigran if you have taken these in the last 2 weeks.
- **St John's Wort** (*Hypericum perforatum*). Taking herbal remedies that contain St John's Wort while you are using Imigran may make side effects more likely.

Pregnancy and breast-feeding

- **If you are pregnant, think you may be pregnant or are planning to have a baby**, ask your doctor for advice before taking this medicine. There is only limited information about the safety of Imigran for pregnant women, though up until now there is no evidence of any increased risk of birth defects. Your doctor will discuss with you whether or not you should use Imigran while you are pregnant.
- **Don't breast-feed your baby for 12 hours after using Imigran.** If you express any breast milk during this time, discard the milk and don't give it to your baby. Some breastfeeding women report breast and/or nipple pain after use of sumatriptan. The pain is usually temporary and disappears in 3 to 12 hours.

Driving and using machines

Either the symptoms of migraine or your medicine may make you drowsy. **If you are affected, don't drive or operate machinery.**

3. How to use Imigran

Only use Imigran after your migraine headache begins

Don't use Imigran to try to prevent an attack.

Always use this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

There's a step-by-step guide to using the spray at the end of this section.

How much to use

Adults aged 18 to 65

- **The usual dose for adults aged 18 to 65** is one Imigran 20 mg Nasal Spray into just one nostril. But one Imigran 10 mg Nasal Spray into one nostril is enough for some people. Don't use more than two sprays in 24 hours.

Adolescents aged 12 to 17

- **The usual dose for adolescents aged 12 to 17** is one Imigran 10 mg Nasal Spray into one nostril.

Children under 12

- **Imigran is not recommended** for children under 12 years old.

Older people (aged over 65)

- **Imigran is not recommended** for people aged over 65.

When to use Imigran

- **It's best to use Imigran as soon as you feel a migraine coming on**, although it can be used at any time during an attack.

If your symptoms start to come back

- You can use a second Imigran after 2 hours, but **don't use more than two sprays in 24 hours.**

If the first spray has no effect

- **Don't use a second spray**, or any other Imigran preparation for the same attack. If Imigran doesn't give you any relief:
→ **Ask your doctor or pharmacist for advice.**

If you use more Imigran than you should

- **Don't use more than two Imigran nasal sprays in 24 hours.** Using too much Imigran could make you ill. If you have used more than two sprays in 24 hours:
→ **Contact your doctor for advice.**

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

Step-by-step guide to using your Imigran


Don't open a blister until you are ready to use a spray.

Each spray is sealed in a blister to keep it clean and safe. If you carry a spray without a blister, or in an open blister, it may not work properly when you need it.


Each spray contains just one dose of Imigran.

Don't press the plunger too soon or you will lose the dose.


Imigran has three parts:



How to use the nasal spray



1. **Remove the nasal spray** from the blister packaging just before you want to use it.
2. Get into a comfortable position. You may prefer to sit down.
3. **Blow your nose** if it feels blocked, or if you have a cold.
4. **Hold the nasal spray** gently with your fingers and thumb, as shown in picture **A**. **Don't press the blue plunger yet.**



5. **Block one nostril** by pressing a finger firmly on the side of your nose. It doesn't matter which nostril you choose.
6. **Put the nozzle of the nasal spray into the other nostril**, as far as feels comfortable - about 1 cm or 1/2 inch (picture **B**). **Breathe out gently through your mouth.** Hold your head upright and close your mouth.



7. **Start to breathe in gently through your nose.** As you breathe in: **Press the blue plunger firmly with your thumb.** The plunger may feel a bit stiff and you may hear it click. Keep breathing in while spraying (picture **C**).
8. **Remove the spray and remove your finger from the side of your nose.** **Keep your head upright for 10-20 seconds**, breathing gently in through your nose and out through your mouth. This helps the medicine stay in your nose. Your nose may feel wet inside and you may notice a slight taste after using the spray - this is normal and will soon pass.
9. **Your nasal spray is now empty.** Throw it away safely and hygienically.

4. Possible side effects

Like all medicines, this medicine can cause side effects, but not everybody gets them. Some symptoms may be caused by the migraine itself. Adverse events reported in adults have also been observed in adolescents. These include very rare reports of heart attacks.

Allergic reaction: get doctor's help straight away

The following side effects have occurred but their frequency is not known.

- **The signs of allergy include rash, hives (itchy rash); wheezing; swollen eyelids, face or lips; complete collapse.**

If you get any of these symptoms soon after using Imigran:

→ **Don't use any more. Contact a doctor straight away.**

Very common side effects

(affect more than **1 in 10** people)

- **Unpleasant taste.**

Common side effects

(affect up to **1 in 10** people)

- **Heaviness, pressure, tightness or pain in the chest**, throat or other parts of the body, or feelings of unusual sensations, including numbness, tingling and warmth or cold. These effects may be intense but generally pass quickly.

If these effects continue or become severe (especially the chest pain):

→ **Get medical help urgently.** In a very small number of people these symptoms can be caused by a heart attack.

Other common side effects include:

- **Irritation** or burning sensation in the nose or throat; nose bleeds.
- **Feeling sick** (nausea) or being sick (vomiting), although this may be due to the migraine itself.
- **Tiredness** or drowsiness.
- **Dizziness**, feeling weak, or getting hot flushes.
- **Temporary increase in blood pressure.**
- **Shortness of breath.**
- **Aching muscles.**

Rare side effects

(affect up to **1 in 1,000** people)

- Breast pain.

Very rare side effects

(affect up to **1 in 10,000** people)

- Liver function changes. If you have a blood test to check your liver function, tell your doctor or nurse that you are using Imigran.

Some patients may have the following side effects but it is not known how often they occur

- Seizures/fits, tremors, short-lived/temporary muscle spasm (which may affect jaw movement), neck stiffness.
- Visual disturbances such as flickering, reduced vision, double vision, loss of vision and in some cases even permanent defects (although these may be due to the migraine attack itself).
- Heart problems, where your heart beat may go faster, slower or change rhythm, chest pains (*angina*) or heart attack.
- Pale blue-tinged skin and/or pain in your fingers, toes, ears, nose or jaw in response to cold or stress (*Raynaud's phenomenon*).
- Feeling faint (blood pressure may go down).
- Pain in the lower left side of the stomach and bloody diarrhoea (*ischaemic colitis*).
- Diarrhoea.
- If you had a recent injury or if you have inflammation (like rheumatism or inflammation of the colon) you may experience pain or pain worsening at the site of injury or inflammation.
- Pain in the joints.
- Feeling anxious.
- Difficulty swallowing.
- Excessive sweating.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at Website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects, you can help provide more information on the safety of this medicine.

5. How to store Imigran

- **Keep this medicine out of the sight and reach of children.**
- Do not store above 30°C. Do not freeze.
- Store in the original container in order to protect from light.
- Don't use this medicine after the expiry date shown on the carton. The expiry date refers to the last day of that month.
- Don't throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.
- If your nasal spray fails to work properly or shows any signs of deterioration, you should seek the advice of your pharmacist.

6. Contents of the pack and other information

What Imigran contains

The active substance is sumatriptan.

Each pre-filled nasal spray device contains 10 mg sumatriptan in 0.1 ml of aqueous solution.

The other ingredients are potassium dihydrogen phosphate, dibasic sodium phosphate anhydrous, sulphuric acid, sodium hydroxide and purified water.

What Imigran looks like and contents of the pack

This medicine is provided in a pre-filled nasal spray device and contains sumatriptan in 0.1 ml of a yellow-coloured solution.

Imigran is packed in a box containing the nasal sprays individually sealed in blisters. They are available in packs of 2 and 6 pre-filled nasal spray devices. Each nasal spray contains one dose of Imigran.

PL: 15814/2012

POM

Manufactured by GlaxoSmithKline Manufacturing S.p.A., Strada Provinciale Asolana, 90, 43056, San Polo di Torrile, Parma, Italy.

Procured from within the EU and repackaged by the Product Licence holder: O.P.D. Laboratories Ltd., Unit 6 Colonial Way, Watford, Herts WD24 4PR.

Leaflet revision and issue date (Ref.) 02.12.2025.

Imigran is a registered trademark of the GSK group of companies.

To request a copy of this leaflet in Braille, large print or audio please call 01923 332 796.

PACKAGE LEAFLET: INFORMATION FOR THE USER
SUMATRIPTAN 10 mg NASAL SPRAY

This medicine is available as the above name but will be referred to as Sumatriptan Nasal Spray throughout this leaflet. Please note that the leaflet also contains information about other strength SUMATRIPTAN 20 mg NASAL SPRAY.

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

Keep this leaflet. You may need to read it again.

If you have any further questions, ask your doctor or pharmacist.

This medicine has been prescribed for you only. Do not pass it on to others - it may harm them, even if their signs of illness are the same as yours.

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

- 1. What Sumatriptan Nasal Spray is and what it is used for**
- 2. What you need to know before you use Sumatriptan Nasal Spray**
- 3. How to use Sumatriptan Nasal Spray**
- 4. Possible side effects**
- 5. How to store Sumatriptan Nasal Spray**
- 6. Contents of the pack and other information**

1. What Sumatriptan Nasal Spray is and what it is used for

Each Sumatriptan Nasal Spray contains a single dose of sumatriptan, which belongs to a group of medicines called triptans (*also known as 5-HT₁ receptor agonists*).

Sumatriptan Nasal Spray is used to treat migraine headache.

Migraine symptoms may be caused by the temporary widening of blood vessels in the head. Sumatriptan Nasal Spray is believed to reduce the widening of these blood vessels. This in turn helps to take away the headache and relieve other symptoms of a migraine attack, such as feeling or being sick (nausea or vomiting) and sensitivity to light and sound.

2. What you need to know before you use Sumatriptan Nasal Spray

Don't use Sumatriptan Nasal Spray:

- **If you're allergic** to sumatriptan, or any of the other ingredients of this medicine (listed in section 6).
- **If you have a heart problem** such as narrowing of the arteries (*Ischaemic heart disease*) or chest pains (*angina*) or have already had a heart attack.
- **If you have circulation problems in your legs** that cause cramp-like pains when you walk (*peripheral vascular disease*).
- **If you have had a stroke** or a mini-stroke (*also called a transient ischaemic attack or TIA*).
- **If you have high blood pressure.** You may be able to use Sumatriptan Nasal Spray if your high blood pressure is mild and is being treated.
- **If you have serious liver disease.**
- **With other migraine medicines**, including those which contain ergotamine, or similar medicines such as methysergide or any triptan/5-HT₁ receptor agonist (medicines which are also used for treating migraine).
- **With anti-depressants called MAOIs** (*monoamine oxidase inhibitors*), or if you have taken these medicines in the last 2 weeks.

If any of these apply to you:

→ **Tell your doctor, and don't use Sumatriptan Nasal Spray.**

Take special care with Sumatriptan Nasal Spray

Talk to your doctor or pharmacist before using Sumatriptan Nasal Spray.

If you have any extra risk factors

- **If you are a heavy smoker**, or are using **nicotine replacement therapy**, and especially
- **If you are a man aged over 40**, or
- **If you are a woman who has been through the menopause.**

In very rare cases, people have developed serious heart conditions after using Sumatriptan Nasal Spray, even though they had no signs of heart disease before. If any of the points above applies to you it could mean you have a greater risk of developing heart disease - so:

→ **Tell your doctor so that your heart function can be checked** before Sumatriptan Nasal Spray is prescribed for you.

If you have a history of fits (seizures)

Or if you have other conditions which might make it more likely that you'll have a fit - for example, a head injury or alcoholism.

→ **Tell your doctor so that you can be supervised more closely.**

If you have liver or kidney disease

→ **Tell your doctor so that you can be supervised more closely.**

If you are allergic to antibiotics called sulphonamides

If so, you may also be allergic to Sumatriptan Nasal Spray. If you know you are allergic to an antibiotic but you are not sure whether it is a sulphonamide:

→ **Tell your doctor or pharmacist before using Sumatriptan Nasal Spray.**

If you are taking anti-depressants called SSRIs

(*Selective Serotonin Reuptake Inhibitors*) or SNRIs (*Serotonin Noradrenaline Reuptake Inhibitors*)

→ **Tell your doctor or pharmacist before using Sumatriptan Nasal Spray.**

Also see *Other medicines and Sumatriptan Nasal Spray*, below.

If you use Sumatriptan Nasal Spray frequently

Using Sumatriptan Nasal Spray too often may make your headaches worse.

→ **Tell your doctor if this applies to you.** He or she may recommend you stop using Sumatriptan Nasal Spray.

If you feel pain or tightness in your chest after you use Sumatriptan Nasal Spray

These effects may be intense but they usually pass quickly. If they don't pass quickly, or they become severe:

→ **Get medical help immediately.** Section 4 of this leaflet has more information about these possible side effects.

Other medicines and Sumatriptan Nasal Spray

Tell your doctor or pharmacist if you're taking, have recently taken or might take any other medicines.

This includes any herbal products or medicines you've bought without a prescription.

Some medicines must not be taken with Sumatriptan Nasal Spray and others may cause adverse effects if they're taken with Sumatriptan Nasal Spray. **You must tell your doctor if you are taking:**

- **ergotamine** also used to treat **migraine**, or similar medicines such as methysergide (see Section 2). Don't use Sumatriptan Nasal Spray at the same time as these medicines. Stop taking these medicines at least 24 hours before using Sumatriptan Nasal Spray. Don't take any medicines which contain ergotamine or compounds similar to ergotamine again for at least 6 hours after using Sumatriptan Nasal Spray.
- **other triptans/5-HT₁ receptor agonists** (such as naratriptan, rizatriptan, zolmitriptan), also used to treat **migraine**, (see Section 2). Don't use Sumatriptan Nasal Spray at the same time as these medicines. Stop taking these medicines at least 24 hours before using Sumatriptan Nasal Spray. Don't take another triptan/5-HT₁ receptor agonist again for at least 24 hours after using Sumatriptan Nasal Spray.
- **SSRIs** (*Selective Serotonin Reuptake Inhibitors*) or **SNRIs** (*Serotonin Noradrenaline Reuptake Inhibitors*) used to treat **depression**. Using Sumatriptan Nasal Spray with these medicines can cause serotonin syndrome (a collection of symptoms which can include restlessness, confusion, sweating, hallucinations, increased reflexes, muscle spasms, shivering, increased heartbeat and shaking). Tell your doctor immediately if you are affected in this way.
- **MAOIs** (*monoamine oxidase inhibitors*) used to treat **depression**. Don't use Sumatriptan Nasal Spray if you have taken these in the last 2 weeks.
- **St John's Wort** (*Hypericum perforatum*). Taking herbal remedies that contain St John's Wort while you are using Sumatriptan Nasal Spray may make side effects more likely.

Pregnancy and breast-feeding

- **If you are pregnant, think you may be pregnant or are planning to have a baby**, ask your doctor for advice before taking this medicine. There is only limited information about the safety of Sumatriptan Nasal Spray for pregnant women, though up until now there is no evidence of any increased risk of birth defects. Your doctor will discuss with you whether or not you should use Sumatriptan Nasal Spray while you are pregnant.
- **Don't breast-feed your baby for 12 hours after using Sumatriptan Nasal Spray.** If you express any breast milk during this time, discard the milk and don't give it to your baby. Some breastfeeding women report breast and/or nipple pain after use of sumatriptan. The pain is usually temporary and disappears in 3 to 12 hours.

Driving and using machines

Either the symptoms of migraine or your medicine may make you drowsy. **If you are affected, don't drive or operate machinery.**

3. How to use Sumatriptan Nasal Spray

Only use Sumatriptan Nasal Spray after your migraine headache begins

Don't use Sumatriptan Nasal Spray to try to prevent an attack.

Always use this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

There's a step-by-step guide to using the spray at the end of this section.

How much to use

Adults aged 18 to 65

- **The usual dose for adults aged 18 to 65** is one Sumatriptan 20 mg Nasal Spray into just one nostril. But one Sumatriptan 10 mg Nasal Spray into one nostril is enough for some people. Don't use more than two sprays in 24 hours.

Adolescents aged 12 to 17

- **The usual dose for adolescents aged 12 to 17** is one Sumatriptan 10 mg Nasal Spray into one nostril.

Children under 12

- **Sumatriptan Nasal Spray is not recommended** for children under 12 years old.

Older people (aged over 65)

- **Sumatriptan Nasal Spray is not recommended** for people aged over 65.

When to use Sumatriptan Nasal Spray

- **It's best to use Sumatriptan Nasal Spray as soon as you feel a migraine coming on**, although it can be used at any time during an attack.

If your symptoms start to come back

- You can use a second Sumatriptan Nasal Spray after 2 hours, but **don't use more than two sprays in 24 hours.**

If the first spray has no effect

- **Don't use a second spray**, or any other Sumatriptan Nasal Spray preparation for the same attack. If Sumatriptan Nasal Spray doesn't give you any relief:
→ **Ask your doctor or pharmacist for advice.**

If you use more Sumatriptan Nasal Spray than you should

- **Don't use more than two Sumatriptan Nasal Sprays in 24 hours.** Using too much Sumatriptan Nasal Spray could make you ill. If you have used more than two sprays in 24 hours:
→ **Contact your doctor for advice.**

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

Step-by-step guide to using your Sumatriptan Nasal Spray


Don't open a blister until you are ready to use a spray.

Each spray is sealed in a blister to keep it clean and safe. If you carry a spray without a blister, or in an open blister, it may not work properly when you need it.


Each spray contains just one dose of Sumatriptan Nasal Spray.

Don't press the plunger too soon or you will lose the dose.


Sumatriptan Nasal Spray has three parts:



How to use the nasal spray



1. **Remove the nasal spray** from the blister packaging just before you want to use it.
2. Get into a comfortable position. You may prefer to sit down.
3. **Blow your nose** if it feels blocked, or if you have a cold.
4. **Hold the nasal spray** gently with your fingers and thumb, as shown in picture **A**. **Don't press the blue plunger yet.**



5. **Block one nostril** by pressing a finger firmly on the side of your nose. It doesn't matter which nostril you choose.
6. **Put the nozzle of the nasal spray into the other nostril**, as far as feels comfortable - about 1 cm or 1/2 inch (picture **B**). **Breathe out gently through your mouth.** Hold your head upright and close your mouth.



7. **Start to breathe in gently through your nose.** As you breathe in: **Press the blue plunger firmly with your thumb.** The plunger may feel a bit stiff and you may hear it click. Keep breathing in while spraying (picture **C**).
8. **Remove the spray and remove your finger from the side of your nose.** **Keep your head upright for 10-20 seconds**, breathing gently in through your nose and out through your mouth. This helps the medicine stay in your nose. Your nose may feel wet inside and you may notice a slight taste after using the spray - this is normal and will soon pass.
9. **Your nasal spray is now empty.** Throw it away safely and hygienically.

4. Possible side effects

Like all medicines, this medicine can cause side effects, but not everybody gets them. Some symptoms may be caused by the migraine itself. Adverse events reported in adults have also been observed in adolescents. These include very rare reports of heart attacks.

Allergic reaction: get doctor's help straight away

The following side effects have occurred but their frequency is not known.

- **The signs of allergy include rash, hives (itchy rash); wheezing; swollen eyelids, face or lips; complete collapse.**

If you get any of these symptoms soon after using Sumatriptan Nasal Spray:

→ **Don't use any more. Contact a doctor straight away.**

Very common side effects

(affect more than **1 in 10** people)

- **Unpleasant taste.**

Common side effects

(affect up to **1 in 10** people)

- **Heaviness, pressure, tightness or pain in the chest**, throat or other parts of the body, or feelings of unusual sensations, including numbness, tingling and warmth or cold. These effects may be intense but generally pass quickly.

If these effects continue or become severe (especially the chest pain):

→ **Get medical help urgently.** In a very small number of people these symptoms can be caused by a heart attack.

Other common side effects include:

- **Irritation** or burning sensation in the nose or throat; nose bleeds.
- **Feeling sick** (nausea) or being sick (vomiting), although this may be due to the migraine itself.
- **Tiredness** or drowsiness.
- **Dizziness**, feeling weak, or getting hot flushes.
- **Temporary increase in blood pressure.**
- **Shortness of breath.**
- **Aching muscles.**

Rare side effects

(affect up to **1 in 1,000** people)

- Breast pain.

Very rare side effects

(affect up to **1 in 10,000** people)

- Liver function changes. If you have a blood test to check your liver function, tell your doctor or nurse that you are using Sumatriptan Nasal Spray.

Some patients may have the following side effects but it is not known how often they occur

- Seizures/fits, tremors, short-lived/temporary muscle spasm (which may affect jaw movement), neck stiffness.
- Visual disturbances such as flickering, reduced vision, double vision, loss of vision and in some cases even permanent defects (although these may be due to the migraine attack itself).
- Heart problems, where your heart beat may go faster, slower or change rhythm, chest pains (*angina*) or heart attack.
- Pale blue-tinged skin and/or pain in your fingers, toes, ears, nose or jaw in response to cold or stress (*Raynaud's phenomenon*).
- Feeling faint (blood pressure may go down).
- Pain in the lower left side of the stomach and bloody diarrhoea (*ischaemic colitis*).
- Diarrhoea.
- If you had a recent injury or if you have inflammation (like rheumatism or inflammation of the colon) you may experience pain or pain worsening at the site of injury or inflammation.
- Pain in the joints.
- Feeling anxious.
- Difficulty swallowing.
- Excessive sweating.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at Website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects, you can help provide more information on the safety of this medicine.

5. How to store Sumatriptan Nasal Spray

- **Keep this medicine out of the sight and reach of children.**
- Do not store above 30°C. Do not freeze.
- Store in the original container in order to protect from light.
- Don't use this medicine after the expiry date shown on the carton. The expiry date refers to the last day of that month.
- Don't throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.
- If your nasal spray fails to work properly or shows any signs of deterioration, you should seek the advice of your pharmacist.

6. Contents of the pack and other information

What Sumatriptan Nasal Spray contains

The active substance is sumatriptan.

Each pre-filled nasal spray device contains 10 mg sumatriptan in 0.1 ml of aqueous solution.

The other ingredients are potassium dihydrogen phosphate, dibasic sodium phosphate anhydrous, sulphuric acid, sodium hydroxide and purified water.

What Sumatriptan Nasal Spray looks like and contents of the pack

This medicine is provided in a pre-filled nasal spray device and contains sumatriptan in 0.1 ml of a yellow-coloured solution.

Sumatriptan Nasal Spray is packed in a box containing the nasal sprays individually sealed in blisters. They are available in packs of 2 and 6 pre-filled nasal spray devices. Each nasal spray contains one dose of Sumatriptan Nasal Spray.

PL: 15814/2012

POM

Manufactured by GlaxoSmithKline Manufacturing S.p.A., Strada Provinciale Asolana, 90, 43056, San Polo di Torriale, Parma, Italy.

Procured from within the EU and repackaged by the Product Licence holder: O.P.D. Laboratories Ltd., Unit 6 Colonial Way, Watford, Herts WD24 4PR.

Leaflet revision and issue date (Ref.) 02.12.2025.

To request a copy of this leaflet in Braille, large print or audio please call 01923 332 796.